



Relations between slices and quotients
of the algebraic cobordism spectrum

Markus Spitzweck

Preprint Nr. 36/2008

RELATIONS BETWEEN SLICES AND QUOTIENTS OF THE ALGEBRAIC COBORDISM SPECTRUM

MARKUS SPITZWECK

ABSTRACT. We prove a relative statement about the slices of the algebraic cobordism spectrum. If the map from MGL to a certain quotient of MGL introduced by Hopkins and Morel is the map to the zero-slice then a relative version of Voevodsky's conjecture on the slices of MGL holds true. We outline the picture for K -theory and rational slices.

CONTENTS

1. Introduction	1
2. Conventions	3
3. Effectivity of MGL	3
4. Quotients and slices of MGL	5
5. K -theory	10
6. Rational slices	12
References	14

1. INTRODUCTION

In this paper we discuss certain aspects of the slice filtration of the algebraic cobordism spectrum MGL . The slice filtration was introduced in [20]. It is a filtration on any motivic spectrum which can be thought of as an analogue of the Postnikov tower of a topological spectrum. We discuss the relation between the slice conjecture on MGL [20, Conjecture (5)] and quotients of MGL . The conjecture describes the slices in terms of the motivic Eilenberg MacLane spectrum and the topological coefficients MU_* . The quotients are defined using the classifying map $L_* \rightarrow MGL_{**}$ of the formal group law induced by the canonical orientation of MGL and canonical generators x_i of L_* . Here L_* denotes the graded Lazard ring. In topology it is well known that the quotient $MU/(x_1, x_2, \dots)MU$ is isomorphic to the Eilenberg MacLane spectrum on the integers. This follows essentially from Quillen's theorem $L_* \cong MU_*$ and the particular structure of L_* . In motivic homotopy theory no direct analog of this argument seems to work. In particular, the filtration on MGL obtained by dividing out the x_i (more precisely ideals of L_* consisting of elements of

Date: May 27, 2009.

degree greater than a given bound) is not directly reflected on the homotopy groups of \mathbf{MGL} , instead this filtration is conjecturally the slice filtration of \mathbf{MGL} . We show that if this holds on the zeroth level then it holds in all levels. The statement is of a purely homotopy theoretic nature and could be formulated in any context of highly structured ring spectra.

Over perfect fields the zero slice of the sphere spectrum is known [19], [9]. An effectivity result for \mathbf{MGL} implies that \mathbf{MGL} has the same zero slice, see Corollary (3.3).

Our main statement implies that if the quotient of \mathbf{MGL} by all the x_i coincides with this zero slice then [20, Conjecture (5)] holds, see Corollary (4.7). As a consequence, under the same assumption, slices of all Landweber exact spectra are given in terms of the Landweber coefficients and the motivic Eilenberg MacLane spectrum, see [15].

The author was inspired by the course notes [7].

Here is an overview of the sections. In the first section we show effectivity and cellularity results for the algebraic cobordism spectrum. Theorem (3.1) states that the cofiber of the unit map of \mathbf{MGL} lies in a certain subcategory of the stable motivic homotopy category spanned by positive Tate-spheres. As a corollary we obtain the effectivity of \mathbf{MGL} and a proof of the observation in [20] that the zero slices of the sphere spectrum and \mathbf{MGL} coincide. These results can be viewed as refinements to the cellularity results of [1]. The effectivity of \mathbf{MGL} was implicitly assumed in [15].

The second paragraph deals with our main observation that if the Hopkins-Morel quotient of \mathbf{MGL} is the zero slice of \mathbf{MGL} then a relative version of Voevodsky's conjecture [20] on slices holds true, Theorem (4.6). This is closely related to the work of Hopkins and Morel on the spectral sequence for \mathbf{MGL} in terms of motivic cohomology, see [8]. In particular it is an unpublished result due to Hopkins and Morel that over fields of characteristic 0 the quotient of \mathbf{MGL} by the x_i is isomorphic to the motivic Eilenberg MacLane spectrum, [7]. It thus follows from Corollary (4.7) that [20, Conjecture (5)] holds over fields of characteristic 0 assuming the Hopkins-Morel isomorphism. As above we get under the same assumption the structure of the slices of Landweber spectra.

We note that the idea of using filtrations of the ideal of the Lazard ring spanned by elements of positive degree in a homotopy way goes back to [7].

The third paragraph covers the relationship of quotients of \mathbf{MGL} to the algebraic K -theory spectrum. The main statement is also contained in [7].

In the last paragraph we consider rational slices. Due to the rational splitting of \mathbf{MGL} this simplifies to understanding the rational Landweber theory \mathbf{LQ} for the additive formal group law over the rationals, see [11]. We obtain these assertions over regular base schemes by comparing the Landweber decomposition of $\mathbf{KGL}_{\mathbb{Q}}$ and a decomposition obtained in [13].

Acknowledgements. I thank the lecturers of [7]. I thank Niko Naumann, Paul Arne Østvær, Ivan Panin and Oliver Röndigs for inspiring communications. I thank Ulrich Bunke and Ansgar Schneider for helpful comments.

2. CONVENTIONS

Throughout the article we work over a Noetherian base scheme of finite Krull dimension S . The stable motivic homotopy category over S is denoted by $\mathbf{SH}(S)$. The standard spheres are denoted by $S^{p,q} \cong S_s^{\wedge(p-q)} \wedge \mathbb{G}_m^{\wedge q}$. The category of smooth schemes over S is denoted by \mathbf{Sm}/S . The tensor unit of $\mathbf{SH}(S)$, i.e. the sphere spectrum, is denoted by $\mathbf{1}$.

The full subcategory of $\mathbf{SH}(S)$ of effective spectra is denoted by $\mathbf{SH}(S)^{eff}$. It is the full localizing triangulated subcategory generated by all $\Sigma_T^\infty X_+$, $X \in \mathbf{Sm}/S$.

The i -th slice of a motivic spectrum \mathbf{E} is denoted $s_i(\mathbf{E})$, see [20].

Throughout the text we will use the language of model categories. An injective model structure will be a model structure e.g. on a diagram category where weak equivalences and cofibrations are understood objectwise. Dually the projective model structure has objectwise fibrations. For homotopy colimits and limits we refer to [3]. For symmetric monoidal model categories we refer to [5].

3. EFFECTIVITY OF MGL

Let $\mathbf{SH}(S)_{\mathcal{T}}$ be the full localizing triangulated subcategory of $\mathbf{SH}(S)$ spanned by $\{S^{p,q} | p, q \in \mathbf{Z}\}$, see [11]. We let $\mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$ the full localizing triangulated subcategory of $\mathbf{SH}(S)$ spanned by $\{S^{p,q} | p \in \mathbf{Z}, q \geq 0\}$.

Theorem 3.1. *The cofiber of the unit map $\mathbf{1} \rightarrow \mathbf{MGL}$ is contained in $\Sigma_T \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$.*

Corollary 3.2. *We have $\mathbf{MGL} \in \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$. In particular \mathbf{MGL} is an effective spectrum.*

Proof. Follows from $\mathbf{1} \in \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$. □

Corollary 3.3. *The unit map $\mathbf{1} \rightarrow \mathbf{MGL}$ induces an isomorphism*

$$s_0(\mathbf{1}) \xrightarrow{\cong} s_0(\mathbf{MGL}).$$

Proof. The functor s_0 is triangulated and $s_0(X) = 0$ for any $X \in \Sigma_T \mathbf{SH}(S)^{eff}$. □

We start with preparations for the proof of theorem (3.1).

Lemma 3.4. *Let r be an integer and let*

$$S^{2r,r} \rightarrow X \rightarrow Y \rightarrow S^{2r,r}[1]$$

and

$$X \rightarrow Z \rightarrow W \rightarrow X[1]$$

be two triangles in $\mathbf{SH}(S)$. Suppose $Y, W \in \Sigma_T^{r+1} \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$. Then the cofiber of $S^{2r,r} \rightarrow Z$ is in $\Sigma_T^{r+1} \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$.

Proof. The cofiber of $S^{2r,r} \rightarrow Z$ is an extension of W by Y . □

Lemma 3.5. *Let $i: Z \rightarrow X$ be a closed immersion in \mathbf{Sm}/S and \mathcal{E} a vector bundle over X . Let $U = X \setminus Z$, and denote the restrictions of \mathcal{E} to Z and U by \mathcal{E}_Z and \mathcal{E}_U , respectively. Let \mathcal{N} be the normal bundle of i . Then the cofiber of $\mathrm{Th}(\mathcal{E}_U) \rightarrow \mathrm{Th}(\mathcal{E})$ is canonically isomorphic to $\mathrm{Th}(\mathcal{E}_Z \oplus \mathcal{N})$ in the \mathbf{A}^1 -homotopy category.*

Proof. This follows from the fact that this cofiber is (as Zariski sheaves) isomorphic to $\mathcal{E}/(\mathcal{E} \setminus j(Z))$, j the composition of i followed by the zero section of \mathcal{E} , and that the normal bundle of j is $\mathcal{E}_Z \oplus \mathcal{N}$. \square

We first quote some facts from [11] about finite Grassmannians. We let $\mathbb{G}(n, d)$ be the scheme parameterizing locally free quotients of rank d of the trivial bundle of rank n . There is a universal short exact sequence

$$(1) \quad 0 \longrightarrow \mathcal{K}_{n,d} \longrightarrow \mathcal{O}_{\mathbb{G}(n,d)}^n \longrightarrow \mathcal{Q}_{n,d} \longrightarrow 0$$

of vector bundles on $\mathbb{G}(n, d)$, and letting $\mathcal{K}'_{n,d}$ the dual of $\mathcal{K}_{n,d}$, the tangent bundle is given by

$$(2) \quad \mathcal{T}_{\mathbb{G}(n,d)} \cong \mathcal{Q}_{n,d} \otimes \mathcal{K}'_{n,d}.$$

The map

$$i : \mathbb{G}(n, d) \hookrightarrow \mathbb{G}(n+1, d+1)$$

classifying $\mathcal{K}_{n,d} \subseteq \mathcal{O}_{\mathbb{G}(n,d)}^n \hookrightarrow \mathcal{O}_{\mathbb{G}(n,d)}^{n+1}$ is a closed immersion. From (2) it follows that the normal bundle $\mathcal{N}(i)$ of i identifies with $\mathcal{K}'_{n,d}$. Next consider the composition on $\mathbb{G}(n+1, d+1)$

$$\alpha : \mathcal{O}_{\mathbb{G}(n+1,d+1)}^n \hookrightarrow \mathcal{O}_{\mathbb{G}(n+1,d+1)}^{n+1} \longrightarrow \mathcal{Q}_{n+1,d+1}$$

for the inclusion into the first n factors. The complement of the support of $\text{coker}(\alpha)$ is an open subscheme $U \subseteq \mathbb{G}(n+1, d+1)$ and there is a map $\pi : U \rightarrow \mathbb{G}(n, d+1)$ classifying $\alpha|_U$. It is easy to see that π identifies with $\mathcal{Q}_{n,d+1} \rightarrow \mathbb{G}(n, d+1)$. An argument with geometric points reveals that $U = \mathbb{G}(n+1, d+1) \setminus i(\mathbb{G}(n, d))$. Moreover the natural map $\iota : \mathbb{G}(n, d+1) \rightarrow \mathbb{G}(n+1, d+1)$ classifying the subbundle $\mathcal{K}_{n,d+1} \oplus \mathcal{O} \subset \mathcal{O}^{n+1}$ is the zero section $\mathbb{G}(n, d+1) \rightarrow \mathcal{Q}_{n,d+1}$ followed by the inclusion $\mathcal{Q}_{n,d+1} \cong U \rightarrow \mathbb{G}(n+1, d+1)$.

We summarize the above with a diagram:

$$(3) \quad \mathbb{G}(n, d) \xrightarrow{i} \mathbb{G}(n+1, d+1) \longleftarrow U \xrightarrow[\pi]{\leftarrow} \mathbb{G}(n, d+1).$$

We note that compositions of the morphisms ι yields a map $\bar{\iota} : \text{pt} \cong \mathbb{G}(d, d) \rightarrow \mathbb{G}(n, d)$ which we consider as the natural pointing of $\mathbb{G}(n, d)$. Note that the unit of MGL is induced via these maps.

Proposition 3.6. *Let \mathcal{E} be a vector bundle of rank r over $\mathbb{G}(n, d)$ which is a finite sum of copies of $\mathcal{K}_{n,d}$, $\mathcal{K}'_{n,d}$ and \mathcal{O} . Then $\bar{\iota}^* \mathcal{E}$ is canonically trivialized. Furthermore the cofiber of the map of suspension spectra of Thom spaces $S^{2r,r} \rightarrow \Sigma^\infty \text{Th}(\mathcal{E})$ lies in $\Sigma_T^{r+1} \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$.*

Proof. We prove the statement by induction on n . It clearly holds for $n = 0$. Suppose $n \geq 0$ and assume the statement holds for n . The statement holds for $\mathbb{G}(n+1, 0)$ and $\mathbb{G}(n+1, n+1)$. Let $0 \leq d < n$. We prove the statement for $\mathbb{G}(n+1, d+1)$. Let \mathcal{E} be a vector bundle on $\mathbb{G}(n+1, d+1)$ of the considered type. It is canonically trivialized over the pointing. We consider the diagram (3). By lemma (3.5) we get an induced exact triangle

$$(4) \quad \Sigma^\infty \text{Th}(\mathcal{E}_U) \longrightarrow \Sigma^\infty \text{Th}(\mathcal{E}) \longrightarrow \Sigma^\infty \text{Th}(\mathcal{E}_{\mathbb{G}(n,d)} \oplus \mathcal{K}'_{n,d}) \longrightarrow \Sigma^\infty \text{Th}(\mathcal{E}_U)[1].$$

We note $\mathcal{K}_{n+1,d+1}|_{\mathbb{G}(n,d)} \cong \mathcal{K}_{n,d}$, hence $\mathcal{E}_{\mathbb{G}(n,d)}$ and $\mathcal{E}_{\mathbb{G}(n,d)} \oplus \mathcal{K}'_{n,d}$ are vector bundles on $\mathbb{G}(n,d)$ of the considered type.

Since $\iota^* \mathcal{K}_{n+1,d+1} \cong \mathcal{K}(n,d+1) \oplus \mathcal{O}$ there is an induced map $\mathrm{Th}(\mathcal{K}_{n,d+1} \oplus \mathcal{O}) \rightarrow \mathrm{Th}(\mathcal{K}_{n+1,d+1})$ which factors through $\mathrm{Th}(\mathcal{K}_{n+1,d+1}|_U)$. Note $\mathrm{Th}(\mathcal{K}_{n,d+1} \oplus \mathcal{O}) \rightarrow \mathrm{Th}(\mathcal{K}_{n+1,d+1}|_U)$ and more generally $\mathrm{Th}(\iota^* \mathcal{E}) \rightarrow \mathrm{Th}(\mathcal{E}_U)$ are motivic weak equivalences: we cover U by the opens for which α restricted to a fixed subset of the summands of \mathcal{O}^n of size $d+1$ surjects onto $\mathcal{Q}_{n+1,d+1}$. Those opens are pullbacks from $\mathbb{G}(n,d+1)$. We claim on such opens the situation trivializes completely. Without loss of generality we can assume the first $d+1$ summands of \mathcal{Q}^n surject onto $\mathcal{Q}_{n+1,d+1}$. Then $\mathcal{K}_{n+1,d+1}$ restricted to this open trivializes by projecting to the last $n-d$ summands of \mathcal{O}^{n+1} . This trivialization is compatible with the one over the corresponding open $V \subset \mathbb{G}(n,d+1)$. Thus $((\iota^* \mathcal{E})|_V)^\circ \rightarrow (\mathcal{E}|_{\pi^{-1}(V)})^\circ$ is a motivic weak equivalence, and the same holds for the map $(\iota^* \mathcal{E})^\circ \rightarrow \mathcal{E}_U^\circ$ by a Mayer-Vietoris argument. This shows the claim that the map of Thom spaces is also a motivic weak equivalence.

We can thus rewrite the sequence (4) as

$$(5) \quad \Sigma^\infty \mathrm{Th}(\iota^* \mathcal{E}) \longrightarrow \Sigma^\infty \mathrm{Th}(\mathcal{E}) \longrightarrow \Sigma^\infty \mathrm{Th}(\mathcal{E}_{\mathbb{G}(n,d)} \oplus \mathcal{K}'_{n,d}) \longrightarrow \Sigma^\infty \mathrm{Th}(\iota^* \mathcal{E})[1]$$

and use the fact that $\iota^* \mathcal{E}$ is of the type considered for $\mathbb{G}(n,d+1)$. By induction hypothesis the cofiber of $S^{2r,r} \cong \Sigma^\infty \mathrm{Th}(\bar{\iota}^* \iota^* \mathcal{E}) \rightarrow \Sigma^\infty \mathrm{Th}(\iota^* \mathcal{E})$ lies in $\Sigma_T^{r+1} \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$. Moreover again by induction hypothesis $\Sigma^\infty \mathrm{Th}(\mathcal{E}_{\mathbb{G}(n,d)} \oplus \mathcal{K}'_{n,d}) \in \Sigma_T^{r+j} \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$ with $j = n-d > 0$. Now the statement follows from lemma (3.4). \square

Proof of Theorem (3.1). We let $\mathbf{BGL}_n = \mathrm{colim}_d \mathbb{G}(n+d,d)$, $\xi_n = \mathrm{colim}_d \mathcal{K}_{n+d,d}$ the universal vector bundle. Then

$$\mathbf{MGL} = \mathrm{hocolim}_n \Sigma^{-2n,-n} \Sigma^\infty \mathrm{Th}(\xi_n) \cong \mathrm{hocolim}_{n,d} \Sigma^{-2n,-n} \Sigma^\infty \mathrm{Th}(\mathcal{K}_{n+d,d}).$$

The unit $\mathbf{1} \rightarrow \mathbf{MGL}$ is induced via the maps

$$\Sigma^{-2n,-n} \Sigma^\infty \mathrm{Th}(\bar{\iota}^* \mathcal{K}_{n+d,d}) \rightarrow \Sigma^{-2n,-n} \Sigma^\infty \mathrm{Th}(\mathcal{K}_{n+d,d}).$$

By proposition (4) the cofibers of these maps are in $\Sigma_T \mathbf{SH}(S)_{\mathcal{T}_{\geq 0}}$. Since cofiber sequences are compatible with homotopy colimits the claim follows. \square

4. QUOTIENTS AND SLICES OF MGL

Let \mathbf{MGL} denote a fibrant and cofibrant model as commutative \mathbb{S} -algebra of the algebraic cobordism spectrum. We work in the simplicial version of the \mathbb{S} -modules of [2], see [14]. In particular \mathbf{MGL} is fibrant and cofibrant as (symmetric) T -spectrum. We let $\mathrm{Mod}(\mathbf{MGL})$ be the symmetric monoidal category with weak unit of \mathbf{MGL} -modules. The homotopy category of $\mathrm{Mod}(\mathbf{MGL})$ is denoted by $\mathcal{D}_{\mathbf{MGL}}$. It is a closed tensor triangulated category.

We denote $\varphi: \mathbf{MU}_* \rightarrow \mathbf{MGL}_*$ the canonical map and fix an isomorphism $\mathbf{MU}_* \cong \mathbf{Z}[x_1, x_2, \dots]$, $|x_i| = i$ (we divide the usual topological grading by 2).

As in topology we can form the quotient $\mathbf{Q} := \mathbf{MGL}/(x_1, x_2, \dots)\mathbf{MGL}$ by taking iterated cofibers of multiplications by the x_i in $\mathcal{D}_{\mathbf{MGL}}$.

This quotient is well-defined up to isomorphism.

We give a construction of a more explicit model of this quotient:

We pick once and for all the following data:

- (1) $S^{2i,i} = T^{\wedge i}$, $i > 0$, a cofibrant model of the $(2i, i)$ -sphere in T -spectra, and denote the corresponding cofibrant sphere $S^{2i,i} \wedge \mathbf{MGL}$ by $\Sigma^{2i,i}\mathbf{MGL}$.
- (2) a map $\Sigma^{2i,i}\mathbf{MGL} \rightarrow \mathbf{MGL}$ representing the element $\varphi_*(x_i) \in \mathbf{MGL}_{2i,i}$, also denoted by x_i .

We let I be the following category: the objects are (commutative) monomials in the x_i , there is a unique map from M to N if N divides M ; the monomial 1 is allowed. We let I° be the full subcategory consisting of all non-constant monomials. The subcategory $I_{\leq 1}$ consists of monomials containing each x_i with multiplicity at most 1, $I_{\leq 1}^\circ$ is the same category with the constant monomial removed.

We let $\underline{1}$ be the category whose diagrams are morphisms, i.e. two objects 0 and 1 and one non identity map. We let $\underline{1}^n \subset I_{\leq 1}$ be the inclusion of the monomials containing only the x_i , $i \leq n$. Via these inclusions $I_{\leq 1}$ is the union of all the $\underline{1}^n$.

We are going to define the following I -diagram of \mathbf{MGL} -modules: a monomial $x_1^{k_1} \dots x_n^{k_n}$, $k_i \geq 0$, is sent to $(\Sigma^{2,1}\mathbf{MGL})^{\wedge_{\mathbf{MGL}} k_1} \wedge_{\mathbf{MGL}} \dots \wedge_{\mathbf{MGL}} (\Sigma^{2n,n}\mathbf{MGL})^{\wedge_{\mathbf{MGL}} k_n}$. The morphisms will be given by multiplications with the x_i , i.e. iterations of the morphisms x_i . We have to be careful since in general the two possible maps $\Sigma^{2i,i}\mathbf{MGL} \wedge_{\mathbf{MGL}} \Sigma^{2i,i}\mathbf{MGL} \rightarrow \Sigma^{2i,i}\mathbf{MGL}$ given by applying the map x_i either on the left or on the right and then composing with a unit morphism (note \mathbf{MGL} only serves as a weak unit) do not coincide in general. Therefore we make the convention that for a map $M \rightarrow N$ in I , $M = x_1^{k_1} \dots x_n^{k_n}$, $N = x_1^{l_1} \dots x_n^{l_n}$, $l_i \leq k_i$, we insert for any $1 \leq i \leq n$ the map $(\Sigma^{2i,i}\mathbf{MGL})^{\wedge_{\mathbf{MGL}} k_i} \rightarrow (\Sigma^{2i,i}\mathbf{MGL})^{\wedge_{\mathbf{MGL}} l_i}$ which applies x_i on the $k_i - l_i$ right most tensor factors of the source followed by appropriate unit maps.

We end up with a diagram of \mathbf{MGL} -modules, denoted \mathbf{D} .

We need the following lemma in which we denote by hofib the homotopy fiber of a map between pointed simplicial sets.

Lemma 4.1. *Let*

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ Z & \longrightarrow & W \end{array}$$

be a diagram of pointed simplicial sets. Let P be the homotopy pullback of the right lower triangle of the square. Then the homotopy fiber of the natural map $X \rightarrow P$ is naturally equivalent to $\text{hofib}(\text{hofib}(X \rightarrow Z) \rightarrow \text{hofib}(Y \rightarrow W))$.

Proof. Replacing the diagram with an injectively fibrant diagram the statement follows from the corresponding strict statement. \square

Let $\mathbf{D}_{\leq 1}$ be the restriction of \mathbf{D} to $I_{\leq 1}$ and $\mathbf{D}_{\leq 1}^\circ$ the one to $I_{\leq 1}^\circ$.

We have the following observations. We use the notion of total cofiber of a diagram with respect to a subdiagram, see [6]. This is defined to be the cofiber of the natural map from the homotopy colimit of the subdiagram to the homotopy colimit of the total diagram. Usually this can be viewed as the total object corresponding to a diagram of a certain shape, e.g. of a cubical diagram. For a functor to be homotopy right cofinal see [3, Definition 19.6.1].

- Lemma 4.2.** (1) *The inclusion $I_{\leq 1}^{\circ} \rightarrow I^{\circ}$ is homotopy right cofinal.*
(2) *The total cofiber of the diagram $D_{\leq 1}$ with respect to the inclusion $D_{\leq 1}^{\circ} \rightarrow D_{\leq 1}$, i.e. the cofiber of the map $\text{hocolim} D_{\leq 1}^{\circ} \rightarrow D(1) = \text{MGL}$, is isomorphic to \mathbb{Q} in \mathcal{D}_{MGL} .*

Proof. (1): Let the inclusion be denoted by j . We have to show that for any object $o \in I^{\circ}$ the under category $o \backslash j$ is contractible. But this under category has the initial object $(o', o \rightarrow o')$, where o' contains x_i with multiplicity 1 if $x_i | o$, otherwise it contains it with multiplicity 0.

(2): We first show the analogous statement for finitely many x_i : the total cofiber of $D|_{\underline{1}^n}$ with respect to the inclusion $(\underline{1}^n)^{\circ} \subset \underline{1}^n$ is equivalent to the quotient $\text{MGL}/(x_1, \dots, x_n)\text{MGL}$. This is proved by induction on n using the following statement (with $J = \underline{1}^n$ for the induction step $n \mapsto n + 1$):

Let J° be a small category and let J be the same category added a terminal object (so if J had already a terminal object this object will no longer be terminal). Let \mathcal{C} be a pointed model category and $G: J \times \underline{1} \rightarrow \mathcal{C}$ be a functor. Let $(J \times \underline{1})^{\circ}$ be again the category obtained by removing the terminal object. Then the total cofiber of G with respect to the inclusion $(J \times \underline{1})^{\circ} \rightarrow J \times \underline{1}$ can be computed as follows: it is the cofiber of the map $\text{Totcof}(G|_{J \times \{0\}}) \rightarrow \text{Totcof}(G|_{J \times \{1\}})$, Totcof denoting the total cofiber with respect to the inclusions of the respective subcategories obtained by removing the terminal object.

By considering pointed mapping spaces this statement reduces to the dual statement for pointed simplicial sets. Here to compute homotopy limits we can use the injective model structure on diagram categories. Now we switch and let J be the opposite of the original J , similarly with J° . The category $(J \times \underline{1})^{\circ}$ this time denotes $J \times \underline{1}$ with the initial object removed. Let $G: J \times \underline{1} \rightarrow \mathbf{sSet}_{\bullet}$ be a functor. Fix an injectively fibrant replacement $G \rightarrow R$. We note that $R|_{(J \times \underline{1})^{\circ}}$, $R|_{J^{\circ} \times \{i\}}$, $i = 0, 1$, are again injectively fibrant: the respective restriction functors are right Quillen functors since the corresponding adjoints preserve objectwise cofibrations.

Let i be the initial object of J . We replace the diagram R by the square

$$\begin{array}{ccc} R(i, 0) & \longrightarrow & \lim R|_{J^{\circ} \times \{0\}} \\ \downarrow & & \downarrow \\ R(i, 1) & \longrightarrow & \lim R|_{J^{\circ} \times \{1\}} \end{array}$$

This yields again an injectively fibrant diagram, the limit of the lower right triangle gives the homotopy limit of $G|_{(J \times \underline{1})^{\circ}}$, whence the statement follows from lemma (4.1).

We are left to prove the statement for infinitely many x_i . The restriction functors from $I_{\leq 1}^{\circ}$ -diagrams to $(\underline{1}^n)^{\circ}$ -diagrams preserve projectively cofibrant diagrams, whence $\text{hocolim} D_{\leq 1}^{\circ} \simeq \text{hocolim}_n (\text{hocolim} D|_{(\underline{1}^n)^{\circ}})$. This shows the claim. \square

For any $n > 0$ let $I_{\text{deg} \geq n}$ be the subcategory of I of monomials of degree $\geq n$, where the degree of a monomial $x_1^{k_1} \cdots x_n^{k_n}$ is $\sum_{i=1}^n i \cdot k_i$. Moreover for a monomial M let $I_{\geq M}$ be the subcategory of all monomials which are divisible by M . We also let $D_{\text{deg} \geq n} = D|_{I_{\text{deg} \geq n}}$ and $D_{\geq M} = D|_{I_{\geq M}}$.

Proposition 4.3. *Let $F: I_{\text{deg} \geq n} \rightarrow \mathcal{C}$ be a diagram in a cofibrantly generated model category \mathcal{C} such that for any monomial M of degree n the natural map $\text{hocolim} F|_{I_{\geq M}} \rightarrow F(M)$ is an equivalence. Then the natural map*

$$\text{hocolim}(F|_{I_{\text{deg} \geq n+1}}) \rightarrow \text{hocolim} F$$

is an equivalence.

Proof. Let $Q \rightarrow F$ be a cofibrant replacement of F for the projective model structure on $\mathcal{C}^{I_{\text{deg} \geq n}}$. Let M be a monomial of degree n . We claim $Q|_{I_{\geq n}}$ is still cofibrant: in fact the right adjoint r to the restriction functor preserves objectwise fibrations: for $o \in I_{\geq M}$ we have $r(H)(o) = H(o)$, for $o \notin I_{\geq M}$ we have $r(H)(o) = \text{pt}$.

Thus if we define Q' by replacing for any M with $\text{deg}(M) = n$ the object $Q(M)$ with $\text{colim} Q|_{I_{\geq M}}$ and leaving the other entries unchanged we do not change the weak homotopy type of Q . As above $Q'|_{I_{\text{deg} \geq n+1}}$ is cofibrant, moreover Q' is cofibrant itself since for any $B \in \mathcal{C}^{I_{\text{deg} \geq n}}$ we have $\text{Hom}(Q'|_{I_{\text{deg} \geq n+1}}, B|_{I_{\text{deg} \geq n+1}}) \cong \text{Hom}(Q', B)$. Thus $\text{hocolim} Q'|_{I_{\text{deg} \geq n+1}} \simeq \text{colim} Q'|_{I_{\text{deg} \geq n+1}} \cong \text{colim} Q' \simeq \text{hocolim} Q' \simeq \text{hocolim} Q$, which finishes the proof. \square

For the proof of the next statement we use the notion of a left Quillen presheaf on ω , where ω is the first infinite ordinal. In this case such a presheaf is given by a model category \mathcal{C}_n for each natural number n and left Quillen functors $f_n: \mathcal{C}_{n+1} \rightarrow \mathcal{C}_n$ for each $n \geq 0$. A section consists of objects $X_n \in \mathcal{C}_n$ for each $n \geq 0$ and maps $f_n(X_{n+1}) \rightarrow X_n$, $n \geq 0$. It is called homotopy cartesian if the maps $(\mathbf{L}f_n)(X_{n+1}) \rightarrow X_n$ are isomorphisms in $\text{Ho}(\mathcal{C}_n)$. The category of sections possesses the inverse model structure where weak equivalences and cofibrations are objectwise. We will use the fact that the mapping space out of a homotopy cartesian section X_\bullet into any Y_\bullet is given as the homotopy limit over the individual mapping spaces $\text{map}(X_n, Y_n)$.

Lemma 4.4. *Let $F: \mathcal{C} \leftrightarrow \mathcal{D}: G$ be a Quillen adjunction between stable pointed model categories. Suppose $\mathbf{R}G: \text{Ho}\mathcal{D} \rightarrow \text{Ho}\mathcal{C}$ preserves sums. Then $\mathbf{R}G$ preserves homotopy colimits.*

Proof. We indicate a proof. First note that a homotopy colimit of a functor $F: I \rightarrow \mathcal{D}$, which we suppose to take values in cofibrant objects, can be computed as the homotopy colimit of the simplicial diagram $[n] \mapsto \coprod_{\varphi: [n] \rightarrow I} F(\varphi(0))$, $[n]$ the ordered set $\{0, \dots, n\}$ viewed as a category. By considering mapping spaces this reduces to the statement that mapping spaces in \mathbf{sSet}^I can be computed by homotopy ends.

Since $\mathbf{R}G$ preserves homotopy coproducts by assumption it thus suffices to show that colimits over Δ^{op} are preserved. Let $\alpha: \Delta^{\text{op}} \rightarrow \mathcal{D}$ be a functor. We claim $\text{hocolim}(\alpha) \simeq \text{hocolim}_n \text{hocolim}(\alpha|_{\Delta_{\leq n}^{\text{op}}})$. This proves the above since $\mathbf{R}G$ preserves finite homotopy colimits since we are dealing with a stable situation and sequential homotopy colimits over ω since $\mathbf{R}G$ preserves sums. Using mapping spaces we reduce the statement about α to the dual statement in simplicial sets. Now we observe that $\text{Ho}(\mathbf{sSet}^{\Delta})$ is equivalent to homotopy cartesian sections in the homotopy category of the category of sections of the left Quillen presheaf on ω , $n \mapsto \mathbf{sSet}^{\Delta_{\leq n}}$, the categories $\mathbf{sSet}^{\Delta_{\leq n}}$ carrying the injective model structure such that the restriction maps preserve cofibrations. Considering mapping spaces out of constant diagrams shows the claim since a mapping space between homotopy cartesian sections is a homotopy limit over ω of the individual mapping spaces. \square

We now turn to the functors $f_i: \mathbf{SH}(S) \rightarrow \Sigma_T^i \mathbf{SH}(S)^{eff} \subset \mathbf{SH}(S)$ introduced in [20]. These are defined as right adjoints to the inclusions $\Sigma_T^i \mathbf{SH}(S)^{eff} \subset \mathbf{SH}(S)$. These functors can be defined on the level of model categories by using colocalization of model categories, see [12]. In particular it makes sense to ask whether these functors preserve homotopy colimits.

Corollary 4.5. *The functors f_i preserve homotopy colimits.*

Proof. This follows from lemma (4.4) and the fact that the f_i preserve sums [20]. \square

We now can state the main observation of this text.

Theorem 4.6. *Suppose the natural map $\mathbf{MGL} \rightarrow \mathbf{Q}$ is the map from \mathbf{MGL} to its zero-slice. Then [20, Conjecture (5)] holds with $H_{\mathbf{MU}_{2q}}$ replaced by $s_0 \mathbf{MGL} \otimes \mathbf{MU}_{2q}$.*

Note the compatibility with the natural homomorphism $\mathbf{MU}_* \rightarrow \mathbf{MGL}_{**}$ still makes sense.

Proof. Lemma (4.2).(1) and (2) and [3, Theorem 19.6.13] imply that the map from \mathbf{MGL} to the total cofiber of \mathbf{D} with respect to $\mathbf{D}^\circ \rightarrow \mathbf{D}$ is the map to \mathbf{Q} , hence by assumption this is the map to the zero-slice. Thus $\mathrm{hocolim} \mathbf{D}^\circ \rightarrow f_1 \mathbf{MGL}$ is an equivalence. By corollary (4.5) the f_i commute with homotopy colimits. We denote by \mathcal{F}_i functorial versions of the f_i on the level of model categories. Then we can rewrite $f_2 \mathrm{hocolim} \mathbf{D}^\circ \simeq \mathrm{hocolim} \mathcal{F}_2 \mathbf{D}^\circ$. Observe the diagram $\mathcal{F}_2 \mathbf{D}^\circ$ satisfies the assumptions of proposition (4.3) for $n = 1$, hence $f_2 \mathbf{MGL} \simeq \mathrm{hocolim} \mathcal{F}_2 \mathbf{D}^\circ \simeq \mathrm{hocolim} \mathbf{D}_{\mathrm{deg} \geq 2}$.

Increasing the degree of the monomials it follows from proposition (4.3) by induction that $f_n \mathbf{MGL} \simeq \mathrm{hocolim} \mathbf{D}_{\mathrm{deg} \geq n}$. In addition the maps

$$f_{n+1} \mathbf{MGL} \rightarrow f_n \mathbf{MGL}$$

are the naturally induced maps

$$\mathrm{hocolim} \mathbf{D}_{\mathrm{deg} \geq n+1} \rightarrow \mathrm{hocolim} \mathbf{D}_{\mathrm{deg} \geq n}.$$

Thus the $s_n \mathbf{MGL}$ are the cofibers of these maps. We rewrite the source again as $\mathrm{hocolim} \mathcal{F}_{n+1} \mathbf{D}_{\mathrm{deg} \geq n}$. Cofibers commute with homotopy colimits, thus $s_n \mathbf{MGL} \simeq \mathrm{hocolim} \mathrm{cofib}(\mathcal{F}_{n+1} \mathbf{D}_{\mathrm{deg} \geq n} \rightarrow \mathbf{D}_{\mathrm{deg} \geq n})$. The value of this cofiber of diagrams at a monomial of degree $> n$ is zero, at a monomial of degree n we exactly have $\Sigma_T^n s_0 \mathbf{MGL}$. Now it is easy to see that the homotopy colimit of such a diagram is the homotopy coproduct of the entries in degree n , which is $\Sigma_T^n s_0 \mathbf{MGL} \otimes \mathbf{MU}_{2n}$.

The compatibility with the natural homomorphism $\mathbf{MU}_* \rightarrow \mathbf{MGL}_{**}$ follows by the choice of the x_i . \square

Recall the natural orientation $\mathbf{MGL} \rightarrow \mathbf{MZ}$ of the motivic Eilenberg MacLane spectrum. It is additive, in particular the x_i all map to zero in \mathbf{MZ}_{**} . Iteratively we get factorizations $\mathbf{MGL}/(x_1, \dots, x_n) \mathbf{MGL} \rightarrow \mathbf{MZ}$ in $\mathbf{SH}(S)$ in a compatible way, which gives a map $\mathbf{Q} \rightarrow \mathbf{MZ}$.

Corollary 4.7. *Suppose S is the spectrum of a perfect field. If the natural map $\mathbf{Q} \rightarrow \mathbf{MZ}$ is an isomorphism then [20, Conjecture (5)] holds.*

Proof. By [9] and [19] $s_0\mathbf{1} \cong \mathbf{MZ}$. Thus by corollary (3.3) the map $\mathbf{MGL} \rightarrow \mathbf{MZ}$ is the map from \mathbf{MGL} to its zero-slice. The statement follows from theorem (4.6). \square

5. K-THEORY

By the Landweber exactness theorem [11] and [10] the spectrum \mathbf{KGL} is the Landweber spectrum associated to the \mathbf{MU}_* -algebra $x_1^{-1}\mathbf{MU}_*/(x_2, x_3, \dots)\mathbf{MU}_* \cong \mathbf{Z}[u, u^{-1}]$. The latter algebra classifies the multiplicative formal group law $x + y - uxy$.

The orientation $\mathbf{MGL} \rightarrow \mathbf{KGL}$ sends all $x_i \in \mathbf{MGL}_{2i,i}$, $i \geq 2$, to $0 \in \mathbf{KGL}_{2i,i}$. Thus we obtain a factorization $\mathbf{MGL}/(x_2, x_3, \dots)\mathbf{MGL} \rightarrow \mathbf{KGL}$. Since x_1 acts invertibly on \mathbf{KGL} this map further factors as

$$B: x_1^{-1}\mathbf{MGL}/(x_2, x_3, \dots)\mathbf{MGL} \rightarrow \mathbf{KGL}.$$

Lemma 5.1. *Let*

$$0 \rightarrow M_* \xrightarrow{f} N_* \rightarrow Q_* \rightarrow 0$$

be a short exact sequence of evenly graded Landweber exact \mathbf{MU}_ -modules. Let $F: \mathbf{E}_M \rightarrow \mathbf{E}_N$ be a map of spectra in $\mathbf{SH}(\mathbf{Z})_{\mathcal{T}}$ representing the homology transformation given by f via the motivic Landweber exact functor theorem. Then the cofiber of F represents the Landweber theory given by Q_* .*

Proof. We let \mathcal{X} be the stack $[\mathbf{Spec}(\mathbf{MU}_*)/\mathbf{Spec}(\mathbf{MU}_*\mathbf{MU})]$ and \widetilde{M}_* etc. the quasi coherent sheaves on \mathcal{X} obtained from the M_* etc. by pushforward along the map $\mathbf{Spec}(\mathbf{MU}_*) \rightarrow \mathcal{X}$. Then by Landweber exactness the sequence

$$0 \rightarrow \widetilde{M}_* \rightarrow \widetilde{N}_* \rightarrow \widetilde{Q}_* \rightarrow 0$$

is a short exact sequence of flat $\mathcal{O}_{\mathcal{X}}$ -modules. In particular tensoring this sequence with a quasi coherent $\mathcal{O}_{\mathcal{X}}$ -module yields again a short exact sequence. However, the Landweber theorem ([11]) is proved by considering the \mathbf{MGL} -homology of a motivic spectrum as a $(\mathbf{MGL}_*, \mathbf{MGL}_*\mathbf{MGL})$ -comodule and via restriction along a map of Hopf algebroids as $(\mathbf{MU}_*, \mathbf{MU}_*\mathbf{MU})$ -comodule, then tensoring this over $\mathcal{O}_{\mathcal{X}}$ with \widetilde{M}_* and finally pulling back to $\mathbf{Spec}(\mathbf{MU}_*)$. This shows that the sequence of motivic homology theories obtained from the original sequence is short exact. In particular if the first map is represented by any map of motivic spectra then the cofiber will represent the homology theory associated with Q_* . \square

Theorem 5.2. *The map of spectra $B: x_1^{-1}\mathbf{MGL}/(x_2, x_3, \dots)\mathbf{MGL} \rightarrow \mathbf{KGL}$ is an isomorphism.*

Proof. We use that $x_1^{-1}\mathbf{MGL}/(x_2, x_3, \dots)\mathbf{MGL}$ can also be constructed by first inverting x_1 and then quotienting out the x_i , $i \geq 2$. Now observe that all the quotients $x_1^{-1}\mathbf{MU}_*/(x_2, \dots, x_n)\mathbf{MU}_*$ are Landweber exact: they are torsionfree, and for a prime p the element v_1 of $(\mathbf{MU}_*)_{(p)}$ already acts invertibly. Thus the

$$\begin{aligned} 0 \rightarrow \Sigma^{n+1}x_1^{-1}\mathbf{MU}_*/(x_2, \dots, x_n)\mathbf{MU}_* &\rightarrow x_1^{-1}\mathbf{MU}_*/(x_2, \dots, x_n)\mathbf{MU}_* \\ &\rightarrow x_1^{-1}\mathbf{MU}_*/(x_2, \dots, x_{n+1})\mathbf{MU}_* \rightarrow 0 \end{aligned}$$

(Σ refers to a shift of evenly graded groups) are short exact sequences of evenly graded Landweber modules, and lemma (5.1) applies. Thus it follows inductively

that the quotients $x_1^{-1}\mathrm{MGL}/(x_2, \dots, x_n)\mathrm{MGL}$ are the Landweber spectra for the modules $x_1^{-1}\mathrm{MU}_*/(x_2, \dots, x_n)\mathrm{MU}_*$. Passing to the colimit shows the claim. \square

Remark 5.3. Fix a prime p and let $\mathrm{BP}^{\mathrm{Top}}$ be the topological Brown-Peterson spectrum. Let BP be the motivic spectrum on the Landweber coefficients $\mathrm{BP}_*^{\mathrm{Top}}$. It can be seen that this coincides with the definition given in [18] since both definitions give rise to the universal oriented ring cohomology theory on compact objects with p -typical formal group law.

Let $\mathrm{E}(n)^{\mathrm{Top}}$ be the topological spectrum for the coefficients

$$v_n^{-1}\mathrm{BP}_*^{\mathrm{Top}}/(v_{n+1}, v_{n+2}, \dots).$$

The corresponding motivic Landweber spectrum is denoted $\mathrm{E}(n)$. Now the quotients $v_n^{-1}\mathrm{BP}_*^{\mathrm{Top}}/(v_{n+1}, \dots, v_{n+k})\mathrm{BP}_*^{\mathrm{Top}}$ are also Landweber exact. Hence by the same method as for theorem (5.2) we see that there is an isomorphism

$$\mathrm{E}(n) \cong v_n^{-1}\mathrm{BP}/(v_{n+1}, v_{n+2}, \dots)\mathrm{BP},$$

compare with [4].

The map from [15, par. 6] is ill-defined since BP is not a direct summand of $\mathrm{MGL}_{(p)}$ as MGL -module.

Next we analyze the relationship to connective or effective K -theory. A version of that has been introduced in [9].

Since $\mathrm{MGL}/(x_2, \dots)\mathrm{MGL}$ is effective, the canonical map $\mathrm{MGL}/(x_2, \dots)\mathrm{MGL} \rightarrow x_1^{-1}\mathrm{MGL}/(x_2, \dots)\mathrm{MGL} \cong \mathrm{KGL}$ factors as

$$B^{\mathrm{eff}} : \mathrm{MGL}/(x_2, \dots)\mathrm{MGL} \rightarrow f_0\mathrm{KGL}.$$

Proposition 5.4. Suppose the map $\mathrm{MGL} \rightarrow \mathbb{Q}$ is the map from MGL to its zero-slice. Then B^{eff} induces isomorphisms on slices. If S is the spectrum of a perfect field and B^{eff} is an isomorphism, then the map $\mathrm{MGL} \rightarrow \mathbb{Q}$ is the map from MGL to its zero-slice.

Proof. Let us assume $\mathrm{MGL} \rightarrow \mathbb{Q}$ is the map to the zero-slice. Then by theorem (4.6) the assumption (SIMGL) of [15] is fulfilled, hence by [15, corollary 4.2] we have $s_0\mathrm{KGL} \cong s_0\mathrm{MGL}$, and the isomorphism is realized by the map $s_0\mathrm{MGL} \rightarrow s_0f_0\mathrm{KGL}$.

By the periodicity of KGL the map $f_1\mathrm{KGL} \rightarrow f_0\mathrm{KGL}$ is isomorphic to the map $\Sigma_T f_0\mathrm{KGL} \rightarrow f_0\mathrm{KGL}$ given by multiplication by the Bott element $u \in (f_0\mathrm{KGL})_{2,1} = \mathrm{KGL}_{2,1}$. Thus the cofiber of this map is isomorphic to $s_0\mathrm{MGL}$, and the same holds true for the cofiber of the map $\Sigma_T\mathrm{MGL}/(x_2, \dots)\mathrm{MGL} \rightarrow \mathrm{MGL}/(x_2, \dots)\mathrm{MGL}$ given by multiplication by x_1 by the assumption. Iterating this process shows the claim.

Suppose now that S is the spectrum of a perfect field. Then by [9] there are isomorphisms $s_0\mathbf{1} \cong s_0\mathrm{KGL} \cong \mathrm{MZ}$, and by corollary (3.3) we also have $s_0\mathrm{MGL} \cong s_0\mathrm{KGL}$. Now if B^{eff} is an isomorphism, then the map $\Sigma_T\mathrm{MGL}/(x_2, \dots)\mathrm{MGL} \rightarrow \mathrm{MGL}/(x_2, \dots)\mathrm{MGL}$ given by multiplication by x_1 is the map $f_1\mathrm{MGL}/(x_2, \dots)\mathrm{MGL} \rightarrow \mathrm{MGL}/(x_2, \dots)\mathrm{MGL}$ showing the claim. \square

6. RATIONAL SLICES

We show in this paragraph that the assumptions from the last sections hold true after rationalization, at least over regular base schemes.

We denote by \mathbf{LQ} the Landweber spectrum associated to the MU_* -module \mathbb{Q} carrying the additive formal group law.

We note that any rational motivic Landweber spectrum has a decomposition into a sum of $\Sigma^{2i,i}\mathbf{LQ}$ for various i . This follows directly from the corresponding decomposition of the topological coefficients.

Since the Landweber coefficients for the rational K -theory spectrum are $\mathbb{Q}[v, v^{-1}]$ (v the Bott element in homological degree 2) we obtain

$$(6) \quad \mathrm{KGL}_{\mathbb{Q}} \cong \mathrm{PLQ} = \bigoplus_{i \in \mathbb{Z}} \Sigma^{2i,i}\mathbf{LQ},$$

compare also with [11, Theorem 10.1].

Using projectors [13, Theoreme IV.72] gives a decomposition

$$(7) \quad \mathrm{KGL}_{\mathbb{Q}} \cong \bigoplus_{i \in \mathbb{Z}} \mathbb{H}_B^{(i)}.$$

This decomposition is first defined for regular base schemes S . For morphisms between regular base schemes it pulls back. It thus makes sense to pull back the decomposition given over $\mathrm{Spec}(\mathbb{Z})$ to any base scheme, which we shall assume.

The next statement follows from the structure of a rational Snaith map

$$\mathrm{colim}_n(\mathbf{LQ} \wedge \Sigma^{-2n,-n}\Sigma^{\infty}\mathbf{P}_+^{\infty}) \rightarrow \mathrm{KGL}_{\mathbb{Q}}$$

(see [16] for the Snaith map) and the way the projectors for (7) are defined.

We call the map

$$\Sigma^{-2i,-i}\Sigma^{\infty}\mathbf{P}_+^{\infty} \rightarrow \mathrm{KGL}_{\mathbb{Q}}$$

the i 'th Snaith map.

Proposition 6.1. *The decompositions (6) and (7) coincide.*

Proof. We can assume the base is $\mathrm{Spec}(\mathbb{Z})$. We first compute the map

$$(8) \quad \mathbb{Q}^{\mathbb{Z}} \rightarrow \prod_{j \in \mathbb{Z}} \mathrm{Hom}(\Sigma^{2j,j}\mathbf{LQ}, \Sigma^{2j,j}\mathbf{LQ}) \rightarrow \mathrm{Hom}(\mathrm{PLQ}, \mathrm{PLQ}) \cong$$

$$\mathrm{Hom}(\mathrm{KGL}_{\mathbb{Q}}, \mathrm{KGL}_{\mathbb{Q}}) \rightarrow \mathrm{Hom}(\Sigma^{-2i,-i}\Sigma^{\infty}\mathbf{P}_+^{\infty}, \mathrm{KGL}_{\mathbb{Q}}) = (\mathrm{KGL}_{\mathbb{Q}}^{**}[x])^{2i,i} = \mathbb{Q}[u].$$

The first map in the second line is precomposition with the i 'th Snaith map. On the right hand side an appropriate Bott shift is applied to powers of x to obtain the powers of u . Note that for the system in i on the right hand side all transition maps are the same and given by $u^j \mapsto -ju^{j-1} + ju^j$ (this follows from the multiplicative formal group law for KGL). Note also that we define the Bott element to be the negative reduced class of \mathbf{P}^1 in \mathbf{P}^{∞} and \mathbf{BGL} .

We let $\mathrm{Hom}(\Sigma^{\infty}\mathbf{P}_+^{\infty}, \mathrm{PLQ}) = \mathbb{Q}[u']$. Here u' is the generator for the additive orientation on PLQ times the Bott element.

The multiplicative isomorphism

$$\mathbb{Q}[u] \cong \mathrm{Hom}(\Sigma^{\infty}\mathbf{P}_+^{\infty}, \mathrm{KGL}_{\mathbb{Q}}) \cong \mathrm{Hom}(\Sigma^{\infty}\mathbf{P}_+^{\infty}, \mathrm{PLQ}) \cong \mathbb{Q}[u']$$

is given by $u = 1 - e^{u'}$, which is the change of formal parameters between the multiplicative and additive formal group law.

Moreover the zero'th Snaith map to $\mathrm{KGL}_{\mathbb{Q}}$ is the element $1 - u \in \mathbb{Q}[[u]] \cong \mathrm{Hom}(\Sigma^{\infty} \mathbf{P}_{+}^{\infty}, \mathrm{KGL}_{\mathbb{Q}})$. Thus the Snaith map translates to

$$e^{u'} \in \mathbb{Q}[[u']] \cong \mathrm{Hom}(\Sigma^{\infty} \mathbf{P}_{+}^{\infty}, \mathrm{PLQ}) \cong \mathrm{Hom}_{\mathrm{Mod}_{\mathrm{SH}(\mathbb{Z})}}(\mathrm{LQ} \wedge \Sigma^{\infty} \mathbf{P}_{+}^{\infty}, \mathrm{PLQ}).$$

We let $\mathrm{LQ} \wedge \Sigma^{\infty} \mathbf{P}_{+}^{\infty} = \mathrm{LQ} \langle b_0, b_1, b_2, \dots \rangle$, where the b_i are dual generators to the powers of the orientation generator. It follows that after these identifications the Snaith map

$$\mathrm{LQ} \langle b_0, b_1, b_2, \dots \rangle \rightarrow \mathrm{PLQ}$$

has the effect

$$\mathrm{LQ} b_j \xrightarrow{\frac{1}{j!}} \Sigma^{2j, j} \mathrm{LQ}$$

on summands.

The Snaith map $\Sigma^{-2i, i} \Sigma^{\infty} \mathbf{P}_{+}^{\infty} \rightarrow \mathrm{PLQ}$ is just the same map with the appropriate Bott shift applied and is thus given on summands by

$$\Sigma^{-2i, -i} \mathrm{LQ} b_j \xrightarrow{\frac{1}{j!}} \Sigma^{-2i+2j, -i+j} \mathrm{LQ}.$$

Now let us start with the n 'th projector $p \in \mathrm{Hom}(\mathrm{PLQ}, \mathrm{PLQ})$, $n \in \mathbb{Z}$. Composition with the i 'th Snaith map gives us a map

$$\Sigma^{-2i, -i} \mathrm{LQ} \langle b_0, b_1, b_2, \dots \rangle \rightarrow \mathrm{PLQ}$$

sending, if $n \geq -i$, $\Sigma^{-2i, -i} b_{n+i}$ to $\frac{1}{(2n+2i)!} \Sigma^{2n, n} \mathrm{LQ}$ (abusing notation) and the other generators to 0. Thus this map is the element

$$\frac{1}{(2n+2i)!} u'^{2n+2i} \in \mathbb{Q}[[u']] = \mathrm{Hom}(\Sigma^{-2i, -i} \Sigma^{\infty} \mathbf{P}_{+}^{\infty}, \mathrm{PLQ}).$$

Applying the equation $u' = \ln(1 - u)$ we get the result that the map (8) sends the projector p , if $n \geq -i$, to

$$\frac{1}{(2n+2i)!} \ln(1 - u)^{2n+2i} \in \mathbb{Q}[[u]] = \mathrm{Hom}(\Sigma^{-2i, -i} \Sigma^{\infty} \mathbf{P}_{+}^{\infty}, \mathrm{KGL}_{\mathbb{Q}}),$$

otherwise to 0.

This is the same formula as used in [13] to define the projectors, see [13, Def. IV.62, after Cor. IV.67, Def. IV.71].

Now it suffices to observe that the Snaith system is indeed the system appearing in loc. cit. to describe endomorphisms of KGL and $\mathrm{KGL}_{\mathbb{Q}}$, see [13, Th. IV.13, before Prop. IV.33]. \square

Lemma 6.2. *The natural map $\mathrm{MGL}_{\mathbb{Q}} \rightarrow \mathrm{LQ}$ factors as an isomorphism $\mathrm{Q}_{\mathbb{Q}} \rightarrow \mathrm{LQ}$.*

Proof. The claim follows from the decompositions

$$\mathrm{MGL}_{\mathbb{Q}} \cong \mathrm{LQ}[b_1, b_2, \dots] \cong \mathrm{LQ}[x_1, x_2, \dots]$$

(see [11] for the definition of the b_i , see also [11, Theorem 10.5]). \square

Proposition 6.3. *Suppose S is regular. Then the map $\mathrm{MGL}_{\mathbb{Q}} \rightarrow \mathrm{LQ}$ is the map to the zero slice.*

Proof. First note that by the decomposition $\mathrm{MGL}_{\mathbb{Q}} \cong \mathrm{LQ}[b_1, b_2, \dots]$ and the effectivity of MGL it follows that LQ is effective. By lemma (6.2) we have to show that $\mathrm{MGL} \rightarrow \mathrm{LQ}$ is the map to the zero slice. The decomposition of $\mathrm{MGL}_{\mathbb{Q}}$ shows that for this it suffices to show that $\mathrm{Hom}_{\mathrm{SH}(S)}(\Sigma^{p,q}\Sigma_+^{\infty}X, \mathrm{LQ}) = 0$ for $X \in \mathrm{Sm}/S$, $p \in \mathbb{Z}$, $q \geq 1$. Equivalently we have to show that every map $\Sigma^{p,q}\Sigma_+^{\infty}X \rightarrow \mathrm{KGL}_{\mathbb{Q}}$, $X \in \mathrm{Sm}/S$, $p \in \mathbb{Z}$, $q \geq 0$, factors through $\mathrm{LQ}[u]$ with respect to the decomposition $\mathrm{KGL}_{\mathbb{Q}} \cong \mathrm{LQ}[u, u^{-1}]$. We can assume $q = 0$ by replacing X by $X \times \mathbb{G}_m^q$. Proposition (6.1) and [13, Corollaire VI.75] imply this for non-negative p . Suppose $p < 0$. Then by the periodicity of PLQ the claim we want to show is equivalent to the statement that for $p + 2q \geq 0$ every map $\Sigma^{p+2q,q}\Sigma_+^{\infty}X \rightarrow \mathrm{LQ}[u, u^{-1}]$ factors through $u^q\mathrm{LQ}[u]$. This follows from the decomposition [17, Corollary (5.5)(ii)] of the algebraic K -theory of the Laurent polynomials over a regular ring. \square

Corollary 6.4. *Suppose S is regular. Then $s_i(\mathrm{MGL}_{\mathbb{Q}}) \cong \Sigma_T^i\mathrm{LQ} \otimes \mathrm{MU}_{2i}$ compatible with the homomorphism $\mathrm{MU}_* \rightarrow \mathrm{MGL}_{\mathbb{Q},**}$.*

Proof. By proposition (6.3) we have $s_0(\mathrm{LQ}) = \mathrm{LQ}$. The claim follows from the decomposition $\mathrm{MGL}_{\mathbb{Q}} \cong \mathrm{LQ}[x_1, x_2, \dots]$. \square

Corollary 6.5. *Suppose S is regular. Then $s_0(\mathbf{1}_{\mathbb{Q}}) = s_0(\mathbf{1})_{\mathbb{Q}} = \mathrm{LQ}$.*

Proof. This follows now from Corollary (3.3). \square

REFERENCES

- [1] Daniel Dugger and Daniel C. Isaksen. Motivic cell structures. *Algebr. Geom. Topol.*, 5:615–652 (electronic), 2005.
- [2] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory*, volume 47 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [3] Philip S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [4] Jens Hornbostel. Localizations in motivic homotopy theory. *Math. Proc. Cambridge Philos. Soc.*, 140(1):95–114, 2006.
- [5] Mark Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [6] Thomas Hüttemann. Total cofibres of diagrams of spectra. *New York J. Math.*, 11:333–343 (electronic), 2005.
- [7] Tylor Lawson. Motivic homotopy. Notes of a course given by Mike Hopkins and Marc Levine at Harvard, <http://www.math.umn.edu/~tlawson/motivic.html>.
- [8] Marc Levine. Comparison of cobordism theories. Preprint, arXiv:0807.2238.
- [9] Marc Levine. The homotopy coniveau tower. *J. Topol.*, 1(1):217–267, 2008.
- [10] Niko Naumann, Markus Spitzweck, and Paul Arne Østvær. Chern classes, K -theory and Landweber exactness over nonregular base schemes. Preprint, arXiv:0809.0267.
- [11] Niko Naumann, Markus Spitzweck, and Paul Arne Østvær. Motivic Landweber exactness. Preprint, arXiv 0806.0274.
- [12] Pablo Pelaez. Multiplicative properties of the slice filtration. PhD thesis, arXiv 0806.1704.
- [13] Joël Riou. Opérations sur la K -théorie algébrique et régulateurs via la théorie homotopique des schémas. Thesis, K -theory Preprint Archives, 793.
- [14] M. Spitzweck. *Operads, Algebras and Modules in Model Categories and Motives*. PhD thesis, University of Bonn, 2001.
- [15] Markus Spitzweck. Slices of motivic Landweber spectra. Preprint, arXiv:0805.3350v1.
- [16] Markus Spitzweck and Paul Arne Østvær. The Bott inverted infinite projective space is homotopy K -theory. Preprint, <http://folk.uio.no/paularne/bott.pdf>.

- [17] V. Srinivas. *Algebraic K-theory*, volume 90 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1996.
- [18] Gabriele Vezzosi. Brown-Peterson spectra in stable \mathbb{A}^1 -homotopy theory. *Rend. Sem. Mat. Univ. Padova*, 106:47–64, 2001.
- [19] V. Voevodsky. On the zero slice of the sphere spectrum. *Tr. Mat. Inst. Steklova*, 246(Algebr. Geom. Metody, Svyazi i Prilozh.):106–115, 2004.
- [20] Vladimir Voevodsky. Open problems in the motivic stable homotopy theory. I. In *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, volume 3 of *Int. Press Lect. Ser.*, pages 3–34. Int. Press, Somerville, MA, 2002.

Fakultät für Mathematik, Universität Regensburg, Germany.
e-mail: Markus.Spitzweck@mathematik.uni-regensburg.de