



---

On modular symbols and the cohomology  
of Hecke triangle surfaces

Gabor Wiese

Preprint Nr. 19/2005

# On modular symbols and the cohomology of Hecke triangle surfaces

Gabor Wiese

4th November 2005

## Abstract

The aim of this article is to give a concise algebraic treatment of the modular symbols formalism, generalised from modular curves to Hecke triangle surfaces. A sketch is included of how the modular symbols formalism gives rise to the standard algorithms for the computation of holomorphic modular forms. Precise and explicit connections are established to the cohomology of Hecke triangle surfaces and group cohomology. In all the note a general commutative ring is used as coefficient ring in view of applications to the computation of modular forms over rings different from the complex numbers.

MSC Classification: 11F67 (primary), 11F75, 11Y40, 20H10 (secondary).

## 1 Introduction

The aim of this article is to give a clear and concise algebraic treatment of the modular symbols formalism and related objects in a form that generalises. We show how the standard algorithms for computing modular forms for congruence subgroups of  $SL_2(\mathbb{Z})$  can be deduced purely algebraically, using only the Eichler-Shimura isomorphism. This includes an algebraic proof of the presentation of modular symbols in terms of Manin symbols. Thus, we avoid the use of the technically rather difficult paper [Šokurov], which is present in all the published treatments known to the author (e.g. [MerelUniversal], [SteinThesis]).

The algebraic formulation that we give generalises immediately to the so-called Hecke triangle groups and we choose that general set-up from the beginning. It should, however, be pointed out that most Hecke triangle groups are non-arithmetic. In that case our treatment only gives an isomorphism between generalised modular symbols and modular forms, but does not yield an algorithm for the computation of the Fourier coefficients of the modular forms due to the absence of a suitable Hecke theory.

There is considerable interest in trying to use the modular symbols formalism over rings different from the complex number in order to compute modular forms over these rings (e.g. using the theory of Katz modular forms, see [EdixJussieu] or [Thesis]). This is the reason why the treatment of the modular symbols formalism in the present article is over any commutative unitary ring. In this generality

(in fact already over the integers) one notices quickly that, although the modular symbols formalism is inspired by the homology of modular curves, it does not quite compute it (compare Theorem 3.5 and Remark 5.10).

Moreover, in order to be able to treat questions like the computation of modular forms over more general rings one needs a geometric or algebraic interpretation of the modular symbols formalism in order to be able to establish a link with modular forms. For this reason we also treat certain cohomology groups on modular curves (as Riemann surfaces) and certain group cohomology groups which are both closely related to the modular symbols formalism. All the objects appearing are described by explicit formulae and a precise comparison is included. The cohomology group of the modular curve considered in the present article is a complex analogue of the étale cohomology group that one uses to define the 2-dimensional  $l$ -adic Galois representation associated to a modular form.

The differences between the various objects come from non-trivially stabilised points on the upper half plane. It is hence natural to use analytic modular stacks instead of modular curves and compare these two via the Leray spectral sequence. This has been carried out in the author's thesis. However, for the sake of the present article a formulation was chosen that uses only homological algebra, but gives the same results.

Apart from some facts about Hecke groups and some cohomology theory of groups and topological spaces, the treatment of the article is essentially self-contained.

## Overview

In Section 2 we present some facts about Hecke triangle groups, as well as two results to be used in the sequel. The following three sections are independent of each other. In Section 3 we introduce the modular symbols formalism extended to subgroups of finite index in Hecke triangle groups, and give a description in terms of Manin symbols. An explicit formula for the parabolic subspace of the group cohomology for subgroups of finite index of Hecke triangle groups is derived in Section 4. The subsequent Section 5 treats a similar cohomology group for a certain sheaf on the modular surface for  $\Gamma$ . An explicit formula is derived, which generalises a result of Merel's to higher weights. In the final section a comparison between the objects is carried out and it is sketched how the Eichler-Shimura theorem together with a theory of Hecke operators and the results from the previous sections can be used to compute modular forms for congruence subgroups of  $SL_2(\mathbb{Z})$ .

## Acknowledgements

This article has grown out of Chapter II of my thesis. I would like to thank my thesis advisor Bas Edixhoven for many useful suggestions, and Theo van den Bogaart for sharing his insights in algebraic geometry.


## 2 Hecke triangle groups and surfaces

For an integer  $n \geq 3$  one defines the  $n$ -th Hecke triangle group  $\Delta_n$  as the subgroup of  $\mathrm{PSL}_2(\mathbb{R})$  generated by

$$\sigma := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \tau := \begin{pmatrix} \lambda_n & -1 \\ 1 & 0 \end{pmatrix}$$

with  $\lambda_n = e^{\pi i/n} + e^{-\pi i/n} = 2 \cos(\pi/n)$ . In abuse of notation when writing a matrix we often mean its class modulo scalar matrices, i.e. an element of the projective linear group. The generation is free and  $\Delta_n$  is the free product of  $\mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$  and  $\mathbb{Z}/n\mathbb{Z} = \langle \tau \rangle$ , which makes the cohomological computations to come very simple. The stabiliser of the element  $\infty \in \mathbb{P}^1(\mathbb{R})$  is  $(\Delta_n)_\infty = \langle T \rangle$  with  $T = \tau\sigma = \begin{pmatrix} 1 & \lambda_n \\ 0 & 1 \end{pmatrix}$ . We denote by  $\Delta_n(\infty)$  the orbit of  $\infty$  under  $\Delta_n$ . As a special case let us note that  $\Delta_3 = \mathrm{PSL}_2(\mathbb{Z})$  and  $\Delta_3(\infty) = \mathbb{P}^1(\mathbb{Q})$ .

The Hecke group  $\Delta_n$  is a Fuchsian group of the first kind (i.e. a discrete subgroup of  $\mathrm{PSL}_2(\mathbb{R})$  of finite covolume) having parabolic elements (namely precisely the conjugates of powers of  $T$ ). Any subgroup  $\Gamma \leq \Delta_n$  of finite index is also a Fuchsian group of the first kind with parabolic elements. Moreover,  $Y_\Gamma := \Gamma \backslash \mathbb{H}$  can be given the structure of an open Riemann surface. It can be compactified  $X_\Gamma := Y_\Gamma \cup \Gamma \backslash \Delta_n(\infty)$ , where the set  $\Gamma \backslash \Delta_n(\infty)$  is the set of (parabolic) cusps of  $\Gamma$ . We write  $\overline{\mathbb{H}} = \mathbb{H} \cup \Delta_n(\infty)$ . The compact Riemann surface  $X_{\Delta_n}$  is called a Hecke triangle surface. The open Riemann surface  $Y_{\Delta_n}$  can be visualised as a fundamental domain with angles  $0$ ,  $\pi/n$  and  $\pi/n$ , as shown in the figure.



By a result of Leutbecher (as cited in [Schmidt]) Hecke triangle groups are non-arithmetic, except for  $n \in \{3, 4, 6\}$ , i.e.  $\Delta_n$  is not commensurable with any  $\mathrm{PSL}_2(\mathcal{O})$  for  $\mathcal{O}$  the ring of integers of a number field. For more details on Hecke triangle groups and surfaces we refer to [Schmidt] and the references therein.

The choice of working inside projective linear groups instead of linear groups was made since it simplifies some formulae and some proofs at nearly no costs.

### Notation

In most of the paper we use the following notation.

**2.1 Notation.** Let  $G = \Delta_n$  be a Hecke triangle group for some integer  $n \geq 3$  and let  $\Gamma \leq G$  be a subgroup of finite index. We use the notations  $Y_\Gamma$ ,  $X_\Gamma$  for the Riemann surfaces introduced above and call  $j : Y_\Gamma \hookrightarrow X_\Gamma$  the natural embedding and  $\pi : \mathbb{H} \twoheadrightarrow Y_\Gamma$  resp.  $\pi : \overline{\mathbb{H}} \twoheadrightarrow X_\Gamma$  the natural projections. Furthermore, the matrices  $\sigma, \tau, T$  defined above will be used.

We let  $R$  be a commutative ring with unit and  $V$  a left  $R[G]$ -module. If  $g \in G$  is some element of finite order  $m$ , we denote by  $N_g$  the element  $1 + g + \dots + g^{m-1}$  of the group ring  $R[G]$ . Similarly, if

$H \leq G$  is a finite subgroup, we write  $N_H = \sum_{h \in H} h \in R[G]$ .

## Mayer-Vietoris and amalgamated products

We assume Notation 2.1.

**2.2 Proposition.** *Let  $M$  be a left  $R[G]$ -module. Then the Mayer-Vietoris sequence gives the exact sequences*

$$\begin{aligned} 0 \rightarrow M^G \rightarrow M^{(\sigma)} \oplus M^{(\tau)} \rightarrow M \rightarrow H^1(G, M) \rightarrow H^1(\langle \sigma \rangle, M) \oplus H^1(\langle \tau \rangle, M) \rightarrow 0, \\ 0 \rightarrow H_1(\langle \sigma \rangle, M) \oplus H_1(\langle \tau \rangle, M) \rightarrow H_1(G, M) \rightarrow M \rightarrow M_{\langle \sigma \rangle} \oplus M_{\langle \tau \rangle} \rightarrow M_G \rightarrow 0 \end{aligned}$$

and for all  $i \geq 2$  isomorphisms

$$\begin{aligned} H^i(G, M) &\cong H^i(\langle \sigma \rangle, M) \oplus H^i(\langle \tau \rangle, M), \\ H_i(G, M) &\cong H_i(\langle \sigma \rangle, M) \oplus H_i(\langle \tau \rangle, M). \end{aligned}$$

**Proof.** Let us write  $G_1 := \langle \sigma \rangle$  and  $G_2 := \langle \tau \rangle$ . By [Brown], II.8.8, we have the exact sequence

$$0 \rightarrow R[G] \rightarrow R[G/G_1] \oplus R[G/G_2] \rightarrow R \rightarrow 0$$

of  $R[G]$ -modules, which are free as  $R$ -modules. Application of the functor  $\text{Hom}_R(\cdot, M)$  gives rise to the exact sequence of  $R[G]$ -modules

$$0 \rightarrow M \rightarrow \text{Hom}_{R[G_1]}(R[G], M) \oplus \text{Hom}_{R[G_2]}(R[G], M) \rightarrow \text{Hom}_R(R[G], M) \rightarrow 0.$$

The central terms, as well as the term on the right, can be identified with coinduced modules. Hence, the statements on cohomology follow by taking the long exact sequence of cohomology and invoking Shapiro's lemma.

Using the functor  $\cdot \otimes_R M$  gives rise to the analogous statements about homology.  $\square$

## Mackey's formula and stabilisers

We now prove Mackey's formula for coinduced modules. If  $H \leq G$  are groups and  $V$  is an  $R[H]$ -module, the coinduced module  $\text{Coind}_H^G V$  can be described as  $\text{Hom}_{R[H]}(R[G], V)$ .

**2.3 Proposition.** *Let  $R$  be a ring,  $G$  be a group and  $H, K$  subgroups of  $G$ . Let furthermore  $V$  be an  $R[H]$ -module. Then Mackey's formula*

$$\text{Res}_K^G \text{Coind}_H^G V \cong \prod_{g \in H \backslash G / K} \text{Coind}_{K \cap g^{-1} H g}^K ({}^g \text{Res}_{H \cap g K g^{-1}}^H V)$$

holds. Here  ${}^g \text{Res}_{H \cap g K g^{-1}}^H V$  denotes the  $R[K \cap g^{-1} H g]$ -module obtained from  $V$  via the conjugated action  $g^{-1} h g \cdot v := h \cdot v$  for  $v \in V$  and  $h \in H$  such that  $g^{-1} h g \in K$ .

**Proof.** We consider the commutative diagram

$$\begin{array}{ccc}
\mathrm{Res}_K^G \mathrm{Hom}_H(R[G], V) & \longrightarrow & \prod_{g \in H \backslash G/K} \mathrm{Hom}_{K \cap g^{-1}Hg} (R[K], {}^g(\mathrm{Res}_{H \cap gKg^{-1}}^H V)) \\
& \searrow & \downarrow \sim \\
& & \prod_{g \in H \backslash G/K} \mathrm{Hom}_{H \cap gKg^{-1}} (R[gKg^{-1}], \mathrm{Res}_{H \cap gKg^{-1}}^H V).
\end{array}$$

The vertical arrow is just given by conjugation and is clearly an isomorphism. The diagonal map is the product of the natural restrictions. From the bijection

$$(H \cap gKg^{-1}) \backslash gKg^{-1} \xrightarrow{gkg^{-1} \mapsto Hgk} H \backslash HgK$$

it is clear that also the diagonal map is an isomorphism, proving the proposition.  $\square$

Applying Shapiro's lemma, one immediately obtains the following two corollaries.

**2.4 Corollary.** *In the situation of Proposition 2.3 one has*

$$\begin{aligned}
H^i(K, \mathrm{Coind}_H^G V) &\cong \prod_{g \in H \backslash G/K} H^i(K \cap g^{-1}Hg, {}^g(\mathrm{Res}_{H \cap gKg^{-1}}^H V)) \\
&\cong \prod_{g \in H \backslash G/K} H^i(H \cap gKg^{-1}, \mathrm{Res}_{H \cap gKg^{-1}}^H V)
\end{aligned}$$

for all  $i \in \mathbb{N}$ .

**2.5 Corollary.** *We now assume Notation 2.1. For  $x \in \overline{\mathbb{H}}$  we denote by  $G_x$  the stabiliser subgroup of  $G$  for the point  $x$ . The image of the  $G$ -orbit of  $x$  in  $X_\Gamma$  is in bijection with the double cosets  $\Gamma \backslash G/G_x$  as follows*

$$\Gamma \backslash G/G_x \xrightarrow{g \mapsto gx} \Gamma \backslash Gx.$$

Moreover, the group  $\Gamma \cap gG_xg^{-1}$  equals  $\Gamma_{gx}$ , the stabiliser subgroup of  $\Gamma$  for the point  $gx$ . Thus, for all  $i \in \mathbb{N}$  Mackey's formula gives an isomorphism

$$H^i(G_x, \mathrm{Coind}_\Gamma^G V) \cong \prod_{y \in \Gamma \backslash Gx} H^i(\Gamma_y, V).$$

### 3 The modular symbols formalism

Modular symbols were systematically studied by [Manin]. In the book [Cremona] it is shown how the modular symbols formalism can be used for computing weight two modular forms. A generalisation to higher weight modular forms was found by [Šokurov], and the resulting “higher weights modular symbols formalism” was used by [MerelUniversal] algorithmically. This formalism is also the subject of the very comprehensive textbook [SteinBook], which is in preparation.

Common to the mentioned treatments is that they use rather difficult computations of homology groups. We show in this section that the description of modular symbols in terms of Manin symbols can be derived purely algebraically with simple and conceptual methods. Together with the Eichler-Shimura isomorphism, the proof of which is also quite easy, this already gives the basic idea of the algorithms of [SteinBook]. This is sketched in Section 6. These algorithms were implemented in Magma and Python by William Stein in the case of the standard congruence subgroups.

## Definition

Modular symbols can be thought of as geodesic paths between two cusps resp. as the associated homology class relative to the cusps. We shall, however, give a combinatorial definition, as is implemented in Magma and like the one in [MereUniversal], [Cremona] and Stein's thesis [SteinThesis], except that we do not factor out torsion, but intend a common treatment for all rings.

We give the definition in the more general context of Hecke triangle groups and also allow general modules. In this section we assume Notation 2.1.

**3.1 Definition.** *We define the  $R$ -modules*

$$\mathcal{M}_R := R[\{\alpha, \beta\} | \alpha, \beta \in G(\infty)] / \langle \{\alpha, \alpha\}, \{\alpha, \beta\} + \{\beta, \gamma\} + \{\gamma, \alpha\} | \alpha, \beta, \gamma \in G(\infty) \rangle$$

and

$$\mathcal{B}_R := R[G(\infty)].$$

We equip both with the natural left  $\Gamma$ -action. Furthermore, we let

$$\mathcal{M}_R(V) := \mathcal{M}_R \otimes_R V \quad \text{and} \quad \mathcal{B}_R(V) := \mathcal{B}_R \otimes_R V$$

for the left diagonal  $\Gamma$ -action.

(a) We call the  $\Gamma$ -coinvariants

$$\mathcal{M}_R(\Gamma, V) := \mathcal{M}_R(V)_\Gamma = \mathcal{M}_R(V) / \langle (x - gx) | g \in \Gamma, x \in \mathcal{M}_R(V) \rangle$$

the space of  $(\Gamma, V)$ -modular symbols.

(b) We call the  $\Gamma$ -coinvariants

$$\mathcal{B}_R(\Gamma, V) := \mathcal{B}_R(V)_\Gamma = \mathcal{B}_R(V) / \langle (x - gx) | g \in \Gamma, x \in \mathcal{B}_R(V) \rangle$$

the space of  $(\Gamma, V)$ -boundary symbols.

(c) We define the boundary map as the map

$$\mathcal{M}_R(\Gamma, V) \rightarrow \mathcal{B}_R(\Gamma, V)$$

which is induced from the map  $\mathcal{M}_R \rightarrow \mathcal{B}_R$  sending  $\{\alpha, \beta\}$  to  $\{\beta\} - \{\alpha\}$ .

(d) The kernel of the boundary map is denoted by  $\mathcal{CM}_R(\Gamma, V)$  and is called the space of cuspidal  $(\Gamma, V)$ -modular symbols.

(e) The image of the boundary map inside  $\mathcal{B}_R(\Gamma, V)$  is denoted by  $\mathcal{E}_R(\Gamma, V)$  and is called the space of  $(\Gamma, V)$ -Eisenstein symbols.

### Manin symbols

Manin symbols provide an explicit description of modular symbols. We stay in the general setting over a ring  $R$  and keep Notation 2.1.

As  $G$  is infinite, the induced module  $R[G]$  is not isomorphic to the coinduced one  $\text{Hom}_R(R[G], R)$  and  $R[G]$  is not cohomologically trivial. However,  $H_{\text{par}}^1(G, R[G]) = 0$  (for the definition see Section 4). This is the essence of the following proposition.

**3.2 Proposition.** *The sequence of  $R$ -modules*

$$0 \rightarrow R[G]N_\sigma + R[G]N_\tau \rightarrow R[G] \xrightarrow{g \mapsto g(1-\sigma)\infty} R[G(\infty)] \xrightarrow{g\infty \mapsto 1} R \rightarrow 0$$

is exact.

**Proof.** We first use that  $R[G]$  is a cohomologically trivial module for both  $\langle\sigma\rangle$  and  $\langle\tau\rangle$ . This gives

$$R[G]N_\sigma = \ker_{R[G]}(1 - \sigma) = R[G]^{\langle\sigma\rangle}, \quad R[G]N_\tau = \ker_{R[G]}(1 - \tau) = R[G]^{\langle\tau\rangle},$$

$$R[G](1 - \sigma) = \ker_{R[G]} N_\sigma \quad \text{and} \quad R[G](1 - \tau) = \ker_{R[G]} N_\tau.$$

Proposition 2.2 yields the exact sequence

$$0 \rightarrow R[G] \rightarrow R[G]_{\langle\sigma\rangle} \oplus R[G]_{\langle\tau\rangle} \rightarrow R \rightarrow 0,$$

since  $H_1(G, R[G]) = 0$  by Shapiro's lemma because  $R[G] \cong \text{Ind}_1^G(R)$ . In fact, this sequence is at the origin of our proof of Proposition 2.2. The injectivity of the first map in the exact sequence means

$$R[G](1 - \sigma) \cap R[G](1 - \tau) = 0.$$

We identify  $R[G]/R[G](1 - T)$  with  $R[G(\infty)]$  by sending  $g$  to  $g\infty$ . Now we show the exactness at  $R[G]$ , which comes down to proving that the equation  $x(1 - \sigma) = y(1 - T)$  for  $x, y \in R[G]$  implies that  $x$  is in  $R[G]^{\langle\sigma\rangle} + R[G]^{\langle\tau\rangle}$ .

Using the formula  $\tau = T\sigma$  we obtain that  $x(1 - \sigma) = y(1 - T) = y(1 - \tau) - yT(1 - \sigma)$ . This yields  $x(1 - \sigma) + yT(1 - \sigma) = y(1 - \tau)$ . This expression, however, is zero. Consequently, there is a  $z \in R[G]$  such that  $y = zN_\tau$ . Hence, using  $T = \tau\sigma$  and consequently  $N_\tau T = N_\tau\sigma$ , we get

$$y(1 - T) = zN_\tau(1 - T) = zN_\tau(1 - \sigma) = y(1 - \sigma).$$

The equation  $x(1 - \sigma) = y(1 - \sigma)$  means that  $x - y$  is in  $R[G]^{(\sigma)}$ . As we know that  $y \in R[G]^{(\tau)}$ , we see that  $x = (x - y) + y$  is in  $R[G]^{(\sigma)} + R[G]^{(\tau)}$ , as required. Note that instead of this explicit calculation we could also have appealed to Proposition 4.3.

The exactness at  $R[G(\infty)]$  can be seen as follows (we avoid here the traditional continued fractions argument, as it does not obviously generalise to Hecke triangle groups and is not in the spirit of the present group theoretic approach). Since  $\sigma$  and  $T = \tau\sigma$  generate  $G$ , the kernel of  $R[G] \xrightarrow{g \mapsto 1} R$  is  $R[G](1 - \sigma) + R[G](1 - T)$ . Taking the quotient by  $R[G](1 - T)$  gives the desired exactness.  $\square$

**3.3 Lemma.** *The sequence of  $R$ -modules*

$$0 \rightarrow \mathcal{M}_R \xrightarrow{\{\alpha, \beta\} \mapsto \beta - \alpha} R[G(\infty)] \xrightarrow{\alpha \mapsto 1} R \rightarrow 0$$

is exact.

**Proof.** The injectivity of the first arrow is clear, since we can write any element in  $\mathcal{M}_R$  as  $\sum_{\alpha \neq \infty} r_\alpha \{\infty, \alpha\}$  with  $r_\alpha \in R$ , using the relations defining  $\mathcal{M}_R$ . The image of this element under the first arrow is  $\sum_{\alpha \neq \infty} r_\alpha \alpha - (\sum_{\alpha \neq \infty} r_\alpha) \infty$ . If this is zero, clearly all  $r_\alpha$  are zero, proving the injectivity of the first arrow.

Suppose now we are given  $\sum_{\alpha} r_\alpha \alpha \in R[G(\infty)]$  in the kernel of the second arrow. Then  $\sum_{\alpha} r_\alpha = 0$  and consequently we have

$$\sum_{\alpha} r_\alpha \alpha = \sum_{\alpha \neq \infty} r_\alpha \alpha - \left( \sum_{\alpha \neq \infty} r_\alpha \right) \infty$$

which is in the image of the first arrow, as noticed before.  $\square$

**3.4 Proposition.** *The homomorphism of  $R$ -modules*

$$R[G] \xrightarrow{\phi} \mathcal{M}_R, \quad g \mapsto \{g, 0, g, \infty\}$$

is surjective and its kernel is given by  $R[G]N_\sigma + R[G]N_\tau$ .

**Proof.** This is a direct consequence of Proposition 3.2 and Lemma 3.3.  $\square$

We are now ready to prove the description of modular symbols in terms of Manin symbols.

**3.5 Theorem.** *Recall that we are assuming Notation 2.1. Let  $M = \text{Ind}_\Gamma^G(V)$ , which we identify with  $(R[G] \otimes_R V)_\Gamma$ . That module carries the right  $R[G]$ -action  $(h \otimes v)g = (hg \otimes v)$ , and the  $\Gamma$ -coinvariants are taken for the diagonal left  $\Gamma$ -action. The following statements hold:*

(a) *The homomorphism  $\phi$  from Proposition 3.4 induces the exact sequence of  $R$ -modules*

$$0 \rightarrow MN_\sigma + MN_\tau \rightarrow M \rightarrow \mathcal{M}_R(\Gamma, V) \rightarrow 0.$$

(b) The homomorphism  $R[G] \rightarrow R[G(\infty)]$  sending  $g$  to  $g.\infty$  induces the exact sequence of  $R$ -modules

$$0 \rightarrow M(1 - T) \rightarrow M \rightarrow \mathcal{B}_R(\Gamma, V) \rightarrow 0.$$

(c) The identifications of (a) and (b) imply the isomorphism

$$\mathcal{CM}_R(\Gamma, V) \cong \ker \left( M / (MN_\sigma + MN_\tau) \xrightarrow{m \mapsto m(1-\sigma)} M / M(1 - T) \right).$$

**Proof.** (a) We derive this from Proposition 3.4, which gives the exact sequence

$$0 \rightarrow R[G]N_\sigma + R[G]N_\tau \rightarrow R[G] \rightarrow \mathcal{M}_2(R) \rightarrow 0.$$

Tensoring with  $V$  over  $R$ , we obtain the exact sequence of left  $R[\Gamma]$ -modules

$$0 \rightarrow (R[G] \otimes_R V)N_\sigma + (R[G] \otimes_R V)N_\tau \rightarrow (R[G] \otimes_R V) \rightarrow \mathcal{M}_R(V) \rightarrow 0.$$

Passing to left  $\Gamma$ -coinvariants yields (a). Part (b) is clear from the definition and Part (c) has already been noticed in the proof of Proposition 3.2.  $\square$

In the literature on Manin symbols one usually finds different versions of the module  $M$ , namely the following. Suppose first that the  $\Gamma$ -action on  $V$  is the restriction of some  $G$ -action on  $V$ . Then we have the isomorphism

$$(R[G] \otimes_R V)_\Gamma \cong R[\Gamma \backslash G] \otimes_R V, \quad g \otimes v \mapsto g \otimes g^{-1}v.$$

The right  $R[G]$ -action carries over to the action  $(\Gamma h \otimes v)g = \Gamma hg \otimes g^{-1}v$ .

We should also mention a slight variant of this. Suppose now that  $\tilde{G}$  and  $\tilde{\Gamma}$  are defined as  $\phi^{-1}(G)$  resp.  $\phi^{-1}(\Gamma)$  for the projection  $\phi : \mathrm{SL}_2(\mathbb{R}) \twoheadrightarrow \mathrm{PSL}_2(\mathbb{R})$ . We also assume that  $-1 \in \tilde{\Gamma}$ , so that  $V$  is an  $R[\tilde{\Gamma}]$ -module, and that the  $\tilde{\Gamma}$ -action on  $V$  is the restriction of some  $\tilde{G}$ -action on  $V$ . Then we have the isomorphism

$$(R[G] \otimes_R V)_\Gamma \cong R[\tilde{\Gamma} \backslash \tilde{G}] \otimes_R V, \quad g \otimes v \mapsto g \otimes g^{-1}v.$$

## 4 Group cohomology

Also in this section we use Notation 2.1.

### Definitions

We define the *parabolic group cohomology group* as the left hand term and the *boundary group cohomology group* as the right hand term in the exact sequence

$$0 \rightarrow H_{\mathrm{par}}^1(\Gamma, V) \rightarrow H^1(\Gamma, V) \xrightarrow{\mathrm{res}} \prod_{c \in \Gamma \backslash G(\infty)} H^1(\Gamma_{\tilde{c}}, V),$$

where  $\Gamma_{\tilde{c}}$  is the stabiliser subgroup of  $\Gamma$  for the cusp  $\tilde{c} \in \overline{\mathbb{H}}$  with  $\pi(\tilde{c}) = c$ .

## Computing group cohomology

In order to compute the group cohomology for  $\Gamma$ , it suffices to compute the cohomology for  $G$  because of Shapiro's lemma, which for any  $R[\Gamma]$ -module  $V$  gives an isomorphism

$$H^1(G, \text{Coind}_\Gamma^G V) \cong H^1(\Gamma, V).$$

Due to Corollary 2.5 it is clear that Shapiro's lemma respects the parabolic subspace.

A first, however, not complete computation of the group cohomology of  $R[G]$ -modules is provided by the Mayer-Vietoris sequence (Proposition 2.2). We now derive an explicit description.

**4.1 Proposition.** *Let  $M$  be a left  $R[G]$ -module. Then the sequence of  $R$ -modules*

$$0 \rightarrow M^G \rightarrow M \rightarrow \ker_M N_\sigma \times \ker_M N_\tau \rightarrow H^1(G, M) \rightarrow 0$$

*is exact.*

**Proof.** We determine the 1-cocycles of  $M$ . Apart from  $f(1) = 0$ , they must satisfy

$$0 = f(\sigma^2) = \sigma f(\sigma) + f(\sigma) = N_\sigma f(\sigma) \text{ and}$$

$$0 = f(\tau^n) = \dots = N_\tau f(\tau).$$

Since these are the only relations in  $G$ , a cocycle is uniquely given by the choices

$$f(\sigma) \in \ker_M N_\sigma \text{ and } f(\tau) \in \ker_M N_\tau.$$

The 1-coboundaries are precisely the cocycles  $f$  which satisfy  $f(\sigma) = (1 - \sigma)m$  and  $f(\tau) = (1 - \tau)m$  for some  $m \in M$ . This proves

$$H^1(G, M) \cong (\ker_M N_\sigma \times \ker_M N_\tau) / (((1 - \sigma)m, (1 - \tau)m) \mid m \in M).$$

Rewriting yields the proposition. □

**4.2 Remark.** *As  $G_\infty = \langle T \rangle < G$  is infinite cyclic, one has  $H^1(G_\infty, \text{Res}_G^{G_\infty} M) \cong M / (1 - T)M$ .*

An explicit presentation of the parabolic group cohomology is the following.

**4.3 Proposition.** *The parabolic group cohomology group sits in the exact sequence*

$$0 \rightarrow M^{(T)} / M^G \rightarrow \ker_M N_\sigma \cap \ker_M N_\tau \xrightarrow{\phi} H_{\text{par}}^1(G, M) \rightarrow 0,$$

*where  $\phi$  maps an element  $m$  to the 1-cocycle  $f$  uniquely determined by  $f(\sigma) = f(\tau) = m$ .*

**Proof.** Using Proposition 4.1, we have the exact commutative diagram

$$\begin{array}{ccccc}
M^{(T)}/M^{G\subset} & \xrightarrow{(\sigma^{-1}-1)} & \ker N_\sigma \cap \ker N_\tau & \longrightarrow & H_{\text{par}}^1(G, M) \\
\downarrow \sigma^{-1} & & \downarrow & & \downarrow \\
M/M^{G\subset} & \xrightarrow{(1-\sigma, 1-\tau)} & \ker N_\sigma \times \ker N_\tau & \twoheadrightarrow & H^1(G, M) \\
\downarrow (1-T)\sigma & & \downarrow (a,b) \mapsto b-a & & \downarrow \\
(1-T)M^{G\subset} & \longrightarrow & M & \twoheadrightarrow & H^1(G_\infty, M).
\end{array}$$

As the bottom left vertical arrow is surjective, the claim follows from the snake lemma.  $\square$

## 5 Cohomology of Hecke triangle surfaces

### The group cohomology presheaf and sheaf

In this section we let  $X$  be a topological space,  $R$  a commutative ring and  $\Gamma$  a group. For any ring  $S$ , not necessarily commutative, we denote by  $S - \mathfrak{Mod}$  the category of  $S$ -modules and by  $\mathfrak{Sh}_X(S - \mathfrak{Mod})$  the category of sheaves of  $S$ -modules on  $X$ .

We collect some well-known, but important, properties in the following proposition.

**5.1 Proposition.** (a) *The category  $\mathfrak{Sh}_X(S - \mathfrak{Mod})$  has enough injectives.*

(b) *Let  $\mathcal{I} \in \mathfrak{Sh}_X(S - \mathfrak{Mod})$  be an injective object. Then  $\mathcal{I}$  is flabby (flasque).*

(c) *Let  $\mathcal{V}$  be an object of  $\mathfrak{Sh}_X(R[\Gamma] - \mathfrak{Mod})$ . Then the cohomology groups  $H^i(X, \mathcal{V})$  for  $i \geq 0$  do not depend on whether they are computed in the category  $\mathfrak{Sh}_X(R[\Gamma] - \mathfrak{Mod})$  or by forgetting the  $\Gamma$ -action in the category  $\mathfrak{Sh}_X(R - \mathfrak{Mod})$ .*

(d) *Let  $\mathcal{I} \in \mathfrak{Sh}_X(S - \mathfrak{Mod})$  be an injective object. Then for all open sets  $U \subseteq X$ , the object  $\mathcal{I}(U)$  of  $S - \mathfrak{Mod}$  is injective.*

(e) *Let  $\mathcal{I} \in \mathfrak{Sh}_X(R[\Gamma] - \mathfrak{Mod})$  be an injective object. Then  $\mathcal{I}^\Gamma$  is an injective object of  $\mathfrak{Sh}_X(R - \mathfrak{Mod})$ .*

**Proof.** We notice that  $X$  together with the constant sheaf  $S$  on  $X$  is a ringed space. The statements (a) and (b) are then [Hartshorne], Proposition 2.2 and Lemma 2.4. We should, however, point out, that Hartshorne works with commutative rings only. But the proofs also work in the non-commutative situation.

(c) follows from (b), as flabby resolutions can be used for computing  $H^i(X, \mathcal{V})$  (see [Hartshorne], Proposition 2.5).

(d) As the sheaf  $\mathcal{I}$  restricted to  $U$  is injective, it suffices to prove the statement for  $U = X$ . The injectivity of  $\mathcal{I}$  means that the functor  $\text{Hom}_{\mathfrak{Sh}_X(S-\mathfrak{Mod})}(\cdot, \mathcal{I})$  is exact. For  $A \in S - \mathfrak{Mod}$ , we denote by  $\underline{A}$  the constant sheaf on  $X$  associated to  $A$ . We have

$$\text{Hom}_{S-\mathfrak{Mod}}(A, \mathcal{I}(X)) = \text{Hom}_{\mathfrak{Sh}_X(S-\mathfrak{Mod})}(\underline{A}, \mathcal{I})$$

for all  $A \in S - \mathfrak{Mod}$ . As taking the constant sheaf is an exact functor,  $\text{Hom}_{S-\mathfrak{Mod}}(\cdot, \mathcal{I}(X))$  is also exact, proving the injectivity of  $\mathcal{I}(X)$ .

(e) Suppose we are given a diagram

$$\begin{array}{ccc} & & \mathcal{I}^\Gamma \\ & \nearrow & \\ A & \hookrightarrow & B \end{array}$$

in  $\mathfrak{Sh}_X(R - \mathfrak{Mod})$ . By composing with the natural injection  $\mathcal{I}^\Gamma \hookrightarrow \mathcal{I}$  and putting a trivial  $\Gamma$ -action on  $A$  and  $B$ , we obtain the commutative diagram

$$\begin{array}{ccc} & & \mathcal{I} \\ & \nearrow & \uparrow \\ A & \hookrightarrow & B \end{array}$$

in  $\mathfrak{Sh}_X(R[\Gamma] - \mathfrak{Mod})$ , since  $\mathcal{I}$  is injective. However, the image of  $B \rightarrow \mathcal{I}$  is contained in  $\mathcal{I}^\Gamma$ , as  $B$  is a trivial  $\Gamma$ -module.  $\square$

Let  $U \subseteq X$  be an open set. We consider the following commutative diagram of categories:

$$\begin{array}{ccccc} & & \mathfrak{Sh}_X(R - \mathfrak{Mod}) & & \\ & \nearrow^{(\cdot)^\Gamma} & & \searrow^{H^0(U, \cdot)} & \\ \mathfrak{Sh}_X(R[\Gamma] - \mathfrak{Mod}) & & & & R - \mathfrak{Mod} \\ & \searrow_{H^0(U, \cdot)} & & \nearrow_{(\cdot)^\Gamma} & \\ & & R[\Gamma] - \mathfrak{Mod} & & \end{array}$$

Let  $\mathcal{V} \in \mathfrak{Sh}_X(R[\Gamma] - \mathfrak{Mod})$ . Due to Proposition 5.1, Grothendieck's theorem on spectral sequences ([Weibel], Theorem 5.8.3) gives rise to the two spectral sequences

$$H^p(\Gamma, H^q(U, \mathcal{V})) \Rightarrow R^{p+q}(H^0(U, \cdot)^\Gamma)(\mathcal{V}) \quad (5.1)$$

and

$$H^p(U, \mathcal{H}^q(\Gamma, \mathcal{V})) \Rightarrow R^{p+q}(H^0(U, \cdot)^\Gamma)(\mathcal{V}), \quad (5.2)$$

where we write

$$\mathcal{H}^q(\Gamma, \mathcal{V}) = R^q((\cdot)^\Gamma)(\mathcal{V}).$$

This sheaf is called the *group cohomology sheaf*. We now consider the following composite of edge morphisms of the above spectral sequences for  $p \geq 1$

$$H^p(\Gamma, H^0(U, \mathcal{V})) \rightarrow R^p(H^0(U, \cdot)^\Gamma)(\mathcal{V}) \rightarrow H^0(U, \mathcal{H}^p(\Gamma, \mathcal{V})). \quad (5.3)$$

Also by [Weibel], Theorem 5.8.3, the edge morphisms are the natural maps. If  $W \subseteq U$  is an open set, then the restriction  $\text{res}_W^U : \mathcal{V}(U) \rightarrow \mathcal{V}(W)$  induces natural maps on each of the three objects, which are also denoted by  $\text{res}_W^U$ . It can be checked that the diagram

$$\begin{array}{ccccc} H^p(\Gamma, H^0(U, \mathcal{V})) & \longrightarrow & R^p(H^0(U, \cdot)^\Gamma)(\mathcal{V}) & \longrightarrow & H^0(U, \mathcal{H}^p(\Gamma, \mathcal{V})) \\ \downarrow \text{res}_W^U & & \downarrow \text{res}_W^U & & \downarrow \text{res}_W^U \\ H^p(\Gamma, H^0(W, \mathcal{V})) & \longrightarrow & R^p(H^0(W, \cdot)^\Gamma)(\mathcal{V}) & \longrightarrow & H^0(W, \mathcal{H}^p(\Gamma, \mathcal{V})) \end{array}$$

is commutative. In other words, the edge morphisms 5.3 give morphisms of presheaves

$$(U \mapsto H^p(\Gamma, H^0(U, \mathcal{V}))) \rightarrow (U \mapsto R^p(H^0(U, \cdot)^\Gamma)(\mathcal{V})) \rightarrow \mathcal{H}^p(\Gamma, \mathcal{V}).$$

We call the first presheaf the *group cohomology presheaf*. This terminology is justified, as the sheafification coincides with the group cohomology sheaf in the cases of interest in the present context.

**5.2 Proposition.** *Assume that  $\Gamma$  is of type  $(FP)_\infty$  over  $R$  (cf. [Bieri], p. 6). The morphisms of presheaves above become isomorphisms on the sheafification. In particular, for  $x \in X$  one has  $\mathcal{H}^i(\Gamma, \mathcal{V})_x \cong H^i(\Gamma, \mathcal{V}_x)$  for all  $i \in \mathbb{N}$ .*

**Proof.** It follows directly from the definition that the second map is sheafification. Indeed, let  $\mathcal{V} \hookrightarrow \mathcal{I}^\bullet$  be an injective resolution. Then

$$\mathcal{H}^p(\Gamma, \mathcal{V}) = \ker((\mathcal{I}^p)^\Gamma \rightarrow (\mathcal{I}^{p+1})^\Gamma) / \text{im}((\mathcal{I}^{p-1})^\Gamma \rightarrow (\mathcal{I}^p)^\Gamma)$$

is by definition the sheafification of

$$U \mapsto R^p(H^0(U, \cdot)^\Gamma)(\mathcal{V}) = \ker(\mathcal{I}^p(U)^\Gamma \rightarrow \mathcal{I}^{p+1}(U)^\Gamma) / \text{im}(\mathcal{I}^{p-1}(U)^\Gamma \rightarrow \mathcal{I}^p(U)^\Gamma).$$

As taking stalks is exact, we have for  $x \in X$  that  $\mathcal{V}_x \hookrightarrow \mathcal{I}_x^\bullet$  is exact in the category of  $R[\Gamma]$ -modules. We claim that this is a  $\Gamma$ -acyclic resolution of  $\mathcal{V}_x$ . By Proposition 5.1 (d) we know that for all  $U \subset X$  open  $\mathcal{I}^i(U)$  is an injective  $R[\Gamma]$ -module for all  $i \geq 0$ , and hence that  $H^q(\Gamma, \mathcal{I}^i(U)) = 0$  for all  $q \geq 1$ . Under the assumption by [Bieri], Proposition 2.4, we know that the functor  $H^i(\Gamma, \cdot)$  commutes with direct limits, whence, indeed,

$$H^q(\Gamma, \mathcal{I}_x^i) = \lim_{\rightarrow U \ni x} H^q(\Gamma, \mathcal{I}^i(U)) = 0$$

for all  $q \geq 1$  and all  $i \geq 0$ , as claimed.

The particular statement follows directly. □

## A spectral sequence for Hecke triangle surfaces

We assume again Notation 2.1 and we let  $\mathcal{C} \in \{\mathbb{H}, \overline{\mathbb{H}}\}$  and  $X = \Gamma \backslash \mathcal{C} \in \{Y_\Gamma, X_\Gamma\}$ .

We recall some facts on  $\overline{\mathbb{H}}$  that we will use in the sequel. The topology on  $\overline{\mathbb{H}}$  extends the topology of  $\mathbb{H}$  and is obtained as follows. Using the action of  $G$  it suffices to give a system of open neighbourhoods of the cusp  $\infty$ , which is provided by the sets  $U_T = \{a + ib \mid b > T\}$  for every real  $T > 0$ . Clearly, the intersection with any open set in  $\mathbb{H}$  is an open set in  $\mathbb{H}$ .

The  $G$ -orbit of every  $x \in \overline{\mathbb{H}}$  is a discrete set. That actually holds for every subgroup of  $\mathrm{SL}_2(\mathcal{O})$  with  $\mathcal{O}$  a ring of integers of a totally real number field. Indeed, as it holds on  $\mathbb{H}$ , we only need to check it on the cusps. We have  $\mathrm{Im}(g(x + iy)) = y/((cx + d)^2 + c^2y^2)$  for  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ . If  $g$  does not stabilise the cusp  $\infty$ , we have  $c \neq 0$ , so that  $\mathrm{Im}(g(x + iy)) \leq 1/(c^2y) \leq C^{-2}y^{-1}$  with  $C$  the smallest non-zero modulus of an element in  $\mathcal{O}$ . This means that  $U_T \cap gU_T = \emptyset$  for any  $T > 1/C$ . Hence, for any  $hU_T \cap gU_T = \emptyset$ , unless  $g\infty = h\infty$ .

Next, we claim that  $\overline{\mathbb{H}}$  is simply connected. It suffices to show that any loop  $L$  starting and ending in  $\infty$  is contractible. We may assume that it does not pass through any other cusp (otherwise, we cut the loop into several loops each one meeting only one cusp). The following homotopy works

$$g : [0, 1] \times [0, 1] \rightarrow \overline{\mathbb{H}}, (s, x) \mapsto \begin{cases} \frac{L(x)}{s} & \text{if } s \neq 0 \text{ and } f(x) \neq \infty, \\ \infty & \text{otherwise.} \end{cases}$$

The map  $g$  is continuous, as the preimage of  $U_T$  is  $[0, 1] \times [0, 1] \cap \{(s, x) \mid \mathrm{Im}(f(x)) > sT\}$ , and, thus, is open in  $[0, 1] \times [0, 1]$ . Clearly,  $g(0, x)$  is the constant path from  $\infty$  to  $\infty$ , whereas  $g(1, x) = L(x)$  is the loop we started with.

Denote by  $\underline{V}$  the constant sheaf on  $\mathcal{C}$  associated to  $V$  together with its natural  $\Gamma$ -action, i.e. for an open set  $U \subset \mathcal{C}$  we let  $\underline{V}(U) = \mathrm{Hom}_{\mathrm{cts}}(U, V)$  (equipping  $V$  with the discrete topology) together with isomorphisms  $\phi_g : \underline{V} \rightarrow g_*\underline{V}$  for each  $g \in \Gamma$  which on  $U$  are given by

$$\mathrm{Hom}_{\mathrm{cts}}(U, V) \rightarrow \mathrm{Hom}_{\mathrm{cts}}(gU, V), \quad f \mapsto (gu \mapsto gf(u) \quad \forall u \in U).$$

We have that  $\pi_*\underline{V}$  is in  $\mathfrak{Sh}_X(R[\Gamma] - \mathfrak{Mod})$ .

**5.3 Lemma.** *For any point  $y \in \mathcal{C}$  and any sheaf  $\mathcal{F} \in \mathfrak{Sh}_{\mathcal{C}}(R - \mathfrak{Mod})$  there is an isomorphism*

$$(\pi_*\mathcal{F})_{\pi(y)} \cong \prod_{\gamma \in \Gamma/\Gamma_y} \mathcal{F}_{\gamma y}$$

of  $R$ -modules. In particular,  $\pi_*$  is an exact functor and for all  $i \geq 0$

$$H^i(\mathcal{C}, \mathcal{F}) \cong H^i(X, \pi_*\mathcal{F}).$$

Moreover, there is an isomorphism

$$(\pi_*\underline{V})_{\pi(y)} \cong \mathrm{Coind}_{\Gamma_y}^{\Gamma} V$$

of  $R[\Gamma]$ -modules.

**Proof.** This follows from the fact that the  $\Gamma$ -orbit of any  $x \in \mathcal{C}$  is a discrete set. □

**5.4 Corollary.** *Let  $\mathcal{F} \in \mathfrak{Sh}_{\mathcal{C}}(R - \mathfrak{Mod})$  and suppose  $H^i(X, \mathcal{F}) = 0$  for all  $i \geq 1$ . Then*

$$H^q(\Gamma, H^0(X, \mathcal{F})) \cong R^q(H^0(X, \cdot)^\Gamma)(\mathcal{F})$$

for all  $q \geq 0$  and, in particular,

$$H^q(\Gamma, V) \cong R^q(H^0(X, \cdot)^\Gamma)(\pi_* \underline{V}).$$

**Proof.** The assumptions mean that the spectral sequence 5.1 degenerates. In the special case this is true since  $H^i(X, \pi_* \underline{V}) = H^i(\mathcal{C}, \underline{V}) = 0$  for all  $i \geq 1$  by Lemma 5.3 and the fact that  $\mathcal{C}$  is simply connected and  $\underline{V}$  is constant.  $\square$

**5.5 Corollary.** *The stalk in  $x \in X$  of the group cohomology sheaf  $\mathcal{H}^q(\Gamma, \pi_* \underline{V})$  is  $H^q(\Gamma_y, V)$  for any  $y \in \mathcal{C}$  with  $\pi(y) = x$ . In particular,  $\mathcal{H}^q(\Gamma, \pi_* \underline{V})$  is a skyscraper sheaf on  $X$  for all  $q \geq 1$ .*

**Proof.** The group  $\Gamma$  is of type  $(\text{FP})_\infty$ . Indeed, by [Bieri], Propositions 2.13 we know that free products of groups of type  $(\text{FP})_\infty$  are of type  $(\text{FP})_\infty$ . As finite groups are clearly  $(\text{FP})_\infty$  (see [Bieri], Example 2.6), it follows that  $G$  is. Finally, by [Bieri], Proposition 2.5, subgroups of finite index in groups of type  $(\text{FP})_\infty$  are  $(\text{FP})_\infty$ , whence  $\Gamma$  is.

Thus, we may apply Proposition 5.2. The first statement now follows from Lemma 5.3 and Shapiro's lemma. The special case is a consequence of the fact that the non-trivially stabilised points of  $\mathcal{C}$  for the action of  $\Gamma$  are discrete.  $\square$

Let us note as a consequence of the case  $q = 0$  that the sheaf  $\pi_*(\underline{V})^\Gamma$  is locally constant on  $X$  if and only if  $V^{\Gamma_y} = V$  for all  $y \in \mathcal{C}$ .

**5.6 Lemma.** *The composition of the edge morphisms 5.3 for  $p \geq 1$*

$$H^p(\Gamma, V) \xrightarrow{\sim} R^p(H^0(X, \cdot)^\Gamma)(\pi_* \underline{V}) \rightarrow H^0(X, \mathcal{H}^p(\Gamma, \pi_* \underline{V}))$$

is the restriction map from the theory of group cohomology, when we identify  $H^0(X, \mathcal{H}^p(\Gamma, \pi_* \underline{V}))$  with  $\prod_{x \in X} H^p(\Gamma_{y_x}, V)$  for a choice of  $y_x \in \mathcal{C}$  with  $\pi(y_x) = x$ .

**Proof.** In Proposition 5.2 we have seen that the composition of the edge maps is sheafification. This means in the case of a skyscraper sheaf that it comes from the natural maps

$$V^\Gamma = (\pi_* \underline{V}(X))^\Gamma \rightarrow ((\pi_* \underline{V})_x)^\Gamma = V^{\Gamma_{y_x}}.$$

Hence, the map on the group cohomology is the restriction.  $\square$

We have now established the following theorem.

**5.7 Theorem.** *In Notation 2.1 with  $\mathcal{C} \in \{\mathbb{H}, \overline{\mathbb{H}}\}$  and  $X = \Gamma \backslash \mathcal{C} \in \{Y_\Gamma, X_\Gamma\}$ , there is a spectral sequence*

$$H^p(X, \mathcal{H}^q(\Gamma, \pi_* \underline{V})) \Rightarrow H^{p+q}(\Gamma, V),$$

in which the edge morphisms (for  $p \geq 1$ )

$$H^p(\Gamma, V) \rightarrow H^0(X, \mathcal{H}^p(\Gamma, \pi_* \underline{V}))$$

become the group theoretic restriction map under the identification

$$H^0(X, \mathcal{H}^q(\Gamma, \pi_* \underline{V})) \cong \prod_{x \in X} H^q(\Gamma_{y_x}, V)$$

for a choice of  $y_x \in \mathcal{C}$  with  $\pi(y_x) = x$ .

### Explicit description of certain cohomology groups on Hecke triangle surfaces

We keep assuming Notation 2.1. Let  $\mathcal{V}$  be a sheaf of  $R$ -modules on  $Y_\Gamma$ . From the Leray spectral sequence associated to  $j : Y_\Gamma \hookrightarrow X_\Gamma$  we get the exact sequence

$$0 \rightarrow H^1(X_\Gamma, j_* \mathcal{V}) \rightarrow H^1(Y_\Gamma, \mathcal{V}) \rightarrow H^0(X_\Gamma, R^1 j_* \mathcal{V}) \rightarrow H^2(X_\Gamma, j_* \mathcal{V}) \rightarrow H^2(Y_\Gamma, \mathcal{V}). \quad (5.4)$$

The *parabolic cohomology group* (for  $Y_\Gamma$  and  $\mathcal{V}$ ) is image of the map  $H_c^i(Y_\Gamma, \mathcal{V}) \rightarrow H^i(Y_\Gamma, \mathcal{V})$ . It is denoted by  $H_{\text{par}}^i(Y_\Gamma, \mathcal{V})$ . Moreover, we call  $H^0(X_\Gamma, R^1 j_* \mathcal{V})$  the *boundary cohomology group* (for  $Y_\Gamma$  and  $\mathcal{V}$ ).

**5.8 Proposition.** *There is the natural isomorphism of  $R$ -modules  $H_{\text{par}}^1(Y_\Gamma, \mathcal{V}) \cong H^1(X_\Gamma, j_* \mathcal{V})$ .*

**Proof.** We consider the exact sequence of sheaves on  $X_\Gamma$

$$0 \rightarrow j_! \mathcal{V} \rightarrow j_* \mathcal{V} \rightarrow C \rightarrow 0,$$

in which  $C$  is defined as the cokernel. It is a skyscraper sheaf, as it is only supported on the cusps. Hence,  $H^1(X_\Gamma, C) = 0$  and the long exact sequence associated to the short exact sequence of sheaves above yields that the upper map is surjective in the commutative diagram

$$\begin{array}{ccc} H_c^1(Y_\Gamma, \mathcal{V}) & \twoheadrightarrow & H^1(X_\Gamma, j_* \mathcal{V}) \\ & \searrow & \downarrow \\ & & H^1(Y_\Gamma, \mathcal{V}), \end{array}$$

in which the vertical map comes from the Leray sequence Equation 5.4. As it is injective, the proposition follows.  $\square$

We can now prove the principal result of this section.

**5.9 Theorem.** *Recall that we are assuming Notation 2.1. Let  $M$  denote the coinduced module  $\text{Coind}_\Gamma^G(V)$ . The following explicit descriptions hold:*

$$H^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \cong M / (M^{(\sigma)} + M^{(\tau)})$$

and

$$H_{\text{par}}^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \cong \ker \left( M / (M^{(\sigma)} + M^{(\tau)}) \xrightarrow{1-\sigma} M / (1-T)M \right).$$

**Proof.** We first use that any non-trivially stabilised point  $x$  of  $\overline{\mathbb{H}}$  is conjugate by some  $g \in G$  to either  $i$ ,  $\zeta_n$  or  $\infty$ . The respective stabiliser groups are  $G_i = \langle \sigma \rangle$ ,  $G_{\sigma\zeta_n} = \langle \tau \rangle$  and  $G_\infty = \langle T \rangle$ . Hence, from Mackey's formula (Corollary 2.5) we obtain

$$\prod_{x \in Y_\Gamma} H^p(\Gamma_{y_x}, V) \cong H^p(\langle \sigma \rangle, M) \oplus H^p(\langle \tau \rangle, M)$$

and

$$\prod_{x \in X_\Gamma} H^p(\Gamma_{y_x}, V) \cong H^p(\langle \sigma \rangle, M) \oplus H^p(\langle \tau \rangle, M) \oplus H^p(\langle T \rangle, M),$$

where again  $y_x \in \overline{\mathbb{H}}$  with  $\pi(y_x) = x$ .

From the spectral sequence from Theorem 5.7 we now obtain the exact sequences

$$0 \rightarrow H^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \rightarrow H^1(\Gamma, V) \rightarrow H^1(\langle \sigma \rangle, M) \oplus H^1(\langle \tau \rangle, M)$$

and

$$0 \rightarrow H^1(X_\Gamma, (\pi_* \underline{V})^\Gamma) \rightarrow H^1(\Gamma, V) \rightarrow H^1(\langle \sigma \rangle, M) \oplus H^1(\langle \tau \rangle, M) \oplus H^1(\langle T \rangle, M).$$

Using Proposition 5.8 and comparing with the Mayer-Vietoris exact sequence (Proposition 2.2) yields the theorem.  $\square$

**5.10 Remark.** *A study of the homology of modular curves (as Riemann surfaces) for a subgroup  $\Gamma \leq \mathrm{PSL}_2(\mathbb{Z}) =: G$  of finite index has been carried out by Merel in [MerelHecke], also in order to compute modular forms. His result [MerelHecke], Proposition 4,*

$$H_1(X_\Gamma, \mathrm{cusps}, R) \cong M / (M^{(\sigma)} + M^{(\tau)})$$

with  $M = \mathrm{Coind}_\Gamma^G(V)$  and  $\mathrm{cusps} = \Gamma \backslash G(\infty)$  is a special case of ours, as one can see as follows.

The general duality theorem (see [Dold], Proposition VIII.7.2; note that in this case Čech cohomology coincides with singular cohomology (see e.g. [Dold], Proposition VIII.6.12)) gives an isomorphism  $H_1(X_\Gamma, \mathrm{cusps}, R) \cong H^1(Y_\Gamma, R)$ , so that we may apply Theorem 5.9.

**5.11 Remark.** *The spectral sequence from Theorem 5.7 has a geometric interpretation in terms of analytic modular stacks (see [Thesis], II.3). Due to the lack of suitable references for stacks we avoid their use here and just state the result:*

*There is an analytic stack, denoted  $[\Gamma \backslash \mathbb{H}]$  together with a projection in the category of analytic stacks  $f : [\Gamma \backslash \mathbb{H}] \rightarrow Y_\Gamma$ . The projection map  $g : \mathbb{H} \rightarrow [\Gamma \backslash \mathbb{H}]$  allows one to define the sheaf  $(g_* \underline{V})^\Gamma$  on the analytic stacks, in analogy to the above treatment. The derived functor cohomology for the functor taking global sections coincides with that of group cohomology, i.e.*

$$H^i([\Gamma \backslash \mathbb{H}], (g_* \underline{V})^\Gamma) \cong H^i(\Gamma, V).$$

*The spectral sequence from Theorem 5.7 for  $\mathcal{C} = \mathbb{H}$  can be identified with the Leray spectral sequence associated to the projection map  $f$ .*

## 6 Computation of modular forms

In this section we sketch how the modular symbols formalism and also the presented cohomology groups can be used for the computation of modular forms.

### Comparison

We start by comparing the various objects discussed so far.

**6.1 Theorem.** *We assume Notation 2.1. The following sequences are exact:*

$$(a) \ 0 \rightarrow H^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \rightarrow H^1(\Gamma, V) \rightarrow \prod_{x \in Y_\Gamma} H^1(\Gamma_{y_x}, V),$$

$$(b) \ 0 \rightarrow H_{\text{par}}^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \rightarrow H_{\text{par}}^1(\Gamma, V) \rightarrow \prod_{x \in Y_\Gamma} H^1(\Gamma_{y_x}, V),$$

$$(c) \ \prod_{x \in Y_\Gamma} (V^{\Gamma_{y_x}} / N_{\Gamma_{y_x}} V) \rightarrow \mathcal{M}_R(\Gamma, V) \rightarrow H^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \rightarrow 0,$$

$$(d) \ \prod_{x \in Y_\Gamma} (V^{\Gamma_{y_x}} / N_{\Gamma_{y_x}} V) \rightarrow \mathcal{CM}_R(\Gamma, V) \rightarrow H_{\text{par}}^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \rightarrow 0,$$

where for all  $x \in Y_\Gamma$  we have chosen  $y_x \in \mathbb{H}$  such that  $\pi(y_x) = x$ .

**Proof.** This follows Theorems 5.7, 5.9 and 3.5, making an identification between the induced and the coinduced module.  $\square$

**6.2 Corollary.** *We use the Notation 2.1 with  $R = \mathbb{Z}$ . The  $\mathbb{Z}$ -modules  $H^1(\Gamma, V)$ ,  $H^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma)$  and  $\mathcal{M}_\mathbb{Z}(\Gamma, V)$  only differ by torsion. The same statement holds for the  $\mathbb{Z}$ -modules  $H_{\text{par}}^1(\Gamma, V)$ ,  $H_{\text{par}}^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma)$  and  $\mathcal{CM}_\mathbb{Z}(\Gamma, V)$ .*

**6.3 Corollary.** *We assume Notation 2.1 and suppose that the order of  $\Gamma_x$  is invertible in  $R$  for all  $x \in \mathbb{H}$ . Then there are isomorphisms*

$$H^1(\Gamma, V) \cong H^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \cong \mathcal{M}_R(\Gamma, V)$$

and

$$H_{\text{par}}^1(\Gamma, V) \cong H_{\text{par}}^1(Y_\Gamma, (\pi_* \underline{V})^\Gamma) \cong \mathcal{CM}_R(\Gamma, V).$$

The statements hold, in particular, for the group  $\Gamma_1(N)$  with  $N \geq 4$ .

**Proof.** This follows from Theorem 6.1. For the last part we use that under the condition  $N \geq 4$  all  $\Gamma_1(N)_x$  are trivial.  $\square$

We should not fail to recall the following well-known facts. The group  $\Gamma_0(N)/\langle \pm 1 \rangle$  can only contain stabiliser groups of order 2 or 3. It contains no stabiliser of even order if and only if  $N$  is divisible by a prime  $q$  which is 3 modulo 4 or by 4. Furthermore, it does not contain any stabiliser of order 3 if and only if  $N$  is divisible by a prime  $q$  which is 2 modulo 3 or by 9 (for a proof see e.g. [Thesis]). Let us also mention that there are techniques for computing the torsion of the objects above explicitly, see e.g. [Thesis], Proposition 2.4.8, and [W].

## Eichler-Shimura

The following theorem, originally by Eichler and Shimura, is valid for any Fuchsian group of the first kind with parabolic elements (see [Knopp], Theorem A, and the discussion there).

**6.4 Theorem.** *Assume Notation 2.1. Let  $k \geq 2$  be an integer and  $\mathbb{C}[X, Y]_{k-2}$  be the  $\mathbb{C}$ -vector space of homogeneous polynomials of degree  $k - 2$ . Then there is an isomorphism, the Eichler-Shimura isomorphism,*

$$H_{\text{par}}^1(\Gamma, \mathbb{C}[X, Y]_{k-2}) \cong S_k(\Gamma) \oplus \overline{S_k(\Gamma)},$$

where  $S_k(\Gamma)$  is the space of holomorphic weight  $k$  cusp forms for  $\Gamma$ .

## Computing modular forms

If the group in question is a congruence subgroup of  $\text{SL}_2(\mathbb{Z})$  there is a theory of Hecke operators on modular forms and on the cohomology groups studied above. The Eichler-Shimura isomorphism (Theorem 6.4) is compatible with the Hecke action (see [Diamond-Im], Theorem 12.2.2). This has the following well-known consequence.

**6.5 Corollary.** *Let  $k \geq 2$  and  $N \geq 1$  be integers and consider  $\Gamma = \Gamma_0(N)$  (or  $\Gamma = \Gamma_1(N)$ ).*

*The  $\mathbb{Z}$ -algebra  $\mathbb{T}$  generated by the Hecke operators inside the endomorphism ring of the  $\mathbb{C}$ -vector space of cusp forms  $S_k(\Gamma)$  of weight  $k$  for  $\Gamma$  equals the  $\mathbb{Z}$ -algebra generated by the Hecke operators  $T_1, T_2, \dots, T_B$  for  $B = \frac{Nk}{12} \prod_{l|N, l \text{ prime}} (1 + \frac{1}{l})$  (resp.  $B = \frac{N^2k}{24} \prod_{l|N, l \text{ prime}} (1 - \frac{1}{l^2})$ ) in the endomorphism ring of either of the  $\mathbb{Q}$ -vector spaces  $H_{\text{par}}^1(\Gamma, \mathbb{Q}[X, Y]_{k-2})$ ,  $\mathcal{CM}_{\mathbb{Q}}(\Gamma, \mathbb{Q}[X, Y]_{k-2})$  or  $H_{\text{par}}^1(Y_{\Gamma}, (\pi_* \mathbb{Q}[X, Y]_{k-2})^{\Gamma})$ . Here  $\mathbb{Q}[X, Y]_{k-2}$  denotes the homogeneous polynomials of degree  $k - 2$  over the rationals.*

**Proof.** Since the coefficients are in a field, the (parabolic) group cohomology of  $\Gamma$  is equal to that of  $\Gamma / (\langle \pm 1 \rangle \cap \Gamma)$ . It is clear that the  $\mathbb{C}$ -vector space  $H_{\text{par}}^1(\Gamma, \mathbb{C}[X, Y]_{k-2})$  contains the natural  $\mathbb{Q}$ -structure  $H_{\text{par}}^1(\Gamma, \mathbb{Q}[X, Y]_{k-2})$ . Any  $\mathbb{Q}$ -structure, however, gives an isomorphic Hecke algebra. Due to Corollary 6.2 any of the other two modules may be used, too. Finally, the Eichler-Shimura isomorphism (Theorem 6.4) implies that the Hecke algebra of  $H_{\text{par}}^1(\Gamma, \mathbb{C}[X, Y]_{k-2})$  is isomorphic to the Hecke algebra of  $S_k(\Gamma)$ . The bounds  $B$  are usually called *Sturm bounds*. For a proof that they suffice see e.g. [EdixJussieu], Proposition 4.2.  $\square$

Because of the isomorphism

$$\text{Hom}_{\mathbb{Z}}(\mathbb{T}, \mathbb{C}) \xrightarrow{\phi \mapsto \sum_{n \geq 1} \phi(T_n) q^n} S_k(\Gamma),$$

the knowledge of the Hecke algebra is equivalent to the knowledge of the modular forms. We use the Fourier expansions of the modular forms at the standard cusp and the usual convention  $q = e^{2\pi i \tau}$  for  $\tau \in \mathbb{H}$ .

The article [W] describes cases in which one may use  $H_{\text{par}}^1(\Gamma, \mathbb{F}[X, Y]_{k-2})$  with a finite fields  $\mathbb{F}$  if one wants to compute Katz modular forms over  $\overline{\mathbb{F}}$ .

## References

- [Bieri] R. Bieri. *Homological dimension of discrete groups*. Queen Mary College Mathematics Notes, London, 1976.
- [Brown] K. S. Brown. *Cohomology of groups*, Springer, New York, 1982.
- [Cremona] J. E. Cremona. *Algorithms for modular elliptic curves*. Second edition. Cambridge University Press, Cambridge, 1997.
- [Diamond-Im] F. Diamond and J. Im. *Modular forms and modular curves*, in *Seminar on Fermat's Last Theorem (Toronto, ON, 1993–1994)*, 39–133, Amer. Math. Soc., Providence, RI, 1995.
- [Dold] A. Dold. *Lectures on algebraic topology*, Springer, New York, 1972.
- [EdixJussieu] S. J. Edixhoven. *Comparison of integral structures on spaces of modular forms of weight two, and computation of spaces of forms mod 2 of weight one*. Accepted for publication in the Journal de l'Institut de Mathématiques de Jussieu.  
See: <http://arxiv.org/abs/math.NT/0312019>
- [Hartshorne] R. Hartshorne. *Algebraic geometry*, Springer, New York, 1977.
- [Knopp] M. I. Knopp. *Some new results on the Eichler cohomology of automorphic forms*. Bull. AMS **80(4)** (1974), 607–632.
- [Manin] Manin, Ju. I. *Parabolic points and zeta functions of modular curves*. Izv. Akad. Nauk SSSR Ser. Mat. **36** (1972), 19–66.
- [MerelHecke] L. Merel. *Opérateurs de Hecke pour  $\Gamma_0(N)$  et fractions continues*, Ann. Inst. Fourier (Grenoble) **41** (1991), no. 3, 519–537.
- [MerelUniversal] L. Merel. *Universal Fourier expansions of modular forms*, in *On Artin's conjecture for odd 2-dimensional representations*, 59–94, Lecture Notes in Math., 1585, Springer, Berlin, 1994.
- [Schmidt] T. A. Schmidt. *Rational Period Functions and Parabolic Cohomology*. Journal of Number Theory **57** (1996), 50–65.
- [Šokurov] V. V. Šokurov. *Shimura integrals of cusp forms*. Izv. Akad. Nauk SSSR Ser. Mat. **44** (1980), no. 3, 670–718, 720.
- [SteinThesis] W. A. Stein. *Explicit approaches to modular abelian varieties*, UC Berkeley, Ph.D. thesis.
- [SteinBook] W. A. Stein. *Algorithms For Computing With Modular Forms*, textbook in preparation.
- [Weibel] C. A. Weibel. *An introduction to homological algebra*, Cambridge Univ. Press, Cambridge, 1994.
- [Thesis] G. Wiese. *Modular Forms of Weight One Over Finite Fields*. Ph.D. thesis, Universiteit Leiden.

[W] G. Wiese. *On the faithfulness of the parabolic cohomology of modular curves as a Hecke module.*  
Preprint.

Gabor Wiese

NWFI - Mathematik

Universität Regensburg

D-93040 Regensburg

Deutschland

E-mail: [gabor@pratum.net](mailto:gabor@pratum.net)

Webpage: <http://homepages.uni-regensburg.de/~wig03492/>