

# Photochemical Redox Reaction of Aqueous Benzylpentacyanocobaltate(III)

ARND VOGLER and RENATE HIRSCHMANN

Institut für Chemie, Universität Regensburg

(Z. Naturforsch. **31b**, 1082-1085 [1976]; eingegangen am 28. April 1976)

Redox Reaction, Benzylpentacyanocobaltate(III), Benzyl Radicals, Photochemistry

Charge transfer excitation of aqueous benzylpentacyanocobaltate(III) led to a redox reaction with formation of  $\text{Co}(\text{CN})_5^{3-}$  and benzyl radicals in the primary photochemical step. In the absence of oxygen the benzyl radicals dimerized to bibenzyl. At 313 nm irradiating wavelength the quantum yield was  $0,13 \pm 0,02$ . In the presence of air oxygen was apparently inserted into the carbon-cobalt bond with the intermediate formation of  $[\text{Co}(\text{CN})_5\text{O}_2\text{CH}_2\text{C}_6\text{H}_5]^{3-}$ , which decomposed to give  $\text{Co}(\text{CN})_5\text{H}_2\text{O}^{2-}$  and benzaldehyde ( $\Phi = 0,15 \pm 0,02$ ). The significance of these results is discussed with regard to the photochemistry of acidopentacyanocobaltates(III) and organocobalamins.

## Introduction

In contrast to acidopentaammine complexes of Co(III) the corresponding acidopentacyanides undergo efficient photosubstitution reactions upon ligand field excitation<sup>1,2</sup>. On the other hand, CT (ligand to metal) excitation leads to redox reactions of the amines with high quantum yields while in the case of the pentacyano complexes the formation of stable redox products has not yet been observed<sup>2,3</sup>. However, it has been shown that CT excitation of acidopentacyanocobaltates(III) is followed by a redox reaction as primary photochemical step<sup>4,5</sup>. The radicals thus produced have been detected by flash photolysis. But a rapid back reaction seems to prevent the formation of stable products. This different photoredox behaviour of the ammine and cyano complexes has been related to the different stability of the metallo fragment, which is generated in the primary photolytic step<sup>3</sup>.  $\text{Co}(\text{NH}_3)_5^{2+}$  is extremely labile and decomposes in aqueous solution before a reoxidation by the ligand radical takes place. On the other hand  $\text{Co}(\text{CN})_5^{3-}$  is a stable complex and its reoxidation by the ligand radical is very efficient. However, under suitable conditions it should be possible to intercept the primary radicals. Stable redox products may then be formed.

Requests for reprints should be sent to Prof. Dr. ARND VOGLER, Institut für Chemie der Universität Regensburg, Universitätsstraße, D-8400 Regensburg.

We have explored this possibility and selected the complex ion benzylpentacyanocobaltate(III) for this study. Our expectation, that  $[\text{Co}(\text{CN})_5\text{CH}_2\text{C}_6\text{H}_5]^{3-}$  may undergo a photochemical redox reaction, was based on the observation that certain other cobalt(III) complexes containing a cobalt-carbon  $\sigma$ -bond, particularly organocobalamins and organocobaloximes, undergo a light-induced homolytic splitting of the cobalt-carbon bond<sup>7-11</sup>.

## Experimental

### Material

$\text{Na}_3[\text{Co}(\text{CN})_5\text{CH}_2\text{C}_6\text{H}_5] \cdot 2 \text{H}_2\text{O}$  was prepared according to a known procedure<sup>6</sup>. The compound was of good spectroscopic purity; its ultraviolet absorption spectrum, shown in Fig. 1, and its IR spectrum agreed well with that reported for an analyzed sample<sup>6</sup>.

### Photolysis procedures

The light source used was a 100 W Osram high-pressure mercury lamp. A Schott filter combination 313 limited the ultraviolet radiation emitted to the wavelength of the mercury line at 313 nm.

The photolyses were carried out at room temperature in 1 cm cylindrical spectrophotometer cells, using concentrations of complex such as to have essentially complete light absorption (about  $2 \times 10^{-4}$  M). Since thermal reactions were slow, corrections were not necessary. The total amount of photolysis was limited to less than 10% to exclude any possible secondary photolysis or inner filter effects. Absorbed light intensities were determined by ferrioxalate actinometry<sup>12</sup>.




Fig. 1. Absorption spectrum of  $[\text{Co}(\text{CN})_5\text{CH}_2\text{C}_6\text{H}_5]^{3-}$ .

### Measurements

Infrared spectra were obtained with a Perkin Elmer 325 IR spectrometer. Visible and ultraviolet spectra were measured with a Cary 14 recording spectrophotometer and a Zeiss PMQ II spectrophotometer for measurements at selected wavelengths. Organic photolysis products were identified by gas chromatography (Perkin Elmer F 20 and Varian 1860-42).


### Results

The electronic spectrum of aqueous




exhibits a single, very intense absorption maximum at 295 nm with  $\epsilon = 1,8 \times 10^4$  (Fig. 1). Upon irradiation into this band (313 nm irradiating wavelength)  $[\text{Co}(\text{CN})_5\text{CH}_2\text{C}_6\text{H}_5]^{3-}$ , dissolved in deaerated water, underwent a photochemical reaction as indicated by the decrease of the absorption band at 295 nm. Simultaneously a new band appeared at 970 nm indicating the formation of  $\text{Co}(\text{CN})_5^{3-}$ <sup>13</sup>. This band was not detected when the photolyzed solution was exposed to air. If the photolysis was carried out in strongly alkaline solution (1 M KOH) a new intense absorption band occurred at 311 nm when the photolyzed solution was saturated with oxygen. This band is characteristic for the superoxo complex  $[(\text{NC})_5\text{CoO}_2\text{Co}(\text{CN})_5]^{6-}$  which may be formed by aerial oxidation of  $\text{Co}(\text{CN})_5^{3-}$ <sup>14</sup>. In large scale runs the organic products of the photolysis carried out in neutral, deaerated solutions, were extracted with  $\text{CH}_2\text{Cl}_2$  and identified by gas chromatography. The predominant product (more than 95%) was bibenzyl.

On the basis of these results the photolysis took place according to




If light absorption by the starting complex was complete the reaction followed a zeroth order kinetics, since the decrease of the extinction at 295 nm was proportional to the time of irradiation. At 313 nm irradiating wavelength the quantum yield was determined by measuring the decrease of the extinction at 295 nm, taking into account that the reaction products have a negligible absorption at 295 nm. The quantum yield was  $\Phi = 0,13 \pm 0,02$ .

When the photolysis was carried out in water in the presence of air the 295 nm absorption of  $[\text{Co}(\text{CN})_5\text{CH}_2\text{C}_6\text{H}_5]^{3-}$  decreased again and the pH of the solution increased.  $\text{Co}(\text{CN})_5^{3-}$  was not formed as indicated by the spectrum of the photolyzed solution which did not show the absorption band at 970 nm. This observation is not surprising since  $\text{Co}(\text{CN})_5^{3-}$  would be oxidized immediately by oxygen with the formation of  $(\text{NC})_5\text{CoO}_2\text{Co}(\text{CN})_5^{6-}$ . This binuclear peroxo complex is stable only in alkaline solution but should decompose under the reaction conditions<sup>14-16</sup>. Hence the photolysis was carried out in an alkaline solution (1 M KOH) saturated with oxygen. However, neither  $[(\text{NC})_5\text{CoO}_2\text{Co}(\text{CN})_5]^{6-}$  nor the superoxo complex  $[(\text{NC})_5\text{CoO}_2\text{Co}(\text{CN})_5]^{5-}$ , which may be formed by further oxidation of the peroxo complex<sup>14</sup>, was formed according to the spectrum of the photolyzed solution, which did not exhibit the typical absorptions of both complexes at 327 nm or 311 nm<sup>14</sup>. The organic product of the photolysis in neutral solutions, mainly benzaldehyde, was extracted with  $\text{CH}_2\text{Cl}_2$  and identified by gas chromatography. These results are consistent with the assumption that the photolysis proceeds by an insertion of oxygen into the cobalt-carbon bond with the intermediate formation of a peroxo complex according to



This peroxo complex was apparently not stable under these conditions, but may decompose according to



Although no attempt was made to detect the complex  $[\text{Co}(\text{CN})_5\text{O}_2\text{CH}_2\text{C}_6\text{H}_5]^{3-}$ , which should have



apparent in the case of organocobalamins. The electronic spectra of cobalamins are determined by the very intense absorptions of the corrin ligand<sup>9</sup>. Any other bands as CT absorptions and LF bands which are of low intensity have not been identified and may be hidden under or obscured by the intense bands which have been assigned to spin-allowed  $\pi\pi^*$  transitions of the corrin ligand. Since the photolysis occurs upon light absorption of the corrin ligand one could assume that an excited state of the corrin ligand is responsible for the photo-reactivity. But there is no obvious explanation how an excited state of corrin may initiate a reaction in which the metal center and an axial ligand are involved while the corrin ligand does not participate in these reactions.

The photolysis of organocobaloximes proceeds apparently by the same mechanism as that of the corresponding cobalamin. On the basis of semi-empirical calculations it has been suggested that the reactive state is indeed of the CT type<sup>11</sup>. In the case of organocobalamins light absorption of the corrin ligand should then be followed by a radiationless deactivation to a low-lying CT state which initiates the redox reaction. It may be objected that such a CT transition should be seen in the absorption spectrum if it is indeed lower in energy than the lowest  $\pi\pi^*$  transition of the corrin ligand. However,

the electron distribution within this CT state should lead to large geometrical changes of the complex, particularly to an extension of the axial cobalt-carbon bond due to the Jahn-Teller distortion which occurs in low-spin Co(II) complexes<sup>23</sup>. As a consequence, although the Franck-Condon transition from the ground state to the CT state can occur at higher energies than that to the lowest  $\pi\pi^*$  state of the corrin ligand, the potential-energy surface of the CT state may cross that of the lowest  $\pi\pi^*$  state near the bottom of the latter and extend to lower energies than that of the  $\pi\pi^*$  state. This would provide a pathway for a radiationless transition from the  $\pi\pi^*$  state to the reactive CT state. A related suggestion has been made to explain the photochemical behaviour of  $\text{Co}(\text{NH}_3)_5\text{NO}_2^{2+}$ <sup>24, 25</sup>.

It has been recently shown that the photoaquation of cyanocobalamin originates from an excited LF state<sup>26</sup>. It seems that generally the photochemistry of cobalamins is closely related to that of other transition metal complexes and particularly to the photochemical reactions of pentacyano complexes of Co(III)<sup>26</sup>.

Support for this research by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie is gratefully acknowledged. We thank E. EIBLER for the analysis of organic products by gas chromatography.

- 1 E. ZINATO, "Concepts in Inorganic Photochemistry," Chapter 4, A. W. ADAMSON and P. D. FLEISCHAUER (eds.), Wiley, New York, N. Y. 1975.
- 2 V. BALZANI and V. CARASSITI, "Photochemistry of Coordination Compounds," Chapter 11, Academic Press, New York, N. Y. 1970.
- 3 J. F. ENDICOTT, "Concepts in Inorganic Photochemistry," Chapter 3, A. W. ADAMSON and P. D. FLEISCHAUER (eds.), Wiley, New York, N. Y. 1975.
- 4 G. FERRAUDI and J. F. ENDICOTT, *J. Amer. Chem. Soc.* **95**, 2371 [1973].
- 5 G. FERRAUDI and J. F. ENDICOTT, *Inorg. Chem.* **12**, 2389 [1973].
- 6 J. HALPERN and J. P. MAHER, *J. Amer. Chem. Soc.* **86**, 2311 [1964].
- 7 J. M. PRATT, *J. Chem. Soc.* **1964**, 5154.
- 8 J. M. PRATT and B. R. D. WHITEAR, *J. Chem. Soc. (A)* **1971**, 252.
- 9 J. M. PRATT, "Inorganic Chemistry of Vitamin B<sub>12</sub>," Academic Press, New York, 1972, and references cited therein.
- 10 G. COSTA, G. MESTRONI, and G. PELLIZER, *J. Organometal Chem.* **15**, 187 [1964].
- 11 G. N. SCHRAUZER, L. P. LEE, and J. W. SIBERT, *J. Amer. Chem. Soc.* **92**, 2997 [1970].
- 12 C. G. HATCHARD and C. A. PARKER, *Proc. Roy. Soc., Ser. A*, **235**, 518 [1956].
- 13 N. N. KING and M. E. WINFIELD, *J. Amer. Chem. Soc.* **83**, 3366 [1961].
- 14 J. H. BAYSTON, R. N. BEALE, N. KELSO KING, and M. E. WINFIELD, *Aust. J. Chem.* **16**, 954 [1963].
- 15 A. HAIM and W. K. WILMARTH, *J. Amer. Chem. Soc.* **83**, 509 [1961].
- 16 J. H. BAYSTON and M. E. WINFIELD, *J. Catalysis* **3**, 123 [1964].
- 17 C. FONTAINE, K. N. V. DUONG, C. MERIENNE, A. GAUDEMER, and C. GIANOTTI, *J. Organometal. Chem.* **38**, 167 [1972].
- 18 C. GIANOTTI, G. MERLE, and J. R. BOLTON, *J. Organometal. Chem.* **99**, 145 [1975], and references cited therein.
- 19 F. R. JENSEN and R. C. KISKIS, *J. Amer. Chem. Soc.* **97**, 5825 [1975].
- 20 V. M. MISKOWSKI and H. B. GRAY, *Inorg. Chem.* **14**, 401 [1975].
- 21 J. K. KOCHI and D. DAVIS, *J. Amer. Chem. Soc.* **86**, 5264 [1964].
- 22 J. NAGY, J. REFFY, A. KUSZMANN-BORBELY, and K. PALOSSY-BECKER, *J. Organometal. Chem.* **7**, 393 [1967].
- 23 M. D. GLICK, J. M. KUSZAJ, and J. F. ENDICOTT, *J. Amer. Chem. Soc.* **95**, 5097 [1973].
- 24 V. BALZANI, R. BALLARDINI, N. SABBATINI, and L. MOGGI, *Inorg. Chem.* **7**, 1398 [1968].
- 25 F. SCANDOLA, C. BARTOCCI, and M. A. SCANDOLA, *J. Phys. Chem.* **78**, 572 [1974].
- 26 A. VOGLER, R. HIRSCHMANN, H. OTTO, and H. KUNKELY, *Ber. Bunsenges. Phys. Chem.* **80**, 420 [1976].