



---

Extensions of profinite duality groups

Alexander Schmidt and Kay Wingberg

Preprint Nr. 25/2008

# Extensions of profinite duality groups

Alexander Schmidt and Kay Wingberg

September 25, 2008

Let  $G$  be a profinite group and let  $p$  be a prime number. By  $Mod_p(G)$  we denote the category of discrete  $p$ -primary  $G$ -modules. For  $A \in Mod_p(G)$  and  $i \geq 0$ , let

$$D_i(G, A) = \varinjlim_U H^i(U, A)^*,$$

where  $*$  is  $\text{Hom}(-, \mathbb{Q}_p/\mathbb{Z}_p)$ , the direct limit is taken over all open subgroups  $U$  of  $G$  and the transition maps are the duals of the corestriction maps.  $D_i(G, A)$  is a discrete  $G$ -module in a natural way. Assume that  $n = cd_p G$  is finite. Then the  $G$ -module

$$I(G) = \varinjlim_{\nu \in \mathbb{N}} D_n(G, \mathbb{Z}/p^\nu \mathbb{Z})$$

is called the **dualizing module** of  $G$  at  $p$ . Its importance lies in the functorial isomorphism

$$H^n(G, A)^* \cong \text{Hom}_G(A, I(G))$$

for all  $A \in Mod_p(G)$ . This isomorphism is induced by the cup-products ( $V \subseteq U$ )

$$H^n(G, A)^* \times_{p^\nu} A^U \longrightarrow H^n(V, \mathbb{Z}/p^\nu \mathbb{Z})^*, (\phi, a) \longmapsto \left( \alpha \mapsto \phi(\text{cor}_G^V(\alpha \cup a)) \right)$$

by passing to the limit over  $\nu$  and  $V$ , and then over  $U$ . The identity-map of  $I(G)$  gives rise to the homomorphism

$$\text{tr} : H^n(G, I(G)) \longrightarrow \mathbb{Q}_p/\mathbb{Z}_p,$$

called the **trace map**.

The profinite group  $G$  is called a **duality group at  $p$  of dimension  $n$**  if for all  $i \in \mathbb{Z}$  and all finite  $G$ -modules  $A \in Mod_p(G)$ , the cup-product and the trace map

$$H^i(G, \text{Hom}(A, I(G))) \times H^{n-i}(G, A) \xrightarrow{\cup} H^n(G, I(G)) \xrightarrow{tr} \mathbb{Q}_p/\mathbb{Z}_p$$

yield an isomorphism

$$H^i(G, \text{Hom}(A, I(G))) \cong H^{n-i}(G, A)^*.$$

**Remark:** In [Ve], J.-L. Verdier used the name **strict Cohen-Macaulay at  $p$**  for what we call a profinite duality group at  $p$  here. In [Pl], A. Pletch defined  $D_p^n$ -groups (and called them duality groups at  $p$  of dimension  $n$ ). The  $D_p^n$ -groups of Pletch are exactly the duality groups at  $p$  (in our sense) which, in addition, satisfy the following finiteness condition:

*FC(G, p):  $H^i(G, A)$  is finite for all finite  $A \in \text{Mod}_p(G)$  and for all  $i \geq 0$ .*

Since any finite, discrete  $G$ -module is trivialized by an open subgroup  $U$  of  $G$ , condition  $FC(G, p)$  can also be rephrased in the form:

*FC(G, p):  $H^i(U, \mathbb{Z}/p\mathbb{Z})$  is finite for all open subgroups  $U$  of  $G$  and all  $i \geq 0$ .*

By a duality theorem due to J. Tate, see [Ta] Thm. 3 or [Ve] Prop. 4.3 or [NSW] (3.4.6), a profinite group  $G$  of cohomological  $p$ -dimension  $n$  is a duality group at  $p$  if and only if

$$D_i(G, \mathbb{Z}/p\mathbb{Z}) = 0 \quad \text{for } 0 \leq i < n.$$

As a consequence we see that every open subgroup of a duality group at  $p$  is a duality group at  $p$  as well (of the same cohomological dimension), and if an open subgroup of  $G$  is a duality group at  $p$  and  $cd_p G < \infty$ , then  $G$  is a duality group at  $p$  of the same cohomological dimension (use [NSW] (3.3.5)(ii)). Furthermore, any profinite group of cohomological  $p$ -dimension 1 is a duality group at  $p$ .

We call a profinite group  $G$  **virtually a duality group at  $p$  of (virtual) dimension  $vcd_p G = n$**  if an open subgroup  $U$  of  $G$  is a duality group at  $p$  of dimension  $n$ .

The objective of this paper is to give a proof of Theorem 1 below, which states that the class of duality groups is closed under group extensions  $1 \rightarrow H \rightarrow G \rightarrow G/H \rightarrow 1$  if the kernel satisfies  $FC(H, p)$ . Weaker forms of Theorem 1 were first proved by A. Pletch (for  $D_p^n$ -groups, see [Pl]<sup>1</sup>) and by the second author (for Poincaré groups, see [Wi]).

<sup>1</sup>The proof given by Pletch in [Pl] is only correct for pro- $p$ -groups as the author assumes that finitely generated projective modules over the complete group ring  $\mathbb{Z}_p[[G]]$  are free.

**Theorem 1.** *Let*

$$1 \longrightarrow H \longrightarrow G \longrightarrow G/H \longrightarrow 1$$

*be an exact sequence of profinite groups such that condition  $FC(H, p)$  is satisfied. Then the following assertions hold:*

- (i) *If  $G$  is a duality group at  $p$ , then  $H$  is a duality group at  $p$  and  $G/H$  is virtually a duality group at  $p$ .*
- (ii) *If  $H$  and  $G/H$  are duality groups at  $p$ , then  $G$  is a duality group at  $p$ .*

*Moreover, in both cases we have:*

$$cd_p G = cd_p H + vcd_p G/H,$$

*and there is a canonical  $G$ -isomorphism*

$$I(G)^\vee \cong I(H)^\vee \hat{\otimes}_{\mathbb{Z}_p} I(G/H)^\vee,$$

*where  $^\vee$  is the Pontryagin dual and  $\hat{\otimes}_{\mathbb{Z}_p}$  is the tensor-product in the category of compact  $\mathbb{Z}_p$ -modules.*

**Remark:** The assumption  $FC(H, p)$  is necessary, as the following examples show:

1. Let  $G$  be the free pro- $p$ -group on two generators  $x, y$  and let  $H \subset G$  be the normal subgroup generated by  $x$ . Then  $H$  is free of infinite rank,  $G/H$  is free of rank one and  $1 \rightarrow H \rightarrow G \rightarrow G/H \rightarrow 1$  is an exact sequence in which all three groups are duality groups of dimension one.
2. Let  $D$  be a duality group at  $p$  of dimension 2,  $F$  a duality group at  $p$  of dimension 1 and  $G = F * D$  their free product. The kernel of the projection  $G \twoheadrightarrow D$  has cohomological  $p$ -dimension 1, hence is a duality group at  $p$  of dimension 1. The group  $G$  has cohomological  $p$ -dimension 2 but is not a duality group at  $p$ .

In the proof of Theorem 1, we make use of the following

**Proposition 2.** *Let*

$$1 \longrightarrow H \longrightarrow G \longrightarrow G/H \longrightarrow 1$$

*be an exact sequence of profinite groups. Assume that  $FC(H, p)$  holds. Then there is a spectral sequence of homological type*

$$E_{ij}^2 = D_i(G/H, \mathbb{Z}/p\mathbb{Z}) \otimes D_j(H, \mathbb{Z}/p\mathbb{Z}) \implies D_{i+j}(G, \mathbb{Z}/p\mathbb{Z}).$$

*Proof.* Let  $g$  run through the open normal subgroups of  $G$ . Then  $gH/H \cong g/g \cap H$  runs through the open normal subgroups of  $G/H$ . For a  $G$ -module  $A \in \text{Mod}_p(G)$ , we consider the Hochschild-Serre spectral sequence

$$E(g, g \cap H, A) : E_2^{ij}(g, g \cap H, A) = H^i(g/g \cap H, H^j(g \cap H, A)) \implies H^{i+j}(g, A).$$

If  $g' \subseteq g$  is another open normal subgroup of  $G$ , then the corestriction yields a morphism

$$\text{cor} : E(g', g' \cap H, A) \longrightarrow E(g, g \cap H, A)$$

of spectral sequences. The map

$$E_2^{ij}(g', g' \cap H, A) \longrightarrow E_2^{ij}(g, g \cap H, A)$$

is the composite of the maps

$$\begin{aligned} H^i(g'/g' \cap H, H^j(g' \cap H, A)) &\xrightarrow{\text{cor}_{g' \cap H}^{g' \cap H}} H^i(g'/g' \cap H, H^j(g \cap H, A)) \\ &\xrightarrow{\text{cor}_{g/g \cap H}^{g'/g' \cap H}} H^i(g/g \cap H, H^j(g \cap H, A)) \end{aligned}$$

and the map between the limit terms is the corestriction

$$\text{cor}_g^{g'} : H^{i+j}(g', A) \longrightarrow H^{i+j}(g, A).$$

For  $2 \leq r \leq \infty$  we set

$$E_{ij}^2 = D_{ij}^r(G, H, A) := \varinjlim_g E_r^{ij}(g, g \cap H, A)^*.$$

As taking duals and direct limits are exact operations, the terms  $D_{ij}^r(G, H, A)$ ,  $2 \leq r \leq \infty$ , establish a homological spectral sequence which converges to  $D_n(G, A)$ . If  $h$  runs through the open subgroups of  $H$  which are normal in  $G$ , then the cohomology groups  $H^j(h, A)$  are  $G$ -modules in a natural way. If  $g$  is open in  $G$  with  $g \cap H \subseteq h$ , then these groups are  $g/g \cap H$ -modules. We see that

$$D_{ij}^2(G, H, A) = \varinjlim_{\substack{h \subseteq H \\ h \trianglelefteq G}} \varinjlim_{\substack{g \subseteq G \\ g \cap H \subseteq h}} H^i(g/g \cap H, H^j(h, A))^*,$$

where for both limits the transition maps are (induced by)  $\text{cor}^*$ . In order to conclude the proof of the proposition, it remains to construct isomorphisms

$$D_{ij}^2(G, H, \mathbb{Z}/p\mathbb{Z}) \cong D_i(G/H, \mathbb{Z}/p\mathbb{Z}) \otimes D_j(H, \mathbb{Z}/p\mathbb{Z})$$

for all  $i$  and  $j$ . To this end note that all occurring abelian groups are  $\mathbb{F}_p$ -vector spaces, so that  $*$  is  $\text{Hom}(-, \mathbb{F}_p)$ . Further note that for vector spaces  $V, W$  over a field  $k$  the homomorphism

$$V^* \otimes W^* \longrightarrow (V \otimes W)^*, \quad \phi \otimes \psi \longmapsto (v \otimes w \mapsto \phi(v)\psi(w))$$

is an isomorphism provided that  $V$  or  $W$  is finite-dimensional. Let  $h$  be an open subgroup of  $H$  which is normal in  $G$  and let  $g' \subseteq g$  be open subgroups of  $G$  such that  $g$  acts trivially on the finite group  $H^j(h, \mathbb{Z}/p\mathbb{Z})$ . Then, by [NSW] (1.5.3)(iv), the diagram

$$\begin{array}{ccc} H^i(g'/g' \cap H, \mathbb{Z}/p\mathbb{Z}) \otimes H^j(h, \mathbb{Z}/p\mathbb{Z}) & \xrightarrow{\cong} & H^i(g'/g' \cap H, H^j(h, \mathbb{Z}/p\mathbb{Z})) \\ \downarrow \text{cor} \otimes \text{id} & & \downarrow \text{cor} \\ H^i(g/g \cap H, \mathbb{Z}/p\mathbb{Z}) \otimes H^j(h, \mathbb{Z}/p\mathbb{Z}) & \xrightarrow{\cong} & H^i(g/g \cap H, H^j(h, \mathbb{Z}/p\mathbb{Z})) \end{array}$$

commutes. For fixed  $h$ , we therefore obtain isomorphisms

$$\begin{aligned} D_i(G/H, \mathbb{Z}/p\mathbb{Z}) \otimes H^j(h, \mathbb{Z}/p\mathbb{Z})^* & \\ & \cong \left( \varinjlim_g H^i(g/g \cap H, \mathbb{Z}/p\mathbb{Z})^* \right) \otimes H^j(h, \mathbb{Z}/p\mathbb{Z})^* \\ & \cong \varinjlim_g H^i(g/g \cap H, \mathbb{Z}/p\mathbb{Z})^* \otimes H^j(h, \mathbb{Z}/p\mathbb{Z})^* \\ & \cong \varinjlim_g \left( H^i(g/g \cap H, \mathbb{Z}/p\mathbb{Z}) \otimes H^j(h, \mathbb{Z}/p\mathbb{Z}) \right)^* \\ & \cong \varinjlim_g H^i(g/g \cap H, H^j(h, \mathbb{Z}/p\mathbb{Z}))^*. \end{aligned}$$

Passing to the limit over  $h$ , we obtain the required isomorphism

$$D_i(G/H, \mathbb{Z}/p\mathbb{Z}) \otimes D_j(H, \mathbb{Z}/p\mathbb{Z}) \cong D_{ij}^2(G, H, \mathbb{Z}/p\mathbb{Z}).$$

□

**Corollary 3.** *Under the assumptions of Proposition 2, let  $i_0$  and  $j_0$  be the smallest integers such that  $D_{i_0}(G/H, \mathbb{Z}/p\mathbb{Z}) \neq 0$  and  $D_{j_0}(H, \mathbb{Z}/p\mathbb{Z}) \neq 0$ , respectively. Then  $D_{i_0+j_0}(G, \mathbb{Z}/p\mathbb{Z}) \neq 0$ .*

*Proof.* The spectral sequence constructed in Proposition 2 induces an isomorphism

$$D_{i_0+j_0}(G, \mathbb{Z}/p\mathbb{Z}) \cong D_{i_0}(G/H, \mathbb{Z}/p\mathbb{Z}) \otimes D_{j_0}(H, \mathbb{Z}/p\mathbb{Z}) \neq 0.$$

□

*Proof of Theorem 1.* Assume that  $G$  is a duality group at  $p$  of dimension  $d$ . Let  $cd_p H = m$  and  $n = d - m$ . Then there exists an open subgroup  $H_1$  of  $H$  such that  $H^m(H_1, \mathbb{Z}/p\mathbb{Z}) \neq 0$ . Let  $G_1$  be an open subgroup of  $G$  such that  $H_1 = G_1 \cap H$ . Then  $G_1$  is a duality group at  $p$  of dimension  $d$ ,  $cd_p H_1 = m$  and  $G_1/H_1$  is an open subgroup of  $G/H$ . We consider the exact sequence

$$1 \longrightarrow H_1 \longrightarrow G_1 \longrightarrow G_1/H_1 \longrightarrow 1.$$

As  $H^m(H_1, \mathbb{Z}/p\mathbb{Z})$  is finite and nonzero, we have  $vcd_p G_1/H_1 = n$ , see [NSW] (3.3.9). Furthermore,  $D_i(G_1, \mathbb{Z}/p\mathbb{Z}) = 0$ ,  $i < n + m$ . Using Corollary 3, we see that  $D_i(G_1/H_1, \mathbb{Z}/p\mathbb{Z}) = 0$  for all  $i < n$  and  $D_j(H_1, \mathbb{Z}/p\mathbb{Z}) = 0$  for all  $j < m$ . Thus  $G_1/H_1$ , hence  $G/H$ , is virtually a duality group at  $p$  of dimension  $n$ , and  $H_1$ , and so  $H$ , is a duality group at  $p$  of dimension  $m$ . This shows (i).

Assume now that  $H$  and  $G/H$  are duality groups at  $p$  of dimension  $m$  and  $n$ . Then,  $cd_p G = n + m$  by [NSW] (3.3.8), and in the spectral sequence of Proposition 2 we have  $E_{ij}^2 = 0$  for  $(i, j) \neq (n, m)$ . Hence  $D_r(G, \mathbb{Z}/p\mathbb{Z}) = 0$  for  $r \neq n + m$  showing that  $G$  is a duality group at  $p$  of dimension  $n + m$ .

In order to prove the assertion about the dualizing modules, let  $h$  run through all open subgroups of  $H$  which are normal in  $G$  and  $g$  runs through the open subgroups of  $G$ . Since  $m = cd_p H$ , the Hochschild-Serre spectral sequence induces isomorphisms

$$H^{m+n}(g, \mathbb{Z}/p^\nu\mathbb{Z}) \cong H^n(g/g \cap H, H^m(g \cap H, \mathbb{Z}/p^\nu\mathbb{Z})),$$

and we obtain

$$\begin{aligned} I(G) &\cong \varinjlim_{\nu} \varinjlim_g H^{m+n}(g, \mathbb{Z}/p^\nu\mathbb{Z})^* \\ &\cong \varinjlim_{\nu} \varinjlim_h \varinjlim_g H^n(g/g \cap H, H^m(h, \mathbb{Z}/p^\nu\mathbb{Z}))^* \\ &\cong \varinjlim_{\nu} \varinjlim_h \varinjlim_{g, \text{res}} H^0(g/g \cap H, \text{Hom}(H^m(h, \mathbb{Z}/p^\nu\mathbb{Z}), I(G/H))) \\ &\cong \varinjlim_{\nu} \varinjlim_h \text{Hom}(H^m(h, \mathbb{Z}/p^\nu\mathbb{Z}), I(G/H)) \\ &\cong \text{Hom}_{cts}(\varprojlim_{\nu} \varprojlim_h H^m(h, \mathbb{Z}/p^\nu\mathbb{Z}), I(G/H)) \\ &\cong \text{Hom}_{cts}((\varinjlim_{\nu} \varinjlim_h H^m(h, \mathbb{Z}/p^\nu\mathbb{Z})^*)^\vee, I(G/H)) \\ &\cong \text{Hom}_{cts}(I(H)^\vee, I(G/H)) \cong (I(H)^\vee \hat{\otimes}_{\mathbb{Z}_p} I(G/H)^\vee)^\vee \end{aligned}$$

(see [NSW] (5.2.9) for the last isomorphism). This completes the proof of the theorem.  $\square$

## References

- [NSW] Neukirch, J., Schmidt, A., Wingberg, K. *Cohomology of Number Fields*. sec.ed. Springer 2008
- [Pl] Pletch, A. *Profinite duality groups I*. J. Pure Applied Algebra **16** (1980) 55–74 and 285–297
- [Ta] Tate, J. *Letter to Serre*. Annexe 1 to Chap.I in Serre, J.-P. *Cohomologie Galoisienne*. Lecture Notes in Mathematics **5**, Springer 1964 (Cinquième édition 1994)
- [Ve] Verdier, J.-L. *Dualité dans la cohomologie des groupes profinis*. Annexe 2 to Chap.I in Serre, J.-P. *Cohomologie Galoisienne*. Lecture Notes in Mathematics **5**, Springer 1964 (Cinquième édition 1994)
- [Wi] Wingberg, K. *On Poincaré groups*. J. London Math. Soc. **33** (1986) 271–278

Alexander Schmidt, NWF I - Mathematik, Universität Regensburg, D-93040 Regensburg, Deutschland. email: alexander.schmidt@mathematik.uni-regensburg.de

Kay Wingberg, Mathematisches Institut, Universität Heidelberg, Im Neuenheimer Feld 288, 69120 Heidelberg, Deutschland. email: wingberg@mathi.uni-heidelberg.de