

# Release of glucose by *Nostoc spec.* isolated from the lichen *Peltigera horizontalis*

Doris Meindl and Eckhard Loos

Lehrstuhl für Zellbiologie und Pflanzenphysiologie, Universitätsstrasse 31, Universität Regensburg, D-8400 Regensburg (F.R.G.)

(Received March 23rd, 1990; revision received July 27th, 1990; accepted July 27th, 1990)

Cultured *Nostoc spec.* from *Peltigera horizontalis* excreted in the light between 1% and 3.4% of fixed  $^{14}\text{CO}_2$  into the medium,  $\leq 0.8\%$  as glucose. A several-fold increase in the rate of release was caused by addition of carbonyl cyanide *p*-trifluoromethoxyphenylhydrazone (FCCP) in the dark, the major component released being glucose. The amount of glucose excreted in the presence of FCCP exceeded the pool of intracellular glucose + glucose 6-phosphate by far, suggesting a glucan as a precursor. A glucan — identified enzymatically with amyloglucosidase — was the main assimilation product which was degraded during excretion of glucose in the dark in the presence of FCCP. Of the free-living cyanobacteria, *Nostoc punctiforme* also showed glucose release whereas *Nostoc muscorum* did not. It is concluded that (i) the symbiotic *Nostoc spec.* can excrete glucose in the absence of the fungal partner, (ii) this release is a passive process being fed by a glucan and (iii) glucose release is not a specific function of symbiotic cyanobacteria.

**Key words:** cyanobacteria (*Nostoc*); glucose transport; lichen (*Peltigera*); symbiosis

## Introduction

Glucose is believed to move in substantial amounts (> 33% of photosynthate) from *Nostoc spec.* to the fungal partner in the thallus of *Peltigera polydactyla* [1,2], a process which seems to be common to most cyanobacteria-containing lichens [3]. Only small amounts of labeled glucose, however, appeared in the medium when *Nostoc spec.* was allowed to assimilate  $^{14}\text{CO}_2$  immediately after isolation from *Peltigera canina* (3.4% of total label); 3 h after isolation, or upon culturing the cells, no radioactivity at all could be detected in this sugar [4, cf. 5]. Obviously, in the intact lichen, glucose excretion of the alga is induced somehow by the fungal partner [4]. As a starting point to unravel this fascinating regulatory phenomenon, the excretion characteristics of a cultured symbiotic cyanobacterium were studied. In

the present work the isolation and growth of *Nostoc spec.* from *Peltigera horizontalis* are briefly described and conditions for glucose release reported. Furthermore, two free-living cyanobacterial species were tested for glucose release, to see whether this is a specific feature of symbiotic blue-green algae.

## Material and Methods


### Plant material

*Nostoc spec.* was isolated from *Peltigera horizontalis* growing on woody cliffs 15 km southwest of Regensburg, F.R.G. Algal filaments were removed from thallus cuts with a fine needle, suspended in a mineral nutrient medium described by Matzke et al. [6] and sonicated in an ultrasonic bath to obtain short fragments. After repeated washings with sterile medium the cell material was plated on the same medium solidified with 1% agar. After incubation for 1—2 weeks at 28°C at 3000 lx colonies appeared which were transferred to fresh agar medium. Axenic colonies were identified by microscopic examination and by plating

**Correspondence to:** Eckhard Loos, Lehrstuhl für Zellbiologie und Pflanzenphysiologie, Universitätsstrasse 31, Universität Regensburg, D-8400 Regensburg, F.R.G.

Abbreviation: FCCP, carbonyl cyanide *p*-trifluoromethoxyphenyl hydrazone.






**Fig. 1.** Time course of  $^{14}\text{C}$ -labelling of cells and of excretion products of *Nostoc spec.*  $\blacktriangle$ — $\blacktriangle$ , radioactivity in cells;  $\bullet$ — $\bullet$ , radioactivity in the medium;  $\circ$ — $\circ$ , radioactivity of uncharged material in the medium. The initial rate of  $^{14}\text{CO}_2$  assimilation was  $47.6 \mu\text{mol} \cdot \text{h}^{-1} \cdot (\text{g wet wt.})^{-1}$  equivalent to  $122 \mu\text{mol} \cdot \text{h}^{-1} \cdot (\text{mg chlorophyll})^{-1}$ .

on *Nostoc* isolated from different species of *Peltigera* [11,12]. The growth proceeded logarithmically up to a density of 0.6 g/l dry weight with a doubling time of 1.33 days; in the stationary phase a maximum yield of 2.9 g/l dry weight was obtained.

#### Kinetics of $^{14}\text{CO}_2$ assimilation; analysis of excretion products; pH dependence of excretion


$^{14}\text{CO}_2$  fixation proceeded in a linear fashion for 1 h (Fig. 1); later it decreased, perhaps because of depletion of  $\text{CO}_2$ . Labeled products accumulated in the medium with a constant rate through 3 h; in several experiments, after 1 h of assimilation, between 1 and 3.4% of total fixed  $^{14}\text{C}$  were found to be released. About one third of the excreted radioactivity was bound to ion exchangers (Fig. 1). Thin-layer-chromatographic analysis of the remaining two thirds uncharged material yielded a complex pattern (Fig. 2a), generally with two conspicuous peaks, one at the origin and the other with a relative mobility similar to glucose. The ratio of the two was variable and seemed to be in favour of the immobile material with younger cultures.



**Fig. 2.** Scans for radioactivity of thin-layer chromatograms. (a) Excretion products of *Nostoc spec.* at pH 4.8; the medium was treated with mixed ion-exchangers before separation. (b) Analysis of the peak at the position of glucose in (a). The peak material was eluted and incubated in a total volume of 0.6 ml 17 mM Na-acetate buffer (pH 5.6) with 62 I.U. glucose oxidase (Boehringer, Mannheim, F.R.G.) for 90 min at 37°C, treated with cation exchanger (Serdolit CS-2, H<sup>+</sup>-form, Serva, Heidelberg, F.R.G.) and was chromatographed. The positions of reference substances are indicated: G, glucose; GA, gluconic acid; GL, gluconolactone; O, origin.

When the putative glucose peak was eluted and treated with glucose oxidase, the major products corresponded to gluconolactone and gluconic acid (Fig. 2b). This is strong evidence for glucose as an excretion product of the cultured *Nostoc spec.* The percentage of total fixed  $^{14}\text{CO}_2$  released as glucose was small, however, and varied between 0.04 and 0.8% after 1 h of assimilation in seven experiments. These results show that the photobiont is capable of releasing glucose also in the absence of the fungus, though in small amounts. Little excretion by cultured *Nostoc spec.* from *Peltigera polydactyla* has been observed also by Green and Smith [4]; they did not detect, however, glucose as an excretion product.

Buffer of pH 4.8 had been used in the preceding experiments, since an acidic pH was expected to favor sugar excretion as in the case of symbiotic *Chlorellae* [6,8,13]. Variation of the pH in the




**Fig. 3.** The effect of FCCP on release of radioactive material by cells of *Nostoc spec.* in the dark. After 1 h of  $^{14}\text{CO}_2$  assimilation (see Material and Methods) the cells were sedimented by centrifugation, resuspended in 0.5 ml fresh buffer, transferred to a darkened test tube with air as gas phase and were shaken. Every hour the medium was removed for counting and was replaced by fresh buffer. The cumulated counts are plotted in the graph. At the beginning of the dark phase FCCP was added in ethanolic solution (final concentration  $10^{-5}$  M), to the control ethanol only. The radioactivity recovered in the cells at the end of the dark phase was 256 100 cpm in the control and 231 960 cpm in the sample with FCCP.

range of pH 4.8–7.8, however, did not reveal a clear optimum, neither for total excreted material nor for excreted glucose (data not shown).

#### The influence of FCCP on glucose excretion

To see whether the release of glucose was energy-dependent, eventually due to an active



**Fig. 4.** Kinetics of glucose release from two *Nostoc* species in the presence (●—●) and absence (○—○) of FCCP. (a) Symbiotic *Nostoc spec.*; (b) *Nostoc punctiforme*. In a 100-ml Erlenmeyer flask, 600 mg filaments (wet weight) suspended in 10 ml of 50 mM citric acid-trisodium citrate buffer (pH 4.8) were shaken in the dark with air as gas phase. After every hour, the suspension was centrifuged, the supernatant was taken for the enzymatic glucose determination (see Material and Methods), and the pellet was resuspended in fresh buffer. FCCP dissolved in ethanol was included in the suspension buffer to give a final concentration of  $10^{-5}$  M; the control contained the corresponding amount of ethanol only. The cumulated amounts of glucose are plotted in the graph. The dashed line in (a) indicates the pool of intracellular glucose + glucose 6-phosphate, which was determined from cells taken at the beginning of the experiment (see Material and Methods).

**Table I.** Distribution of radioactivity after 1 h  $^{14}\text{CO}_2$  assimilation by *Nostoc* after a consecutive 4-h incubation of the cells in the dark in the presence of  $10^{-5}$  M FCCP. Filaments of 120 mg wet weight suspended in 2 ml 50 mM citric acid-trisodium citrate buffer (pH 4.8) were allowed to assimilate  $^{14}\text{CO}_2$  for 1 h (see Material and Methods). The suspension was then divided into two halves; in one half the distribution of label between medium and cell fractions was determined immediately, in the other half after shaking for 4 h in the dark in the presence of  $10^{-5}$  M FCCP (28°C, air as gas phase). Cell fractions were obtained as outlined in Material and Methods.

Fraction	Incubation condition	
	1 h light, $^{14}\text{CO}_2$ (cpm)	1 h light, $^{14}\text{CO}_2$ ; 4 h dark, air, $10^{-5}$ M FCCP (cpm)
Medium	4 050	92 540
Ethanol soluble	12 230	59 150
Ethanol insoluble	207 550	42 900
Degraded by amyloglucosidase	178 178	24 233

export step, experiments were carried out with the uncoupler FCCP. After 1 h of  $^{14}\text{CO}_2$  assimilation the filaments were transferred into fresh buffer and the excretion of labeled material was followed in the dark, in the presence and absence of FCCP. Surprisingly, the uncoupler caused a several-fold increase in excreted radioactivity (Fig. 3), a large part of which (56%) consisted of glucose as identified by thin-layer chromatography (cf. Fig. 2). In other experiments, unlabeled *Nostoc spec.* was incubated in the dark and extracellular glucose was determined enzymatically. In the presence of FCCP, glucose was released at a much higher rate as compared to the control (Fig. 4a). This effect was not specific for uncouplers like FCCP, since a several-fold stimulation of glucose efflux could be achieved also with  $\text{NaN}_3$ , an inhibitor of respiration.

These observations support the idea that the glucose transport step is not an active process, but rather a passive one, e.g. a carrier-mediated diffusion. Such a passive process was suggested from work with thallus disks of *Peltigera polydactyla*, in which dinitrophenol and sodium azide showed no effect on the transfer of algal photosynthate to the fungus [14; cf. 15]. In contrast, an energy requirement for the release of glucose, probably in the transport step proper, was indicated for a *Chlorella* symbiotic with a freshwater sponge [8].

Which pools are feeding the FCCP-induced glucose efflux? In the course of 4 h, 600 mg (wet weight) *Nostoc spec.* had excreted 0.46  $\mu\text{mol}$  of glucose (Fig. 4a), whereas the pool of cellular glu-

cose + glucose 6-phosphate at the beginning was only 0.022  $\mu\text{mol}$  (Fig. 4a, dotted line). From this it is evident that some glucose precursor has to be degraded, probably an intracellular glucan. Experimental evidence for this was obtained by determining enzymatically the amount of labeled glucan (i) after a period of  $^{14}\text{CO}_2$  assimilation and (ii) after an additional incubation in the dark in the presence of FCCP. Table I shows that after 1 h of  $^{14}\text{CO}_2$  assimilation most of the label was recovered in insoluble material. Of this, 86% were degraded to glucose by amyloglucosidase. The insoluble cell fraction, therefore, seems to consist largely of a starch- or glycogen-like polysaccharide [16]. At the end of the incubation in the dark, this labeled glucan had diminished drastically with a concomitant increase of radioactivity in the medium and in the soluble cellular material (Table I). Thin-layer chromatographic analysis revealed [ $^{14}\text{C}$ ]glucose as a major component in the medium (49% of the medium's radioactivity). Another experiment gave a similar result. Thus, a glucan seems to be the source of the excreted glucose. Evidence for a glucan as an intermediate stage in the flow of carbon from the lichenized *Nostoc* to the fungus has been presented by Hill [17,18].

The reason for the strong increase in glucose efflux caused by FCCP (Figs. 3 and 4a) is unclear. One could imagine that in *Nostoc spec.* a turnover of glucan occurs [17,18, cf. 19] and that this turnover is unbalanced. A lower rate of glucan synthesis might be caused by an increased level of phosphate in the uncoupled state, since phosphate

allosterically inhibits ADP-glucose pyrophosphorylase, the key enzyme of glucan biosynthesis [20,21]. Alternatively or in addition, the rate of glucan degradation might be increased by still unknown mechanisms [cf.22].

*Is the release of glucose specific for symbiotic Nostoc species?*

When two free-living species of cyanobacteria were tested for glucose release, one (*Nostoc muscorum*) was found to be incompetent and the other (*Nostoc punctiforme*) showed a release which could be stimulated by FCCP similar as in the symbiotic *Nostoc spec.*, at least during the first 2 h (Fig. 4b). The ability for glucose release, therefore, is not a special character of symbiotic strains of cyanobacteria. The close relationship between symbiotic *Nostoc spec.* and free-living *Nostoc punctiforme* already suggested on grounds of their morphology [cf. 23] and life cycles [12] is further underlined by their common capability for glucose release.

**Acknowledgement**

This work was supported by the Deutsche Forschungsgemeinschaft. Thanks are due to Dr. W. Lockau for critical reading of the manuscript.

**References**

- 1 E.A. Drew and D.C. Smith, Studies in the physiology of lichens VIII. Movement of glucose from alga to fungus during photosynthesis in the thallus of *Peltigera polydactyla*. *New Phytol.*, 66 (1967) 389—400.
- 2 D.J. Hill and D.C. Smith, Lichen physiology. XII. The "inhibition technique". *New Phytol.*, 71 (1972) 15—30.
- 3 D.H.S. Richardson, D.J. Hill and D.C. Smith, Lichen physiology. XI. The role of the alga in determining the patterns of carbohydrate movement between lichen symbionts. *New Phytol.*, 67 (1986) 469—486.
- 4 G.A. Green and D.C. Smith, Lichen physiology. XIV. Differences between lichen algae in symbiosis and in isolation. *New Phytol.*, 73 (1974) 753—766.
- 5 E.A. Drew and D.C. Smith, Studies in the physiology of lichens. VII. The physiology of the *Nostoc* symbiont of *Peltigera polydactyla* compared with cultured and free-living forms. *New Phytol.*, 66 (1967) 379—388.
- 6 B. Matzke, E. Schwarzmeier and E. Loos, Maltose excretion by the symbiotic *Chlorella* of the heliozoan *Acanthocystis turfacea*. *Planta*, 181 (1990) 593—598.
- 7 R.Y. Stanier, R. Kunisawa, M. Mandel and G. Cohen-Bazire, Purification and properties of unicellular blue-green algae (order *Chroococcales*). *Bacteriol. Rev.*, 35 (1971) 171—205.
- 8 A. Fischer, D. Meindl and E. Loos, Glucose excretion by the symbiotic *Chlorella* of *Spongilla fluviatilis*. *Planta*, 170 (1989) 251—256.
- 9 H.U. Bergmeyer, E. Bernt, F. Schmidt and H. Stork, D-Glucose:Bestimmung mit Hexokinase und Glucose-6-phosphat-Dehydrogenase, in: H.U. Bergmeyer (Ed.), *Methoden der Enzymatischen Analyse*, Vol. II, Verlag Chemie, 1974, pp. 1241—1246.
- 10 R.L. Bielecki, Sugar alcohols, in: F.A. Loewus and W. Tanner (Eds.), *Encyclopedia of Plant Physiology*, New Series, Vol. 13A, Springer-Verlag, 1982, pp. 158—192.
- 11 B. Bergmann and L. Hällbom, *Nostoc* of *Peltigera canina* when lichenized and isolated. *Can. J. Bot.*, 60 (1982) 2092—2098.
- 12 J.C. Boissière, N.C. Boissière, P. Champion-Arnaud and R. Lallemand, Le cycle des *Nostoc* des genres *Peltigera* et *Collema* en cultures in vitro et dans le thalle lichénique. *Can. J. Bot.*, 65 (1987) 1468—1477.
- 13 L. Muscatine, Symbiosis of hydra and algae. III. Extracellular products of the algae. *Comp. Biochem. Physiol.*, 16 (1965) 77—92.
- 14 D.C. Smith, Symbiosis and the biology of lichenised fungi, in *Symbiosis*, 29th Symp. Soc. Exp. Biol., Cambridge University Press, London, 1975, pp. 373—405.
- 15 S. Chambers, M. Morris and D.C. Smith, Lichen physiology. XV. The effect of digitonin and other treatments on biotrophic transport of glucose from alga to fungus in *Peltigera polydactyla*. *New Phytol.*, 76 (1976) 485—500.
- 16 J. H. Pazar and T. Ando, The action of an amyloglucosidase of *Aspergillus niger* on starch and maltose-oligosaccharides. *J. Biol. Chem.*, 234 (1959) 1966—1970.
- 17 D.J. Hill, The movement of carbohydrate from the alga to the fungus in the lichen *Peltigera polydactyla*. *New Phytol.*, 71 (1972) 31—39.
- 18 D.J. Hill, The physiology of lichen symbiosis, in: D.H. Brown, D.L. Hawksworth and R.H. Bailey (Eds.), *Lichenology: Progress and Problems*, Academic Press, 1976, pp. 457—496.
- 19 M. Lehmann and G. Wöber, Accumulation, mobilization and turnover of glycogen in the blue-green bacterium *Anacystis nidulans*. *Arch. Microbiol.*, 111 (1976) 93—97.
- 20 J. Preiss, H.P. Ghosh and J. Wittkop, Regulation of the biosynthesis of starch in spinach leaf chloroplasts, in: T.W. Goodwin (Ed.), *Biochemistry of Chloroplasts*, Vol. II, Academic Press, 1967, pp. 131—153.
- 21 C. Levi and J. Preiss, Regulatory properties of the ADP-glucose pyrophosphorylase of the blue-green bacterium *Synechococcus 6301*. *Plant Physiol.*, 58 (1976) 753—756.
- 22 A.J. Smith, Modes of cyanobacterial carbon metabolism, in: N.G. Carr and B.A. Whitton (Eds.), *The Biology of Cyanobacteria*, Botanical Monographs, Vol. 19, Blackwell Scientific Publications, 1982, pp. 47—85.
- 23 L. Geitler, Cyanophyceae, in: R. Kolkwitz (Ed.), Dr. L. Rabenhorst's Kryptogamenflora von Deutschland, Österreich und der Schweiz, Vol. 14, Akademische Verlagsgesellschaft m.b.H., 1932, pp. 1—1196.