



The conformal Yamabe constant of
product manifolds

Bernd Ammann, Mattias Dahl and Emmanuel Humbert

Preprint Nr. 17/2011

THE CONFORMAL YAMABE CONSTANT OF PRODUCT MANIFOLDS

BERND AMMANN, MATTIAS DAHL, AND EMMANUEL HUMBERT

ABSTRACT. Let (V, g) and (W, h) be compact Riemannian manifolds of dimension at least 3. We derive a lower bound for the conformal Yamabe constant of the product manifold $(V \times W, g + h)$ in terms of the conformal Yamabe constants of (V, g) and (W, h) .

CONTENTS

1.	Introduction	1
2.	Yamabe constants of product metrics	4
3.	Applications	7
	References	11

1. INTRODUCTION

1.1. The Yamabe functional, constant scalar curvature metrics, and Yamabe metrics. For a Riemannian manifold (M, G) we denote the scalar curvature by s^G , Laplace operator Δ^G , volume form dv^G . In general the dependence on the Riemannian metric is denoted by the metric as a superscript.

For integers $m \geq 3$ we set $a_m := \frac{4(m-1)}{m-2}$ and $p_m := \frac{2m}{m-2}$. Let $C_c^\infty(M)$ denote the space of compactly supported smooth functions on M . For a Riemannian manifold (M, G) of dimension $m \geq 3$ we define the Yamabe functional by

$$\mathcal{F}^G(u) := \frac{\int_M (a_m |du|_G^2 + s^G u^2) dv^G}{\left(\int_M |u|^{p_m} dv^G\right)^{\frac{2}{p_m}}},$$

where $u \in C_c^\infty(M)$ does not vanish identically. The *conformal Yamabe constant* $\mu(M, G)$ of (M, G) is defined by

$$\mu(M, G) := \inf_{u \in C_c^\infty(M), u \neq 0} \mathcal{F}^G(u).$$

The conformal Yamabe constant is usually defined only for compact manifolds, here we allow also non-compact manifolds in the definition. This will turn out to be essential for studying surgery formulas for Yamabe invariants of compact manifolds, see Subsection 3.2. Also notice that the conformal Yamabe constant for non-compact manifolds has been studied for instance in [11] and [9].

For compact M one easily sees that $\lim_{\varepsilon \rightarrow 0} \mathcal{F}^G(\sqrt{u^2 + \varepsilon^2}) = \mathcal{F}^G(u)$, thus we obtain

$$\mu(M, G) = \inf_{u \in C_+^\infty(M)} \mathcal{F}^G(u) > -\infty,$$

Date: March 10, 2011.

2000 Mathematics Subject Classification. 35J60 (Primary), 35P30, 58J50, 58C40 (Secondary).

Key words and phrases. Yamabe constant, Yamabe invariant, product manifolds.

where $C_+^\infty(M)$ denotes the space of positive smooth functions. According to the resolution of the Yamabe problem [20, 4, 18], see for example [12] for a good overview article, this infimum is always attained by a positive smooth function if M is a compact manifold.

For a compact manifold M one also defines for any metric G the (normalized) Einstein-Hilbert functional \mathcal{E} as

$$\mathcal{E}(G) := \frac{\int_M s^G dv^G}{\text{vol}^G(M)^{\frac{m-2}{m}}}.$$

These functionals are closely related to each other, namely if $u > 0$ and $\tilde{G} = u^{4/(m-2)}G$, then

$$\mathcal{E}(\tilde{G}) = \mathcal{F}^G(u).$$

From the discussion above it follows that the functional \mathcal{E} always attains its infimum in each conformal class $[G]$. Such minimizing metrics are called *Yamabe metrics*. Obviously \tilde{G} is a Yamabe metric if and only if $\lambda\tilde{G}$ is a Yamabe metric for any $\lambda > 0$. Thus any conformal class on a compact manifold carries a Yamabe metric of volume 1. Yamabe metrics \tilde{G} are stationary points of \mathcal{E} , restricted to the conformal class, and thus satisfy an Euler-Lagrange equation. This Euler-Lagrange equation says precisely that the scalar curvature of \tilde{G} is constant. One also sees that $\mu(M, G)$ is positive if and only if $[G]$ contains a metric of positive scalar curvature.

We denote the standard flat metric on \mathbb{R}^m by ξ^m . On the sphere $S^m \subset \mathbb{R}^{m+1}$ the standard round metric ρ^m is a Yamabe metric, and the whole orbit of ρ^m under the action of the Möbius group $\text{Conf}(S^m) = \text{PSO}(m+1, 1)$ consists of Yamabe metrics. Thus $\mathbb{S}^m := (S^m, \rho^m)$ carries a non-compact space of Yamabe metrics of volume 1.

In contrast to this, there is only one metric of constant scalar curvature and of volume 1 in the conformal class $[G]$, if at least one of the following conditions is satisfied.

- M is compact and $\mu(M, G) \leq 0$. The unicity then follows from the maximum principle.
- (M, G) is a connected compact Einstein manifold, and (M, G) is non-isometric to $(S^m, \lambda\rho^m)$ for any $\lambda > 0$. This case is one of Obata's theorems [13, Prop. 6.2].
- (M, G) is close in the $C^{2,\alpha}$ -topology to such an Einstein metric, see [8, Theorem C].

In particular in these cases there is exactly one Yamabe metric of volume 1, and any metric of constant scalar curvature is a Yamabe metric. In $[\rho^m]$ any metric of constant scalar curvature κ is in the orbit of the Möbius group acting on $\lambda\rho^m$, where $\kappa = m(m-1)/\lambda$. As a consequence, on the round sphere any constant scalar curvature metric is a Yamabe metric as well.

However, in general, the functionals $\mathcal{E}|_{[G]}$ and $\mathcal{F}^G|_{C_+^\infty(M)}$ may have non-minimizing stationary points. These stationary points are thus metrics of constant scalar curvature which are not Yamabe metrics. The simplest such example, extensively discussed by Schoen [19] for $w = 1$, is the metric $G = \rho^v + \lambda\rho^w$ on $S^v \times S^w$, $v \geq 2$, which has constant scalar curvature $v(v-1) + w(w-1)$, but which is not a Yamabe metric for sufficiently large λ . This is due to the fact that $\mu(M, G) \leq \mu(\mathbb{S}^m)$, which follows from a standard test function argument, whereas $\mathcal{E}(\rho^v + \lambda\rho^w) \rightarrow \infty$ as $\lambda \rightarrow \infty$ when $v \geq 2$.

In conclusion, if (M, G) is an explicitly given compact manifold of constant scalar curvature, the calculation of $\mu(M, G)$ is easy if either (M, G) is Einstein or if $\mu(M, G) \leq 0$, but in general it can be a hard problem.

$\varepsilon_{v,w}$	w=3	w=4	w=5	w=6	w=7
v=3	0.625	0.7072..	0.7515..	0.7817..	0.8042..
4	0.7072..	0.7777..	0.8007..	0.8367..	0.8537..
5	0.7515..	0.8007..	0.8427..	0.8631..	0.8772..
6	0.7817..	0.8367..	0.8631..	0.88	0.8921..
7	0.8042..	0.8537..	0.8772..	0.8921..	0.9027..

FIGURE 1. Values of $\varepsilon_{v,w}$

1.2. Product manifolds. We now consider Riemannian product manifolds, that is for Riemannian manifolds (V, g) and (W, h) of dimensions v and w , we equip $M = V \times W$ with the product metric $G = g + h$, or more generally $G = g + \lambda h$ where $\lambda > 0$. We ask the following question.

Question. *Suppose V and W are compact and equipped with Yamabe metrics g and h . Let $\lambda > 0$. Is then $g + \lambda h$ also a Yamabe metric?*

From the discussion on unicity above it follows that the answer is yes,

- if $v, w \geq 3$, $\mu(V, g) \leq 0$ and $\mu(W, h) \leq 0$;
- or if $v, w \geq 3$, $\mu(V, g) > 0$ and $\mu(W, h) < 0$ for $\lambda > 0$ small enough;
- or if (V, g) and (W, h) are both Einstein with $\frac{1}{v}s^g$ close to $\frac{1}{\lambda w}s^h$.

If the answer to the above question is yes, then one deduces

$$\mu(V \times W, g + \lambda h) = \left(\frac{\mu(V, g)}{\text{vol}^g(V)^{2/v}} + \frac{\mu(W, h)}{\text{vol}^h(W)^{2/w}} \right) \left(\text{vol}^g(V) \text{vol}^h(W) \right)^{\frac{2}{v+w}}. \quad (1)$$

On the other hand if g has positive scalar curvature, then $\mathcal{E}(g + \lambda h) \rightarrow \infty$ for $\lambda \rightarrow \infty$, thus $g + \lambda h$ is not a Yamabe metric for large λ . This applies, in particular, to the cases $\mu(V, g) > 0$, $v \geq 3$, or if $(V, g) = (S^2, \rho^2)$.

1.3. An intuitive—but incorrect—argument in the positive case. Now we assume $v, w \geq 3$, $\mu(V, g) > 0$, and $\mu(W, h) > 0$. We already explained why $g + \lambda h$ is not a Yamabe metric for large (and small) $\lambda > 0$, as a consequence Equation (1) cannot be true for all $\lambda > 0$. Despite of this fact, assume for a moment that (1) were true for all $\lambda > 0$. We then could minimize over λ , and we would obtain

$$\inf_{\lambda \in (0, \infty)} \mu(V \times W, g + \lambda h) = (v + w) \left(\frac{\mu(V, g)}{v} \right)^{\frac{v}{v+w}} \left(\frac{\mu(W, h)}{w} \right)^{\frac{w}{v+w}} \quad (2)$$

1.4. Main result. Although the *naive derivation* of formula (2) used incorrect assumptions, our main result, Theorem 2.3 will tell us that the *formula itself* is correct up to a factor

$$\varepsilon_{v,w} = \frac{a_{v+w}}{a_v^{v/(v+w)} a_w^{w/(v+w)}} < 1.$$

assuming the mild condition (4).

More precisely, we assume that V and W are compact manifolds of dimension at least 3, with Yamabe metrics g and h of positive conformal Yamabe constant. In particular, condition (4) is satisfied. Then Theorem 2.3 implies

$$\varepsilon_{v,w} \leq \frac{\inf_{\lambda \in (0, \infty)} \mu(V \times W, g + \lambda h)}{(v + w) \left(\frac{\mu(V, g)}{v} \right)^{\frac{v}{v+w}} \left(\frac{\mu(W, h)}{w} \right)^{\frac{w}{v+w}}} \leq 1.$$

Note that $\varepsilon_{v,w} \rightarrow 1$ for $v, w \rightarrow \infty$. See Figure 1 for some values of $\varepsilon_{v,w}$.

The main theorem also applies to many non-compact manifolds, see Theorem 2.3.

1.5. Further comments on related literature. Our main motivation to study Yamabe constants of products is the application sketched in Subsection 3.2.

Fundamental results on Yamabe constants on products have been found in the interesting article [1] where it is, in particular, shown that the conformal Yamabe constant of the product $V \times \mathbb{R}^w$ is a lower bound for $\sigma(V \times W)$. This article also emphasized the importance of the question under which conditions a function $u \in C^\infty(V \times W)$ minimizing \mathcal{E} is a function of only one of the factors. If V is compact and of constant scalar curvature 1, it was shown that the conformal Yamabe constant of manifolds $V \times \mathbb{R}^w$ is up to a constant the inverse of an optimal constant in a Gagliardo-Nirenberg type estimate.

In related research Petean [15] derived a lower bound for the conformal Yamabe constant of product manifolds $V \times \mathbb{R}$, where V is compact of positive Ricci curvature. If additionally we require V to be Einstein, any minimizer $u \in C^\infty(V \times \mathbb{R})$ of \mathcal{E} only depends on \mathbb{R} . As a corollary Petean obtained lower bounds for the smooth Yamabe invariant $\sigma(V \times S^1)$ in this case.

This result of Petean contrasts nicely to Theorem 2.3. Whereas Petean's result requires that one of the factor is 1-dimensional, our Theorem 2.3 requires both factors to be of dimension at least 3.

In [17] an explicit lower bound for $\mu(S^2 \times \mathbb{R}^2, \rho^2 + \xi^2)$ is obtained: $\mu(S^2 \times \mathbb{R}^2, \rho^2 + \xi^2) \geq 0.68 \cdot Y(S^4)$. A similar, but weaker result was obtained in [14].

Several recent publications study multiplicity phenomena on products $S^v \times W$ equipped with product metric of the standard metric on S^v with a metric of constant scalar curvature $s > 0$ on W . Explicit lower bounds for the number of metrics of constant scalar curvature 1 in the conformal class $[g_0]$ are derived, and these bounds grow linearly in \sqrt{s} . The case $v = 1$ was studied in [7, 6], the general case then treated in [16]. In the recent preprint [10] isoparametric hypersurfaces are used in order to obtain new metrics of constant scalar curvature in the conformal class of products of riemannian manifolds, e.g. the conformal class of $(S^3 \times S^3, \rho^3 + \lambda\rho^3)$.

1.6. Structure of the present article. In Section 2 we derive the main techniques and the main result of the article. We use mixed $L^{p,q}$ -spaces in order to obtain a lower bound of the conformal Yamabe constants in the case that both factors have dimension at least 3. We start with a proof of an iterated Hölder inequality in Subsection 2.1 which is well-adapted for the proof of our product formula in Subsection 2.3 which is the main result of the article.

In Section 3 we discuss applications. In Subsection 3.1 we find an estimate for the smooth Yamabe invariant of product manifolds. Subsection 3.2 explains our original motivation for the subject, which is to find better estimates for the constants appearing in the surgery formula in [2]. In Subsection 3.3 we define a stable Yamabe invariant and show that a similar surgery formula as in the unstable situation holds true.

Acknowledgements. We want to thank the organizers of the Conference “Contributions in Differential Geometry – a round table in occasion of the 65th birthday of Lionel Bérard Bergery, 2010”, in particular thanks to A. Besse, A. Altomani, T. Krantz and M.-A. Lawn. During that conference we found central ingredients for the present article.

B. Ammann was partially supported by DFG Sachbeihilfe AM 144/2-1. E. Humbert was partially supported by ANR-10-BLAN 0105.

2.1. Iterated Hölder inequality for product manifolds. Let (V, g) and (W, h) be Riemannian manifolds of dimensions $v := \dim V$ and $w := \dim W$. We set

$$(M, G) := (V \times W, g + h),$$

so that $m := \dim M = v + w$. We do not assume that the manifolds are complete. The first result we will need is a kind of iterated Hölder inequality for $(M, G) := (V \times W, g + h)$.

Lemma 2.1. *For any function $u \in C_c^\infty(M)$ we have*

$$\left(\int_M |u|^{p_m} dv^G \right)^{\frac{2}{p_m}} \leq \left(\int_V \left(\int_W |u|^{p_w} dv^h \right)^{\frac{2}{p_w}} dv^g \right)^{\frac{w}{m}} \left(\int_V \left(\int_W |u|^2 dv^h \right)^{\frac{p_v}{2}} dv^g \right)^{\frac{v-2}{m}}.$$

The lemma is actually a special case of the Hölder inequality for mixed $L^{p,q}$ -spaces. See [5] for further information on such spaces.

Proof. By the Hölder inequality we have

$$\int_W |u|^{p_m} dv^h \leq \left(\int_W |u|^{p_w} dv^h \right)^{\frac{w-2}{m-2}} \left(\int_W |u|^2 dv^h \right)^{\frac{v}{m-2}}.$$

We integrate this inequality over (V, g) , and use the following Hölder inequality

$$\int_V \alpha \beta dv^g \leq \left(\int_V |\alpha|^{\frac{m-2}{w}} dv^g \right)^{\frac{w}{m-2}} \left(\int_V |\beta|^{\frac{m-2}{v-2}} dv^g \right)^{\frac{v-2}{m-2}}$$

with

$$\alpha := \left(\int_W |u|^{p_w} dv^h \right)^{\frac{w-2}{m-2}} \quad \text{and} \quad \beta := \left(\int_W |u|^2 dv^h \right)^{\frac{v}{m-2}}.$$

This proves Lemma 2.1 □

2.2. A Lemma about integration and derivation. Second we need a Lemma concerning the interchange of derivation and taking (partial) L^2 -norm.

Lemma 2.2. *Let $u \in C_c^\infty(M)$, $u \not\equiv 0$, and set*

$$\gamma := \left(\int_W u^2 dv^h \right)^{\frac{1}{2}}.$$

Then

$$\int_V |d\gamma|_g^2 dv^g \leq \int_M |du|_g^2 dv^G. \quad (3)$$

Proof. Take any vector field X on M tangent to V . One has g -almost everywhere (except on the boundary of $\gamma^{-1}(0)$)

$$|X\gamma|^2 \leq \left(\frac{\int_W uXu dv^h}{\left(\int_W u^2 dv^h \right)^{\frac{1}{2}}} \right)^2 \leq \int_W (Xu)^2 dv^h,$$

where we used the Cauchy-Schwartz inequality

$$\int_W uXu dv^h \leq \left(\int_W (Xu)^2 dv^h \right)^{\frac{1}{2}} \left(\int_W u^2 dv^h \right)^{\frac{1}{2}}.$$

Integrating over V , we deduce that

$$\int_V |X\gamma|^2 dv^g \leq \int_M |Xu|^2 dv^G.$$

Since this holds for any X tangent to V , inequality (3) follows. □

2.3. Conformal Yamabe constant of product metrics. We now state and prove our main theorem. It will turn out that the following modified invariant is convenient when studying products of Riemannian manifolds with non-negative Yamabe constant. If $\mu(M, G) \geq 0$ we set

$$\nu(M, G) := \left(\frac{\mu(M, G)}{ma_m} \right)^m.$$

Theorem 2.3. *Let (V, g) and (W, h) be Riemannian manifolds of dimensions $v, w \geq 3$. Assume that $\mu(V, g), \mu(W, h) \geq 0$ and that*

$$\frac{s^g + s^h}{a_m} \geq \frac{s^g}{a_v} + \frac{s^h}{a_w}. \quad (4)$$

Then,

$$\mu(M, G) \geq \frac{ma_m}{(va_v)^{\frac{v}{m}}(wa_w)^{\frac{w}{m}}} \mu(V, g)^{\frac{v}{m}} \mu(W, h)^{\frac{w}{m}},$$

or, equivalently,

$$\nu(M, G) \geq \nu(V, g)\nu(W, h).$$

Note that we do not assume that the manifolds are complete.

Proof. Take any non-negative function $u \in C_c^\infty(M)$ normalized by

$$\int_M |u|^{p_m} dv^G = 1. \quad (5)$$

We then have

$$\frac{1}{a_m} \mathcal{F}^G(u) = \int_M \left(|du|_G^2 + \frac{s^G}{a_m} u^2 \right) dv^G.$$

Using $|du|_G^2 = |du|_g^2 + |du|_h^2$ and $s^G = s^g + s^h$ together with (4) we obtain

$$\frac{1}{a_m} \mathcal{F}^G(u) \geq \int_M \left(|du|_g^2 + \frac{s^g}{a_v} u^2 \right) dv^G + \int_V \int_W \left(|du|_h^2 + \frac{s^h}{a_w} u^2 \right) dv^h dv^g. \quad (6)$$

We set $\gamma := \left(\int_W u^2 dv^h \right)^{\frac{1}{2}}$. For the first term here, Lemma 2.2 and the definition of $\mu(V, g)$ imply that

$$\begin{aligned} \int_M \left(|du|_g^2 + \frac{s^g}{a_v} u^2 \right) dv^G &\geq \int_V \left(|d\gamma|_g^2 + \frac{s^g}{a_v} \gamma^2 \right) dv^g \\ &\geq \frac{1}{a_v} \mu(V, g) \left(\int_V \gamma^{p_v} dv^g \right)^{\frac{2}{p_v}} \\ &= \frac{1}{a_v} \mu(V, g) \left(\int_V \left(\int_W |u|^2 dv^h \right)^{\frac{p_v}{2}} dv^g \right)^{\frac{v-2}{v}}. \end{aligned} \quad (7)$$

For the second term we have

$$\int_V \int_W \left(|du|_h^2 + \frac{s^h}{a_w} u^2 \right) dv^h dv^g \geq \frac{1}{a_w} \mu(W, h) \int_V \left(\int_W u^{p_w} dv^h \right)^{\frac{2}{p_w}} dv^g \quad (8)$$

by the definition of $\mu(W, h)$. Plugging (7) and (8) in (6) we get

$$\begin{aligned} \mathcal{F}^G(u) &\geq \frac{a_m}{a_v} \mu(V, g) \left(\int_V \left(\int_W |u|^2 dv^h \right)^{\frac{p_v}{2}} dv^g \right)^{\frac{v-2}{v}} \\ &\quad + \frac{a_m}{a_w} \mu(W, h) \int_V \left(\int_W u^{p_w} dv^h \right)^{\frac{2}{p_w}} dv^g \end{aligned} \quad (9)$$

Set

$$r := ma_m \nu(V, g)^{\frac{1}{m}} \nu(W, h)^{\frac{1}{m}}$$

For $a, b > 0$ we compute

$$\begin{aligned}
ra^{\frac{v-2}{m}}b^{\frac{w}{m}} &= r \left(\left(\frac{\nu(V, g)^w}{\nu(W, h)^v} \right)^{\frac{1}{m^2}} a^{\frac{v-2}{m}} \right) \left(\left(\frac{\nu(W, h)^v}{\nu(V, g)^w} \right)^{\frac{1}{m^2}} b^{\frac{w}{m}} \right) \\
&\leq r \left[\frac{v}{m} \left(\frac{\nu(V, g)^{\frac{w}{v}}}{\nu(W, h)} \right)^{\frac{1}{m}} a^{\frac{v-2}{v}} + \frac{w}{m} \left(\frac{\nu(W, h)^{\frac{v}{w}}}{\nu(V, g)} \right)^{\frac{1}{m}} b \right] \\
&= ma_m \nu(V, g)^{\frac{1}{m}} \nu(W, h)^{\frac{1}{m}} \frac{v}{m} \left(\frac{\nu(V, g)^{\frac{w}{v}}}{\nu(W, h)} \right)^{\frac{1}{m}} a^{\frac{v-2}{v}} \\
&\quad + ma_m \nu(V, g)^{\frac{1}{m}} \nu(W, h)^{\frac{1}{m}} \frac{w}{m} \left(\frac{\nu(W, h)^{\frac{v}{w}}}{\nu(V, g)} \right)^{\frac{1}{m}} b \\
&= a_m v \nu(V, g)^{\frac{1}{v}} a^{\frac{v-2}{v}} + a_m w \nu(W, h)^{\frac{1}{w}} b \\
&= \frac{a_m}{a_v} \mu(V, g) a^{\frac{v-2}{v}} + \frac{a_m}{a_w} \mu(W, h) b
\end{aligned}$$

where we in the second line used Young's inequality

$$cd \leq \frac{v}{m} c^{\frac{m}{v}} + \frac{w}{m} d^{\frac{m}{w}},$$

which is valid for any $c, d \geq 0$. Using the above in (9) with

$$a := \int_V \left(\int_W |u|^2 dv^h \right)^{\frac{pv}{2}} dv^g \quad \text{and} \quad b := \int_V \left(\int_W |u|^{p_w} dv^h \right)^{\frac{2}{p_w}} dv^g,$$

we get

$$\mathcal{F}^G(u) \geq r \left(\int_V \left(\int_W |u|^2 dv^h \right)^{\frac{pv}{2}} dv^g \right)^{\frac{v-2}{m}} \left(\int_V \left(\int_W |u|^{p_w} dv^h \right)^{\frac{2}{p_w}} dv^g \right)^{\frac{w}{m}}.$$

Using Lemma 2.1 and Relation (5) we deduce

$$\mathcal{F}^G(u) \geq r = ma_m \nu(V, g)^{\frac{1}{m}} \nu(W, h)^{\frac{1}{m}} = \frac{ma_m}{(va_v)^{\frac{v}{m}} (wa_w)^{\frac{w}{m}}} \mu(V, g)^{\frac{v}{m}} \mu(W, h)^{\frac{w}{m}}.$$

Since this holds for all u , Theorem 2.3 follows. \square

3. APPLICATIONS

3.1. The smooth Yamabe invariant of product manifolds. Let M be a compact manifold of dimension $m \geq 3$. Then its *smooth Yamabe invariant* is defined as

$$\sigma(M) := \sup \mu(M, G)$$

where the supremum runs over all Riemannian metrics G on M . This invariant of differentiable manifolds has the property that $\sigma(M) \leq \sigma(S)$ for all M and $\sigma(M) > 0$ if and only if M admits a metric with positive scalar curvature.

From Theorem 2.3 we obtain the following corollary.

Corollary 3.1. *Let V, W be compact manifolds of dimensions $v, w \geq 3$. Assume $\sigma(V) \geq 0$. Then*

$$\sigma(V \times W) \geq \frac{ma_m}{(va_v)^{\frac{v}{m}} (wa_w)^{\frac{w}{m}}} \sigma(V)^{\frac{v}{m}} \sigma(W)^{\frac{w}{m}},$$

where $m = v + w$.

Proof. We first consider the case $\sigma(V) > 0$. In [1, Theorem 1.1] it is proven that

$$\lim_{t \rightarrow \infty} \mu(V \times W, g + t^2 h) = \mu(V \times \mathbb{R}^w, g + \xi^w)$$

if g is a metric on V with positive scalar curvature and h is any metric on W . Since $a_v \geq a_m$ we see that (4) holds, so Theorem 2.3 together with $\mu(\mathbb{R}^w, \xi^w) = \mu(S^w, \rho^w)$ imply the corollary if $\sigma(V) > 0$.

In the case $\sigma(V) = 0$ there is a sequence of metrics g_i on V such that $\text{vol}^{g_i}(V) = 1$, $\mu(V, g_i) \leq 0$, and $\mu(V, g_i) \rightarrow 0$ as $i \rightarrow \infty$. From the solution of the Yamabe problem we can assume that all g_i have constant scalar curvature $s^{g_i} = \mu(V, g_i)$. Choose $\varepsilon_i > 0$ such that $\varepsilon_i \rightarrow 0$ and $\varepsilon_i^{-w} \mu(V, g_i) \rightarrow 0$ for $i \rightarrow \infty$. For a metric h on W with constant scalar curvature s^h , the metric $G_i := \varepsilon_i^w g_i + \varepsilon_i^{-v} h$ has $\text{vol}^{G_i}(V \times W) = 1$ and constant scalar curvature $\varepsilon_i^{-w} \mu(V, g_i) + \varepsilon_i^v s^h \rightarrow 0$. It follows that $\mu(V \times W, G_i) \rightarrow 0$ and thus $\sigma(V \times W) \geq 0$. \square

3.2. Surgery formulas. Assume that M is a compact m -dimensional manifold, and that $i : S^k \times \overline{B^{m-k}} \rightarrow M$ is an embedding. We define

$$N := (M \setminus i(S^k \times B^{m-k})) \cup_{\partial} (B^{k+1} \times S^{m-k-1})$$

where \cup_{∂} means that we identify $x \in S^k \times S^{m-k-1} = \partial(B^{k+1} \times S^{m-k-1})$ with $i(x) \in \partial i(S^k \times B^{m-k})$. After a smoothing procedure N is again a compact manifold without boundary, and we say that N is *obtained from M by m -dimensional surgery along i* .

In [2, Corollary 1.4] we found the following result.

Theorem 3.2. *Let N be obtained from M via surgery of dimension $k \in \{0, 1, \dots, m-3\}$, then there is a constant $\Lambda_{m,k} > 0$ with*

$$\sigma(N) \geq \min\{\sigma(M), \Lambda_{m,k}\}.$$

Furthermore, for $k = 0$ this statement is true for $\Lambda_{m,0} = \infty$.

It is helpful to consider how the constant $\Lambda_{m,k}$ was obtained in [2] in the case $k \geq 1$. We showed that Theorem 3.2 holds for a constant $\Lambda_{m,k}$ satisfying

$$\Lambda_{m,k} \geq \min\left\{\Lambda_{m,k}^{(1)}, \Lambda_{m,k}^{(2)}\right\}.$$

We will not recall the definition $\Lambda_{m,k}^{(1)}$ and $\Lambda_{m,k}^{(2)}$ here in detail, as it is not needed, but we will explain some relevant facts for $\Lambda_{m,k}^{(1)}$ and $\Lambda_{m,k}^{(2)}$.

For $c \in [0, 1]$ let \mathbb{H}_c^{k+1} be the simply connected $(k+1)$ -dimensional complete Riemannian manifold of constant sectional curvature $-c^2$, for $c = 0$ it is \mathbb{R}^{k+1} and for $c > 0$ it is hyperbolic space rescaled by a factor c^{-2} . One defines

$$\Lambda_{m,k}^{(0)} := \inf_{c \in [0,1]} \mu(\mathbb{H}_c^{k+1} \times \mathbb{S}^{m-k-1}).$$

It was shown in [2, Corollary 1.4] that $\Lambda_{m,k}^{(1)} \geq \Lambda_{m,k}^{(0)}$ for $k \in \{1, \dots, m-3\}$. Furthermore $\Lambda_{m,k}^{(2)} \geq \Lambda_{m,k}^{(1)}$ will be shown in our publication [3] provided that $k+3 \leq m \leq 5$ or $k+4 \leq m$. Thus Theorem 3.2 holds for $\Lambda_{m,k} := \Lambda_{m,k}^{(0)}$ if $k+3 \leq m \leq 5$ or $k+4 \leq m$.

Thus in many cases we have obtained, using Corollary 3.1, an explicit number $\Lambda_{m,k}$ for which Theorem 3.2 holds.

Corollary 3.3. *If $2 \leq k \leq m-4$, then Theorem 3.2 holds for*

$$\Lambda_{m,k} = \frac{m a_m}{((k+1)a_{k+1})^{\frac{k+1}{m}} ((m-k-1)a_{m-k-1})^{\frac{m-k-1}{m}}} \sigma(S^{k+1})^{\frac{k+1}{m}} \sigma(S^{m-k-1})^{\frac{m-k-1}{m}}.$$

It follows for example: If M is an m -dimensional compact manifold, obtained from S^m by performing successive surgeries of dimension k , $0 \leq k \leq m-4$, $k \neq 1$, then $\sigma(M) \geq \Lambda_m$, where $\Lambda_6 = 54.779$, $\Lambda_7 = 74.504$, $\Lambda_8 = 92.242$, $\Lambda_9 = 109.426$, etc.

3.3. A stable Yamabe invariant. In this section we will define and discuss a “stabilized” Yamabe invariant, obtained by letting the dimension go to infinity for a given compact Riemannian manifold by multiplying with Ricci-flat manifolds of increasing dimension. Very optimistically, such a stabilization could be related to the linear eigenvalue problem obtained by formally letting the dimension tend to infinity in the Yamabe problem. The stable invariant can also be viewed as a quantitative refinement of the property that a given manifold admit stably positive scalar curvature.

For a compact manifold M with $\sigma(M) \geq 0$ we define

$$\Sigma(M) := \left(\frac{\sigma(M)}{ma_m} \right)^m,$$

then

$$\Sigma(M) = \sup \nu(M, G)$$

where the supremum runs over all Riemannian metrics G on M . The conclusion of Corollary 3.1 can be formulated as

$$\Sigma(V \times W) \geq \Sigma(V)\Sigma(S^w). \quad (10)$$

Let (B, β) be a compact Ricci-flat manifold of dimension b . We could for example choose B to be the 1-dimensional circle S^1 , or an 8-dimensional Bott manifold equipped with a metric with holonomy $\text{Spin}(7)$. From (10) we then get

$$\frac{\Sigma(S^{v+bi})}{\Sigma(S^{bi})} \geq \frac{\Sigma(V \times B^i)}{\Sigma(S^{bi})} \geq \Sigma(V), \quad (11)$$

where the upper bound comes from $\Sigma(V \times B^i) \leq \Sigma(S^{v+bi})$. We define the stable Yamabe invariant of V as the limit superior of the middle term,

$$\bar{\Sigma}(V) := \limsup_{i \rightarrow \infty} \frac{\Sigma(V \times B^i)}{\Sigma(S^{bi})}$$

To see that the stable Yamabe invariant is finite we need to study the upper bound in (11), and the function $v \mapsto \Sigma(S^v)$. We have

$$\sigma(S^v) = v(v-1)\omega_v^{2/v}, \quad \omega_v = \frac{2\pi^{\frac{v+1}{2}}}{\Gamma\left(\frac{v+1}{2}\right)},$$

where ω_v is the volume of \mathbb{S}^v , so

$$\Sigma(S^v) = 4\pi \left(\frac{\pi(v-2)}{4} \right)^v \frac{1}{\Gamma\left(\frac{v+1}{2}\right)^2}.$$

Stirling’s formula tells us that

$$\Gamma(z) = \sqrt{\frac{2\pi}{z}} \left(\frac{z}{e} \right)^z \left(1 + O\left(\frac{1}{z}\right) \right)$$

and therefore

$$\begin{aligned}\Sigma(S^v) &= 4\pi \left(\frac{\pi(v-2)}{4}\right)^v \frac{v+1}{4\pi} \left(\frac{2e}{v+1}\right)^{v+1} \left(1 + O\left(\frac{1}{v}\right)\right) \\ &= 2e \left(\frac{\pi e}{2}\right)^v \frac{(1-2/v)^v}{(1+1/v)^v} \left(1 + O\left(\frac{1}{v}\right)\right) \\ &= 2e^{-2} \left(\frac{\pi e}{2}\right)^v \left(1 + O\left(\frac{1}{v}\right)\right).\end{aligned}$$

We see that

$$\lim_{i \rightarrow \infty} \frac{\Sigma(S^{v+bi})}{\Sigma(S^{bi})} = \lim_{i \rightarrow \infty} \left(\frac{\pi e}{2}\right)^v \left(1 + O\left(\frac{1}{bi}\right)\right) = \left(\frac{\pi e}{2}\right)^v,$$

so from (11) we get the following bound on the stable Yamabe invariant

$$\left(\frac{\pi e}{2}\right)^v \geq \bar{\Sigma}(V) \geq \Sigma(V).$$

We conclude that the stable invariant is a non-trivial invariant.

The stable Yamabe invariant is not strictly speaking a stable invariant in the sense that it gives the same value for V and $V \times B^i$. These values are however related by a simple identity, as we will see next. Taking the limit superior as $j \rightarrow \infty$ in

$$\frac{\Sigma(V \times B^i \times B^j)}{\Sigma(S^{bj})} = \frac{\Sigma(V \times B^{i+j})}{\Sigma(S^{bi+bj})} \frac{\Sigma(S^{bi+bj})}{\Sigma(S^{bj})}$$

we conclude

$$\bar{\Sigma}(V \times B^i) = \bar{\Sigma}(V) \left(\frac{\pi e}{2}\right)^{bi}$$

and further

$$\bar{\Sigma}(V) \geq \Sigma(V \times B^i) \left(\frac{\pi e}{2}\right)^{-bi} \tag{12}$$

for all $i \geq 0$.

The next simple proposition tells us that positivity of $\Sigma(V)$ is equivalent to V having stably metrics of positive scalar curvature.

Proposition 3.4. *Let V be a compact manifold. The following three statements are equivalent.*

- (a) $\bar{\Sigma}(V) > 0$.
- (b) There is $i_0 > 0$ such that $V \times B^{i_0}$ admits a positive scalar curvature metric.
- (c) There is a $i_0 > 0$ such that $V \times B^i$ admits a positive scalar curvature metric for all $i \geq i_0$.

Proof. The implications (a) \Rightarrow (b) and (b) \Leftrightarrow (c) are easy to show. The implication (b) \Rightarrow (a) is a consequence of (12). \square

We also obtain a stable version of Theorem 3.2 for surgeries of codimension at least 4. A similar result holds for surgeries of codimension 3, but with a less explicit constant.

Theorem 3.5. *Assume that N is obtained from the compact m -dimensional manifold M by surgery of dimension k , where $0 \leq k \leq m - 4$, then*

$$\bar{\Sigma}(N) \geq \min \left\{ \bar{\Sigma}(M), \bar{\Sigma}(S^m), \left(\frac{\pi e}{2}\right)^{k+1} \Sigma(S^{m-k-1}) \right\}.$$

Proof. The manifold N after surgery is obtained by a connected sum of M and S^m along embeddings of a k -dimensional sphere with trivial normal bundle. Similarly $N \times B^i$ is obtained by a connected sum of $M \times B^i$ and $S^m \times B^i$ by a connected sum along embeddings of $S^k \times B^i$ with trivial normal bundle. Thus [2, Theorem 1.3] together with Corollary 3.3 tells us that

$$\begin{aligned} \Sigma(N \times B^i) &\geq \min \left\{ \Sigma(M \times B^i), \Sigma(S^m \times B^i), \left(\frac{\Lambda_{m+bi, k+bi}}{(m+bi)a_{m+bi}} \right)^{m+bi} \right\} \\ &\geq \min \{ \Sigma(M \times B^i), \Sigma(S^m \times B^i), \Sigma(S^{k+bi+1})\Sigma(S^{m-k-1}) \} \end{aligned}$$

and this yields the statement of the theorem. \square

For the smooth Yamabe invariant the value of the sphere is a universal upper bound. One can ask if the same holds for the stable invariant, is $\bar{\Sigma}(M) \leq \bar{\Sigma}(S^m)$ for all M ?

REFERENCES

- [1] K. Akutagawa, L. Florit, and J. Petean, *On the Yamabe constant of riemannian products*, Comm. Anal. Geom. **15** (2007), 947–969.
- [2] B. Ammann, M. Dahl, and E. Humbert, *Smooth Yamabe invariant and surgery*, Preprint, ArXiv 0804.1418, 2008.
- [3] ———, *Square-integrability of solutions of the Yamabe equation*, Preprint in preparation, 2011.
- [4] T. Aubin, *Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire.*, J. Math. Pur. Appl., IX. Ser. **55** (1976), 269–296.
- [5] A. Benedek and R. Panzone, *The space L^p , with mixed norm*, Duke Math. J. **28** (1961), 301–324.
- [6] L. Bérard Bergery and G. Kaas, *Examples of multiple solutions for the Yamabe problem on scalar curvature*, Preprint <http://hal.archives-ouvertes.fr/hal-00143495/>.
- [7] ———, *Remark on an example by R. Schoen concerning the scalar curvature*, Preprint <http://hal.archives-ouvertes.fr/hal-00143485/>.
- [8] C. Böhm, M. Wang, and W. Ziller, *A variational approach for compact homogeneous Einstein manifolds*, Geom. Funct. Anal. **14** (2004), 681–733.
- [9] N. Große, *The Yamabe equation on manifolds of bounded geometry*, Preprint, arxiv 0912.4398, 2009.
- [10] G. Henry and J. Petean, *Isoperimetric hypersurfaces and metrics of constant scalar curvature*, Tech. report, Preprint CIMAT Mexico, 2011.
- [11] S. Kim, *An obstruction to the conformal compactification of Riemannian manifolds*, Proc. Amer. Math. Soc. **128** (2000), no. 6, 1833–1838.
- [12] J. M. Lee and T. H. Parker, *The Yamabe problem*, Bull. Am. Math. Soc., New Ser. **17** (1987), 37–91.
- [13] M. Obata, *The conjectures on conformal transformations of Riemannian manifolds*, J. Diff. Geom. **6** (1971/72), 247–258.
- [14] J. Petean, *Best Sobolev constants and manifolds with positive scalar curvature metrics*, Ann. Global Anal. Geom. **20** (2001), 231–242.
- [15] ———, *Isoperimetric regions in spherical cones and Yamabe constants of $M \times S^1$* , Geom. Dedicata **143** (2009), 37–48.
- [16] ———, *Metrics of constant scalar curvature conformal to Riemannian products*, Proc. Amer. Math. Soc. **138** (2010), 2897–2905.
- [17] J. Petean and J. M. Ruiz, *Isoperimetric profile comparisons and Yamabe constants*, preprint, 2010, ArXiv 1010.3642.
- [18] R. Schoen, *Conformal deformation of a Riemannian metric to constant scalar curvature*, J. Diff. Geom. **20** (1984), 479–495.
- [19] ———, *On the number of constant scalar curvature metrics in a conformal class*, Differential geometry, Pitman Monogr. Surveys Pure Appl. Math., vol. 52, Longman Sci. Tech., Harlow, 1991, pp. 311–320.
- [20] N. S. Trudinger, *Remarks concerning the conformal deformation of Riemannian structures on compact manifolds*, Ann. Sc. Norm. Super. Pisa, Sci. Fis. Mat., III. Ser. **22** (1968), 265–274.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY
E-mail address: `bernd.ammann@mathematik.uni-regensburg.de`

INSTITUTIONEN FÖR MATEMATIK, KUNGLIGA TEKNISKA HÖGSKOLAN, 100 44 STOCKHOLM, SWEDEN
E-mail address: `dahl@math.kth.se`

INSTITUT ÉLIE CARTAN, BP 239, UNIVERSITÉ DE NANCY 1, 54506 VANDOEUVRE-LÈS-NANCY
CEDEX, FRANCE
E-mail address: `humbert@iecn.u-nancy.fr`