

# Universität Regensburg Mathematik



## Higher congruence companion forms

Rajender Adibhatla and Jayanta Manoharmayum

Preprint Nr. 26/2011

# HIGHER CONGRUENCE COMPANION FORMS

RAJENDER ADIBHATLA AND JAYANTA MANOHARMAYUM

ABSTRACT. For a rational prime  $p \geq 3$  we consider  $p$ -ordinary, Hilbert modular newforms  $f$  of weight  $k \geq 2$  with associated  $p$ -adic Galois representations  $\rho_f$  and  $\bmod p^n$  reductions  $\rho_{f,n}$ . Under suitable hypotheses on the size of the image, we use deformation theory and modularity lifting to show that if the restrictions of  $\rho_{f,n}$  to decomposition groups above  $p$  split then  $f$  has a companion form  $g$  modulo  $p^n$  (in the sense that  $\rho_{f,n} \sim \rho_{g,n} \otimes \chi^{k-1}$ ).

## 1. INTRODUCTION

Let  $F$  be a totally real number field and let  $p$  be an odd prime. Suppose we are given a Hilbert modular newform  $f$  over  $F$  of level  $\mathfrak{n}_f$ , character  $\psi$  and (parallel) weight  $k \geq 2$ . For a prime  $\mathfrak{q}$  not dividing  $\mathfrak{n}_f$ , let  $c(\mathfrak{q}, f)$  denote the eigenvalue of the Hecke operator  $T(\mathfrak{q})$  acting on  $f$ ; denote by  $K_f$  the number field generated by the  $c(\mathfrak{q}, f)$ 's and  $\psi(\text{Frob}_{\mathfrak{q}})$ 's, and by  $\mathcal{O}_f$  the integer ring of  $K_f$ . Then for each prime  $\wp|p$  of  $\mathcal{O}_f$  one has a continuous, odd, absolutely irreducible representation  $\rho_{f,\wp} : G_F \rightarrow GL_2(\mathcal{O}_{f,\wp})$  characterized by the following property:  $\rho_{f,\wp}$  is unramified outside primes dividing  $p\mathfrak{n}_f$ , and at a prime  $\mathfrak{q} \nmid p\mathfrak{n}_f$  the characteristic polynomial of  $\rho_{f,\wp}(\text{Frob}_{\mathfrak{q}})$  is  $X^2 - c(\mathfrak{q}, f)X + \psi(\text{Frob}_{\mathfrak{q}})\text{Nm}(\mathfrak{q})^{k-1}$ . We denote the  $p$ -adic cyclotomic character by  $\chi$ . Thus the determinant of  $\rho_{f,\wp}$  is  $\psi\chi^{k-1}$ .

From here on we assume that the character  $\psi$  is unramified at  $p$ . Suppose that  $f$  is ordinary at  $p$ . Then, by Wiles [18], and Mazur-Wiles [14], for every prime  $\mathfrak{p}|p$  we have

$$\rho_{f,\wp}|_{G_{\mathfrak{p}}} \sim \begin{pmatrix} \psi_{1\mathfrak{p}}\chi^{k-1} & * \\ 0 & \psi_{2\mathfrak{p}} \end{pmatrix}$$

where  $\psi_{1\mathfrak{p}}, \psi_{2\mathfrak{p}}$  are unramified characters. In fact, with  $a(\mathfrak{p}, f)$  defined to be the unit root  $X^2 - c(\mathfrak{p}, f)X + \psi(\text{Frob}_{\mathfrak{p}})\text{Nm}(\mathfrak{p})^{k-1} = 0$ , we have  $\psi_{2\mathfrak{p}}(\text{Frob}_{\mathfrak{p}}) = c(\mathfrak{p}, f)$ . A natural question is to ask when the restriction(s)  $\rho_{f,\wp}|_{G_{\mathfrak{p}}}$  actually split. We note that if  $\rho_{f,\wp} \bmod \wp$  is absolutely irreducible then the splitting (or not) of  $\rho_{f,\wp} \bmod \wp^n$  is independent of the choice of a lattice used to define  $\rho_{f,\wp}$ . Indeed, if for some  $M \in GL_2(K_{f,\wp})$  the conjugate  $M\rho_{f,\wp}M^{-1}$  is integral and stabilises the upper triangular decomposition at  $\mathfrak{p}$ , then  $M$  is a scalar multiple of  $\begin{pmatrix} u & v \\ 0 & 1 \end{pmatrix}$  where  $u \equiv 1 \pmod{\wp}$  and  $v \equiv 0 \pmod{\wp}$ . If we denote by  $c_{\mathfrak{p}} \in H^1(G_{\mathfrak{p}}, \mathcal{O}_{f,\wp}(\psi_{1\mathfrak{p}}\psi_{2\mathfrak{p}}^{-1}\chi^{k-1}))$  the cohomology class for  $\rho_{f,\wp}|_{G_{\mathfrak{p}}}$ , then the cohomology class for the extension at  $\mathfrak{p}$  determined by  $M\rho_{f,\wp}M^{-1}$  is  $uc_{\mathfrak{p}}$ . Hence, if  $\rho_{f,\wp} \bmod \wp$  is absolutely irreducible we can speak of  $\rho_{f,\wp} \bmod \wp^n$  being split without any ambiguity.

Now suppose we are given a second newform  $g$  which is also ordinary at  $p$ . Fix a  $p$ -adic integer ring  $\mathcal{O}$  in which  $\mathcal{O}_f$  and  $\mathcal{O}_g$  embed, and let  $\pi$  be a uniformiser. We say that  $g$  is a  $\bmod \pi^n$

---

Date: Aug 2011.

2010 *Mathematics Subject Classification.* 11F33, 11F80.

*weak companion form* for  $f$  if  $c(\mathfrak{q}, f) \equiv c(\mathfrak{q}, g) \text{Nm}(\mathfrak{q})^{k-1} \pmod{\pi^n}$  for all but finitely many primes  $\mathfrak{q}$ . The weight  $k'$  of  $g$  is then determined by  $k$  via the congruence  $\chi^{k'-1} \equiv \chi^{1-k} \pmod{\pi^n}$  on each decomposition group above  $p$ . Note that we do not enforce any optimality requirement on the level of  $g$  (and hence the prefix ‘weak’).

Classically, companion forms  $\pmod{p}$  played an important part in the weight optimisation part of Serre’s Modularity Conjecture. Serre’s predicted equivalence between local splitting for the residual modular representation (tame ramification) and the existence of companions was established by Gross in [8]. In much the same spirit, the main result of this paper, which we now state, proves the equivalence between splitting  $\pmod{p^n}$  and the existence of  $\pmod{p^n}$  weak companion forms.

**Main Theorem.** *Let  $F$  be a totally real number field,  $p$  be an odd prime unramified in  $F$ , and let  $f$  be a  $p$ -ordinary Hilbert modular newform  $f$  of squarefree level  $\mathfrak{n}$ , character  $\psi$  with order coprime to  $p$  and unramified at  $p$ , and weight  $k \geq 2$ . Let  $n \geq 2$  and set  $\rho_{f,n} := \rho_{f,\wp} \pmod{p^n}$ ,  $\bar{\rho}_f := \rho_{f,\wp} \pmod{\wp}$ ,  $\mathbf{k} := \mathcal{O}_f/\wp$ . Assume the following hypotheses:*

- *Global conditions.*

(GC1)  $\rho_{f,n}$  takes values in  $GL_2(W/p^n)$  where  $W$  is the Witt ring of  $\mathbf{k} := \mathcal{O}_f/\wp$  (under the natural injection  $W/p^n \hookrightarrow \mathcal{O}_f/\wp^n$ ).

(GC2) The image of  $\rho_{f,\wp} \pmod{\wp}$  contains  $SL_2(\mathbf{k})$ . Furthermore, if  $p = 3$  then the image  $\rho_{f,n}$  contains a transvection  $\begin{pmatrix} 1 & \\ 0 & 1 \end{pmatrix}$ .

- *Local conditions.*

(LC1) If  $\mathfrak{p}$  is a prime dividing  $p$  then  $c(\mathfrak{p}, f)^2 \not\equiv \psi(\text{Frob}_{\mathfrak{p}}) \pmod{\wp}$ .

(LC2) Let  $\mathfrak{q}$  be a prime dividing the level  $\mathfrak{n}$  where  $\bar{\rho}_f$  is unramified. If  $\text{Nm}(\mathfrak{q}) \equiv 1 \pmod{p}$  then  $p$  divides the order of  $\bar{\rho}_f(\text{Frob}_{\mathfrak{q}})$ .

Let  $k' \geq 2$  be the smallest integer such that  $k + k' \equiv 2 \pmod{(p-1)p^{n-1}}$ . Then  $f$  splits  $\pmod{p^n}$  if and only if it has a  $p$ -ordinary  $\pmod{p^n}$  weak companion form  $g$  of weight  $k'$  and character  $\psi$ .

The proof, given in section 3.2, relies on being able to lift  $\rho_{f,n} \otimes \chi^{1-k}$  to characteristic 0 with prescribed local properties and then proving that the lift is modular by using results of Skinner and Wiles in [17] along with the existence of companion forms  $\pmod{p}$  over totally real fields due to Gee ([5, Theorem 2.1]). The construction of characteristic 0 lifts for certain classes of  $\pmod{p^n}$  representations is carried out in section 3.1. (See Theorem 3.2 for the statement.)

The Main Theorem, in practice, is not useful for checking when a given newform fails to split  $\pmod{p^n}$  because we have very little control over the level of the weak companion form. However, as we show in Section 4, in the situation when the dimension of the tangent space  $\mathfrak{t}_{\mathcal{D}}$  (associated to a deformation condition  $\mathcal{D}$  of  $\bar{\rho}$ ) is 0, we can use higher companion forms to computationally verify a conjecture of Greenberg connecting local splitting with complex multiplication. We conclude by giving examples in support of this conjecture.

## 2. TOOLKIT

The method we use for obtaining a fine structure on deformations of a  $\pmod{p^n}$  representation is an adaptation of the more familiar  $\pmod{p}$  case and has two key components: the existence of sufficiently well behaved local deformations; and, for the existence of characteristic 0 liftings, being able to place local constraints so that the dual Selmer group vanishes. Naturally, both of these present difficulties in the general  $\pmod{p^n}$  case. In this section, we discuss the tools that will enable us to manage the difficulties for certain classes of  $\pmod{p^n}$  representations.

Throughout this section  $p$  is an odd prime,  $\mathbf{k}$  is a finite field of characteristic  $p$  and  $W$  is the Witt ring of  $\mathbf{k}$ .

**2.1. Deformations and substantial deformation conditions.** In the main, we follow Mazur's treatment of deformations and deformation conditions in [13]. Given a residual representation, a deformation condition is simply a collection of liftings satisfying some additional properties (closure under projections, a Mayer-Vietoris property etc). The fundamental consequence then is the existence of a (uni)versal deformation.

We expand on this: Suppose we are given a 'nice' profinite group  $\Gamma$ , and a continuous representation  $\bar{\rho} : \Gamma \rightarrow GL_2(\mathbf{k})$ . If  $\mathcal{D}$  is a deformation condition for  $\bar{\rho}$  then there is a complete local Noetherian  $W$ -algebra  $R$  with residue field  $\mathbf{k}$  and a lifting  $\rho : \Gamma \rightarrow GL_2(R)$  in  $\mathcal{D}$  with the following property: If  $\rho' : \Gamma \rightarrow GL_2(A)$  is a lifting of  $\bar{\rho}$  in  $\mathcal{D}$  then there is a morphism  $R \rightarrow A$  which gives, on composition with  $\rho$ , a representation strictly equivalent to  $\rho'$ . In addition, we require that the morphism above is unique when  $A$  is the ring of dual numbers  $\mathbf{k}[\epsilon]/(\epsilon^2)$ . If the projective image of  $\bar{\rho}$  has trivial centralizer then  $R$ , together with  $\rho$ , represents the functor that assigns type  $\mathcal{D}$  deformations to a coefficient ring. We shall use the natural identification of the tangent space  $\mathfrak{t}_{\mathcal{D}}$  with a subspace of  $H^1(\Gamma, \text{ad}\bar{\rho})$  (and as a subspace of  $H^1(\Gamma, \text{ad}^0\bar{\rho})$  when considering deformations with a fixed determinant). The (uni)versal deformation ring  $R$  then has a presentation  $W[[T_1, \dots, T_n]]/J$  where  $n = \dim_{\mathbf{k}} \mathfrak{t}_{\mathcal{D}}$ . We will be particularly interested in smooth deformation conditions (so the ideal of relations  $J$  will be (0)).

As hinted in the beginning of this section, the method we use for constructing smooth global deformation conditions depends upon being able to find local (uni)versal deformation rings smooth in a large number of variables. It will be convenient to make the following definition:

**Definition 2.1.** Let  $F$  be a local field and let  $\bar{\rho} : G_F \rightarrow GL_2(\mathbf{k})$  be a residual representation.

- (a) We call a deformation condition for  $\bar{\rho}$  with fixed determinant *substantial* if it is smooth and its tangent space  $\mathfrak{t}$  satisfies the inequality

$$\dim_{\mathbf{k}} \mathfrak{t} \geq \dim_{\mathbf{k}} H^0(G_F, \text{ad}^0\bar{\rho}) + [F : \mathbb{Q}_p]\delta$$

where  $\delta$  is 1 when  $F$  has residue characteristic  $p$  and 0 otherwise.

- (b) A deformation  $\rho : G_F \rightarrow GL_2(A)$  of  $\bar{\rho}$  is substantial if it is part of a substantial deformation condition.

We now give examples of substantial deformation conditions. From here on, for the rest of the section,  $F$  is a finite extension of  $\mathbb{Q}_l$  for some prime  $l$ . As in the definition above, let  $\bar{\rho} : G_F \rightarrow GL_2(\mathbf{k})$  be a residual representation.

**Example 2.2.** Assume that the residue characteristic of  $F$  is different from  $p$ . Suppose that the order of  $\bar{\rho}(I_F)$  is co-prime to  $p$ , and let  $d : G_F \rightarrow W^\times$  be a character lifting  $\det\bar{\rho}$ . The collection of liftings of  $\bar{\rho}$  which factor through  $G_F/(I_F \cap \ker\bar{\rho})$  and have determinant  $d$  is a substantial deformation condition. The tangent space has dimension  $\dim_{\mathbf{k}} H^0(G_F, \text{ad}^0\bar{\rho})$ .

**Example 2.3.** Suppose that

$$\bar{\rho} \sim \begin{pmatrix} \bar{\chi} & * \\ 0 & 1 \end{pmatrix}_{\bar{\epsilon}}$$

for some character  $\bar{\varepsilon} : G_F \rightarrow \mathbf{k}^\times$ . Moreover, assume that if  $\bar{\rho}$  is semi-simple then  $\bar{\chi}$  is non-trivial. Fix a character  $\varepsilon : G_F \rightarrow W^\times$  lifting  $\bar{\varepsilon}$ . Then the collection of liftings strictly equivalent to

$$\begin{pmatrix} \chi & * \\ 0 & 1 \end{pmatrix} \varepsilon$$

is a substantial deformation condition. Note that  $\bar{\rho}$  is equivalent to a representation of the form considered above only if  $p$  divides the order of  $\bar{\rho}(I_F)$ . (See Example 3.3 of [11].)

**Example 2.4.** We now assume that the residue characteristic of  $F$  is  $p$ . Suppose we are given an integer  $k \geq 2$  and a representation  $\bar{\rho} : G_F \rightarrow GL_2(\mathbf{k})$  such that

$$\bar{\rho} = \begin{pmatrix} \bar{\chi}^{k-1} \bar{\psi}_1 & * \\ 0 & \bar{\psi}_2 \end{pmatrix}$$

where  $\bar{\psi}_1, \bar{\psi}_2$  are unramified characters. Let  $\psi$  be the Teichmüller lift of  $\bar{\psi}_1 \bar{\psi}_2$ . If  $A$  is a coefficient ring, we shall call a lifting  $\rho_A : G_F \rightarrow GL_2(A)$  of  $\bar{\rho}$  a  $\bar{\psi}_2$ -good weight  $k$  lifting with character  $\psi$  if  $\rho_A$  is strictly equivalent to a representation of the form

$$\begin{pmatrix} \widetilde{\psi}_1 \chi^{k-1} & * \\ 0 & \widetilde{\psi}_2 \end{pmatrix}$$

for some unramified characters  $\widetilde{\psi}_1, \widetilde{\psi}_2 : G_F \rightarrow A^\times$  lifting  $\bar{\psi}_1, \bar{\psi}_2$  and  $\widetilde{\psi}_1 \widetilde{\psi}_2 = \psi$ .

We then have the following property of weight  $k$  liftings (proof immediate, Example 3.4 in [11]):

**Proposition 2.5.** *Let  $\bar{\rho} : G_F \rightarrow GL_2(\mathbf{k})$  be as above in Example 2.4, and further assume that  $\bar{\chi}^{k-1} \bar{\psi}_1 \neq \bar{\chi} \bar{\psi}_2$ . Then the deformation condition consisting of weight  $k$  liftings of  $\bar{\rho}$  is a smooth deformation condition. The dimension of its tangent space is equal to  $[F : \mathbb{Q}_p] + \dim_{\mathbf{k}} H^0(G_F, \text{ad}^0 \bar{\rho})$ .*

We conclude by determining all substantial deformation conditions for residual representations of a particular shape. For the remainder of this section, we assume  $F$  has residue field of order  $q$  with  $p \nmid q$  and let  $\bar{\rho} : G_F \rightarrow GL_2(\mathbf{k})$  be an unramified representation with  $\bar{\rho}(\text{Frob}) = \begin{pmatrix} q\alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix}$ . Of course, any lift of  $\bar{\rho}$  necessarily factors through the maximal tamely ramified extension  $F^{\text{tr}}$  of  $F$ . We fix a generator  $\tau$  of the tame inertia, a lift  $\sigma$  of Frobenius to  $\text{Gal}(F^{\text{tr}}/F)$ . One then has the following description:

$$\text{Gal}(F^{\text{tr}}/F) = \langle \sigma, \tau \mid \sigma \tau \sigma^{-1} = \tau^q \rangle.$$

First define polynomials  $h_n(T) \in \mathbb{Z}[T], n \geq 1$ , by the recursion  $h_{n+2} = T h_{n+1} - h_n$  and initial values  $h_1 := 1, h_2 := T$ . The following properties of  $h_n$  are easily verified by induction:

- $h_n(2) = n$
- If  $M$  is a 2 by 2 matrix over any commutative ring with trace  $t$  and determinant 1 then,
 
$$M^n = h_n(t)M - h_{n-1}(t)I$$
- $h_n^2 - T h_n h_{n-1} + h_{n-1}^2 = 1$

We then have the following proposition.

**Proposition 2.6.** *Let  $\bar{\rho}$  be as above with  $q\alpha \neq \alpha^{-1}$ . Denote by  $\hat{\alpha}$  the Teichmüller lift of  $\alpha$ . Let  $R$ , resp.  $\rho : \text{Gal}(F^{\text{tr}}/F) \rightarrow \text{GL}_2(R)$ , be the versal deformation ring, resp. the versal representation, for liftings of  $\bar{\rho}$  with determinant  $\chi$ .*

- (i) *Suppose  $\alpha^2 \neq 1$  and  $q^2\alpha^2 \neq 1$ . If  $q \equiv 1 \pmod{p}$  then  $R \cong W(k)[[S, T]]/\langle(1+T)^q - (1+T)\rangle$  and*

$$\rho(\sigma) = \begin{pmatrix} q\hat{\alpha}(1+S) & 0 \\ 0 & (\hat{\alpha}(1+S))^{-1} \end{pmatrix}, \quad \rho(\tau) = \begin{pmatrix} 1+T & 0 \\ 0 & (1+T)^{-1} \end{pmatrix}.$$

*If  $q \not\equiv 1 \pmod{p}$  then  $R \cong W[[S]]$  and  $\rho(\sigma) = \begin{pmatrix} q\hat{\alpha}(1+S) & 0 \\ 0 & (\hat{\alpha}(1+S))^{-1} \end{pmatrix}$ . In any case, a deformation condition is substantial if and only if it is unramified.*

- (ii) *If  $\alpha^2 = 1$  and  $q^2 \not\equiv 1 \pmod{p}$  then  $R \cong W(k)[[S, T]]/(ST)$  and*

$$\rho(\sigma) = \hat{\alpha} \begin{pmatrix} q(1+S) & 0 \\ 0 & (1+S)^{-1} \end{pmatrix}, \quad \rho(\tau) = \begin{pmatrix} 1 & T \\ 0 & 1 \end{pmatrix}.$$

*A deformation condition for  $\bar{\rho}$  with determinant  $\chi$  is substantial if and only if it is either unramified or of the type considered in Example 2.3.*

- (iii) *If  $\alpha^2 = 1$  and  $q \equiv -1 \pmod{p}$ , then  $R := W(k)[[S, T_1, T_2]]/J$  where*

$$J := \langle T_1(q(1+S)^2 - h_q(2\sqrt{1+T_1T_2})), T_2(1 - q(1+S)^2 h_q(2\sqrt{1+T_1T_2})) \rangle, \quad \text{and}$$

$$\rho(\sigma) = \hat{\alpha} \begin{pmatrix} q(1+S) & 0 \\ 0 & (1+S)^{-1} \end{pmatrix}, \quad \rho(\tau) = \begin{pmatrix} \sqrt{1+T_1T_2} & T_1 \\ T_2 & \sqrt{1+T_1T_2} \end{pmatrix}.$$

*The only ramified substantial deformation condition for  $\bar{\rho}$  is the of the type given Example 2.3: it corresponds to the quotient  $W(k)[[S, T_1, T_2]]/(S, T_2)$ .*

*Proof.* Let  $A$  be a coefficient ring with maximal ideal  $\mathfrak{m}_A$ , and let be  $\rho_A$  a lifting of  $\bar{\rho}$  with determinant  $\chi$ . By Hensel's Lemma, we can assume that  $\rho_A(\sigma)$  is diagonal. Let

$$(2.1) \quad \rho_A(\sigma) = \begin{pmatrix} q\hat{\alpha}(1+s) & 0 \\ 0 & (\hat{\alpha}(1+s))^{-1} \end{pmatrix}, \quad \rho_A(\tau) = \begin{pmatrix} a & t_1 \\ t_2 & d \end{pmatrix}$$

with  $s, t_1, t_2, a-1, d-1 \in \mathfrak{m}_A$  and  $ad - t_1t_2 = 1$ . Since  $\sigma\tau\sigma^{-1} = \tau^q$ , we have

$$(2.2) \quad \begin{pmatrix} a & t_1q(\hat{\alpha}(1+s))^2 \\ t_2q^{-1}(\hat{\alpha}(1+s))^{-2} & d \end{pmatrix} = \begin{pmatrix} ah_q(t) - h_{q-1}(t) & t_1h_q(t) \\ t_2h_q(t) & dh_q(t) - h_{q-1}(t) \end{pmatrix}$$

where  $t = a + d$  is the trace. Note that  $t \equiv 2 \pmod{\mathfrak{m}_A}$  and so  $h_q(t) \equiv q \pmod{\mathfrak{m}_A}$ .

If  $\alpha^2 \neq 1$  then  $q(\hat{\alpha}(1+s))^2 - h_q(t)$  is a unit and we get  $t_1 = 0$ . Similarly, if  $q^2\alpha^2 \neq 1$  then  $t_2 = 0$ . The claims made in part (i) of the proposition are now immediate.

We now continue our analysis of  $\rho_A$  under the assumption that  $\alpha^2 = 1$  and  $q \not\equiv \pm 1 \pmod{p}$ . For ease of notation, we shall in fact assume that  $\hat{\alpha} = 1$ . Since  $1 - h_q(t)$  is a unit, taking the difference of the diagonal entries on both sides of (2.2) gives  $a - d = 0$  and so  $a = d = \sqrt{1 + t_1 t_2}$ . Comparison of the off-diagonal entries of (2.2) (followed by multiplication) produces  $t_1 t_2 (1 - h_q(t)^2) = 0$ .

Suppose now that  $q \not\equiv -1 \pmod{p}$ . Then  $t_1 t_2 = 0$  and so  $t = 2, h_q(t) = q$ . We can now simplify the two relations from the off-diagonal entries to get  $t_1 = t_1(1 + s)^2$  and  $t_2 = t_2 q^2(1 + s)^2$ , and finally deduce that  $st_1 = 0, t_2 = 0$ . Part (ii) of the proposition now follows easily.

Finally, we consider the case  $q \equiv -1 \pmod{p}$ . The presentation for  $R$  and  $\rho$  follows from the presentation of an arbitrary lift along with the fact that  $\dim H^1(G_F, \text{ad}^0 \bar{\rho}) = 3$ . We now indicate how to determine the substantial deformation conditions. Take  $A$  to be characteristic 0 (and  $\hat{\alpha} = 1$ ). In the presentation (2.1) the trace of  $\rho_A(\tau)$  is  $2\sqrt{1 + t_1 t_2}$ . If  $t_1 t_2 \neq 0$  then  $\rho_A(\tau)$  has distinct eigenvalues - contradicting the fact that  $\rho_A$  is twist equivalent to  $\begin{pmatrix} \chi & * \\ 0 & 1 \end{pmatrix}$  over its field of fractions. Hence we must have  $t_2 = 0$  and  $s = 0$ .  $\square$

**2.2. Subgroups of  $GL_2(W/p^n)$ .** We now derive some properties of certain subgroups of  $GL_2(W/p^n)$  which will be of relevance in constructing global deformations. Let's recall that  $p$  is an odd prime, and that  $\mathbf{k}$  is the residue field  $W/p$ . We denote by  $\text{ad}^0$  the the vector space of  $2 \times 2$ -matrices over  $\mathbf{k}$  with  $GL_2(W/p^n)$  acting by conjugation, and by  $\text{ad}^0(i)$  its twist by the  $i$ -th power of the determinant. For convenience, we record the following useful identity

$$(2.3) \quad \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \begin{pmatrix} 1 & -x \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} a + cx & b - 2ax - cx^2 \\ c & -a - cx \end{pmatrix}.$$

It is well known—see Lemma 2.48 of [3], for instance—that  $H^1(SL_2(\mathbf{k}), \text{ad}^0) = 0$  except when  $\mathbf{k} = \mathbb{F}_5$ . We now state and proof (for completeness) the following result in the exceptional case.

**Lemma 2.7.**  $H^1(GL_2(\mathbb{F}_5), \text{ad}^0(i)) = 0$  if  $i = 0, 1$  or  $3$ .

*Proof.* Let  $B \supset U$  be the subgroups of  $GL_2(\mathbb{F}_5)$  consisting of matrices of the form  $\begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$  respectively. We then need to verify that  $H^1(B, \text{ad}^0(i)) \cong H^1(U, \text{ad}^0(i))^{B/U} = (0)$ .

Let  $\sigma := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \tau := \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}$ . From (2.3) it follows that  $(\sigma - 1)\text{ad}^0$  is the subspace of upper triangular matrices in  $\text{ad}^0$ . Thus if  $0 \neq \xi \in H^1(U, \text{ad}^0(i))$  then we can assume that  $\xi(\sigma) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ , and  $\xi$  is fixed by  $B/U$  if and only if  $(\tau * \xi)(\sigma) - \xi(\sigma)$  is upper triangular. Now  $(\tau * \xi)(\sigma) = \tau \xi(\sigma^2) \tau^{-1} = \begin{pmatrix} 1 & 2 \\ -1 & 2 \end{pmatrix}$  and so  $\xi \in H^1(U, \text{ad}^0(i))^{B/U}$  if and only if  $1 + 3^i = 0$ .  $\square$

**Proposition 2.8.** *Let  $G$  be a subgroup of  $GL_2(W/p^n)$ . Suppose the mod  $p$  reduction of  $G$  contains  $SL_2(\mathbf{k})$ . Furthermore, assume that if  $p = 3$  then  $G$  contains a transvection  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . Then the following statements hold.*

- (a)  $G$  contains  $SL_2(W/p^n)$ .
- (b) Suppose that  $p \geq 5$ . If  $\mathbf{k} = \mathbb{F}_5$  assume further that  $G \pmod{5} = GL_2(\mathbb{F}_5)$ . Then  $H^1(G, \text{ad}^0(i)) = 0$  for  $i = 0, 1$ .
- (c) The restriction map  $H^1(G, \text{ad}^0(i)) \rightarrow H^1\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \text{ad}^0(i)\right)$  is an injection (for all  $p \geq 3$ ).

*Proof.* Part (a). The claim is likely to be familiar. Certainly the case  $\mathbf{k} = \mathbb{F}_p$  is well known and can be found in Serre's book [16].

We shall only verify that if  $G$  is a subgroup of  $SL_2(W/p^n)$  whose mod  $p^{n-1}$  reduction is  $SL_2(W/p^{n-1})$  then  $G = SL_2(W/p^n)$ . The kernel of the reduction map  $G \rightarrow SL_2(W/p^{n-1})$

consists of matrices of the form  $I + p^{n-1}A$  with  $A$  an element of some additive subgroup of  $\text{ad}^0$  stable under the action of  $G$ . Consequently either  $G = SL_2(W/p^n)$  or else the reduction map  $G \rightarrow SL_2(W/p^{n-1})$  is an isomorphism.

We will now discount the second possibility. So suppose that  $G \rightarrow SL_2(W/p^{n-1})$  is an isomorphism. Since  $G$  contains a transvection when  $p = 3$  we must have  $p \geq 3$ . Now let  $g \in G$  be the pre-image of  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in SL_2(W/p^{n-1})$ . So  $g$  must have order  $p^{n-1}$ . If we write  $g$  as  $(I + p^{n-1} \begin{pmatrix} a & b \\ c & -a \end{pmatrix}) \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ , a simple calculation using (2.3) shows that  $g^p = \begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$  and so  $g$  has order  $p^n$ —a contradiction.

*Part (b).* The hypothesis implies that  $H^1(G \bmod p, \text{ad}^0(i)) = 0$ . So let's assume that  $n \geq 2$  and that  $H^1(G \bmod p^{n-1}, \text{ad}^0(i)) = 0$ , and suppose that  $0 \neq \xi \in H^1(G, \text{ad}^0(i))$ . Then the restriction  $\xi$  to  $H := \ker(G \rightarrow G \bmod p^{n-1})$  is a group homomorphism compatible with the action of  $SL_2(W/p^{n-1})$ . It follows from part (a) that  $H$  is in fact  $\ker(SL_2(W/p^n) \rightarrow SL_2(W/p^{n-1}))$ . Since  $H$  is naturally identified with  $\text{ad}^0$ , it follows that  $\xi|_H$  is an isomorphism. Let's denote by  $W^\nu$  the ring  $W/p^{n-1} \oplus \mathbf{k}\epsilon$  where  $\epsilon^2 = p\epsilon = 0$ . (Or equivalently  $W^\nu \cong W[\epsilon]/(p^{n-1}, \epsilon^2, p\epsilon)$ .) We then see that the homomorphism  $SL_2(W/p^n) \rightarrow SL_2(W^\nu)$  given by

$$g \rightarrow (I + \epsilon\xi(g))(g \bmod p^{n-1})$$

is an isomorphism. To finish off, we proceed as in part (a): The transvection  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in SL_2(W^\nu)$  has order  $p^{n-1}$  while its pre-image in  $SL_2(W/p^n)$ , a matrix of the form  $(I + p^{n-1} \begin{pmatrix} a & b \\ c & -a \end{pmatrix}) \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ , has order  $p^n$ .

*Part (c).* Suppose  $0 \neq \xi \in H^1(G, \text{ad}^0(i))$  restricts to a trivial cohomology class in  $H^1(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \text{ad}^0(i))$ . Then the restriction of  $\xi$  to  $\begin{pmatrix} 1 & p^{n-1} \\ 0 & 1 \end{pmatrix}$  is trivial. Set  $N := \ker(G \rightarrow G \bmod p^{n-1})$ . Then  $\xi|_N$  has a non-trivial kernel, and hence  $\xi|_N$  is trivial. Thus  $\xi$  is a non-zero element of  $H^1(G \bmod p^{n-1}, \text{ad}^0(i))$ . We are thus reduced to the case when  $n = 1$ . Now  $H^1(SL_2(\mathbf{k}), \text{ad}^0) = 0$  except when  $\mathbf{k} = \mathbb{F}_5$ , so we are reduced to the case when  $G$  is a subgroup of  $GL_2(\mathbb{F}_5)$  containing  $SL_2(\mathbb{F}_5)$ . But in this case  $(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix})$  is the Sylow 5-subgroup of  $G$ , and hence if  $\xi|_{\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}} = 0$  then  $\xi = 0$ .  $\square$

### 3. CONSTRUCTING CHARACTERISTIC 0 LIFTS OF $\bmod p^n$ GALOIS REPRESENTATIONS

We can now formulate precise conditions under which a given  $\bmod p^n$  Galois representation can be lifted to characteristic 0, and use the lifts constructed to prove the existence of weak companion forms.

**3.1. Deformations of  $\bmod p^n$  representations to  $W(k)$ .** We now suppose we are given a totally real number field  $F$  and continuous odd representations  $\bar{\rho} : G_F \rightarrow GL_2(\mathbf{k})$ ,  $\rho_n : G_F \rightarrow GL_2(W/p^n)$ ,  $n \geq 2$ , with  $\bar{\rho} = \rho_n \bmod p$ . We shall also assume that the  $\bar{\rho}, \rho_n$  satisfy the following.

**Hypothesis A.** *The image of  $\bar{\rho}$  contains  $SL_2(\mathbf{k})$ . Furthermore, if  $p = 3$  then the image of  $\rho_n$  contains the transvection  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ .*

Fix a character  $\epsilon : G_F \rightarrow W^\times$  lifting the determinant of  $\rho_n$ . We wish to consider global deformation conditions  $\mathcal{D}$  for  $\bar{\rho}$  with determinant  $\epsilon$  such that  $\rho_n$  is a deformation of type  $\mathcal{D}$ . We shall abbreviate this and call  $\mathcal{D}$  a deformation condition for  $\rho_n$ . Except for a change in choice of lettering for primes of  $F$  we keep the notation of [11]. Thus  $\mathcal{D}_q$  is the local component at a prime  $q$ ,  $\mathfrak{t}_{\mathcal{D}_q}$  is the tangent space there, and  $\mathfrak{t}_{\mathcal{D}_q}^\perp \subseteq H^1(G_{F_q}, \text{ad}^0 \bar{\rho}(1))$  is the orthogonal complement of  $\mathfrak{t}_{\mathcal{D}_q}$  under the pairing induced by

$$\text{ad}^0 \bar{\rho} \times \text{ad}^0 \bar{\rho}(1) \xrightarrow{\text{trace}} \mathbf{k}(1).$$

The tangent space for  $\mathcal{D}$  is the Selmer group  $H_{\{\mathfrak{t}_{\mathcal{D}_q}\}}^1(F, \text{ad}^0\bar{\rho})$ ; the dual Selmer group  $H_{\{\mathfrak{t}_{\mathcal{D}_q}^\perp\}}^1(F, \text{ad}^0\bar{\rho}(1))$  is determined by the local conditions  $\mathfrak{t}_{\mathcal{D}_q}^\perp$ . (See for instance [15, Definition 8.6.19].) We also set

$$\delta(\mathcal{D}) := \dim_{\mathbf{k}} H_{\{\mathfrak{t}_{\mathcal{D}_q}\}}^1(F, \text{ad}^0\bar{\rho}) - \dim_{\mathbf{k}} H_{\{\mathfrak{t}_{\mathcal{D}_q}^\perp\}}^1(F, \text{ad}^0\bar{\rho}(1)).$$

**Proposition 3.1.** *Suppose we are given a deformation condition  $\mathcal{D}$  for  $\rho_n$  with determinant  $\epsilon$ . Let  $S$  be a fixed finite set of primes of  $F$  including primes where  $\mathcal{D}$  is ramified and all the infinite primes. If  $\delta(\mathcal{D}) \geq 0$  we can find a deformation condition  $\mathcal{E}$  for  $\rho_n$  with determinant  $\epsilon$  such that:*

- The local conditions  $\mathcal{E}_q$  and  $\mathcal{D}_q$  are the same at primes  $q \in S$ ;
- $\mathcal{E}_q$  is a substantial deformation condition for  $q \notin S$ ; and,
- $H_{\{\mathfrak{t}_{\mathcal{E}_q}^\perp\}}^1(F, \text{ad}^0\bar{\rho}(1)) = (0)$ .

*Proof.* Let  $K$  be the splitting field of  $\rho_n$  adjoined  $p^n$ -th roots of unity. We claim that we can find elements  $g, h \in \text{Gal}(K/F)$  such that

- (R1)  $\rho_n(g) \sim \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $\chi(g) = -1 \pmod{p^n}$ ;
- (R2)  $\rho_n(h) \sim a \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $\chi(h) = 1 \pmod{p^n}$ .

For R1, we can take  $g$  to be complex conjugation. For R2, by considering  $\epsilon = \chi(\epsilon\chi^{-1})$  or otherwise, we can write  $\epsilon = \chi\epsilon_0\epsilon_1^2$  where  $\epsilon_0$  is a finite order character of order co-prime to  $p$ . Our assumptions on the size of  $\bar{\rho}$  and  $\rho_n$  (when  $p = 3$ ) along with Proposition 2.8 imply that the image of the twist of  $\rho_n \otimes \epsilon_1^{-1}$  contains  $SL_2(W/p^n)$ . Thus we can find  $h_1 \in \text{Gal}(K/F)$  such that  $\rho_n(h_1) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \epsilon_1(h_1)$  and we get  $\epsilon_0(h_1)\chi(h_1) = 1$ . We can then take  $h$  to be  $h_1^{p^k-1}$  where  $p^k$  is the cardinality of  $\mathbf{k}$ .

We first adjust  $\mathcal{D}$  and define a deformation condition  $\mathcal{E}_0$  for  $\rho_n$  with determinant  $\epsilon$  as follows. We make no change if  $p \geq 5$  and the projective image of  $\bar{\rho}$  strictly contains  $PSL_2(\mathbb{F}_5)$ ; so  $\mathcal{E}_0$  is  $\mathcal{D}$ . Now for the remaining cases: Suppose that either  $p = 3$  or the projective image of  $\bar{\rho}$  is  $A_5$  (so  $\mathbf{k}$  is necessarily  $\mathbb{F}_5$ ). Using the Chebotarev Density Theorem and R2 above, we can find a prime  $q_0 \notin S$  with  $q_0 \equiv 1 \pmod{p^n}$  and  $\rho_n(\text{Frob}_{q_0}) = a \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . Let  $\mathcal{E}_0$  be the deformation condition of  $\bar{\rho}$  with determinant  $\epsilon$  characterized by the following local conditions:

- at primes  $q \neq q_0$ ,  $\mathcal{E}_{0q} = \mathcal{D}_q$ ;
- at  $q_0$ ,  $\mathcal{E}_{0q_0}$  consists of deformations of the form

$$\begin{pmatrix} \chi & * \\ 0 & 1 \end{pmatrix} \epsilon'$$

where  $\epsilon' : G_{v_0} \rightarrow W^\times$  is unramified and  $\epsilon|_{G_{q_0}} = \chi\epsilon'^2$ .

By our choice of  $q_0$ ,  $\mathcal{E}_0$  is a deformation condition for  $\rho_n$ . Further,  $\mathcal{E}_{0q_0}$  is a substantial deformation and all non-zero cohomology classes in  $\mathfrak{t}_{\mathcal{E}_{0q_0}}, \mathfrak{t}_{\mathcal{E}_{0q_0}}^\perp$  are ramified.

We claim that the restriction maps

$$H_{\{\mathfrak{t}_{\mathcal{E}_{0q}}\}}^1(F, \text{ad}^0\bar{\rho}) \rightarrow H^1(G_K, \text{ad}^0\bar{\rho}) \quad \text{and} \quad H_{\{\mathfrak{t}_{\mathcal{E}_{0q}}^\perp\}}^1(F, \text{ad}^0\bar{\rho}(1)) \rightarrow H^1(G_K, \text{ad}^0\bar{\rho}(1))$$

are injective. When  $p \geq 5$  and the projective image of  $\bar{\rho}$  strictly contains  $A_5$  an easy calculation using Proposition 2.8 shows that  $H^1(\text{Gal}(K/F), \text{ad}^0\bar{\rho})$  and  $H^1(\text{Gal}(K/F), \text{ad}^0\bar{\rho}(1))$  are trivial, and so the injectivity follows. In the case when  $p = 3$  or the projective image of  $\bar{\rho}$  is  $A_5$ , we argue as follows: If  $\xi \in \ker\left(H_{\{\mathfrak{t}_{\mathcal{E}_{0q}}\}}^1(F, \text{ad}^0\bar{\rho}) \rightarrow H^1(G_K, \text{ad}^0\bar{\rho})\right)$ , then  $\xi$  is naturally an element of  $H^1(\text{Gal}(K/F), \text{ad}^0\bar{\rho})$ . Thus  $\xi$  is unramified at  $q_0$  and so the restriction of  $\xi$  to the decomposition

group at  $\mathfrak{q}_0$  must be trivial. Using Proposition 2.8 it follows that  $\xi \in H^1(\text{Gal}(K/F), \text{ad}^0 \bar{\rho})$  is trivial. A similar argument works for  $\text{ad}^0 \bar{\rho}(1)$ .

The proof is now standard: If the dual Selmer group for  $\mathcal{E}_0$  is non-trivial then we can find

$$0 \neq \xi \in H^1_{\{\mathfrak{t}_{\varepsilon_0 \mathfrak{q}}\}}(F, \text{ad}^0 \bar{\rho}), \quad 0 \neq \psi \in H^1_{\{\mathfrak{t}_{\varepsilon_0 \mathfrak{q}}\}}(F, \text{ad}^0 \bar{\rho}(1)).$$

Take  $g \in \text{Gal}(K/L)$  as in R1, consider pairs  $(M_1, N_1), (M_2, N_2)$  where  $\{(\begin{smallmatrix} 0 & * \\ * & 0 \end{smallmatrix})\} = N_1 \subset M_1 = \text{ad}^0 \bar{\rho}$ ,  $\{(\begin{smallmatrix} * & * \\ 0 & * \end{smallmatrix})\} = N_2 \subset M_2 = \text{ad}^0 \bar{\rho}(1)$  and apply Proposition 2.2 of [11]. One can then find a prime  $\mathfrak{r} \notin S \cup \{\mathfrak{q}_0\}$  lifting  $g$  such that the restrictions of  $\xi, \psi$  to  $G_{\mathfrak{r}}$  are not in  $H^1(G_{\mathfrak{r}}, N_1), H^1(G_{\mathfrak{r}}, N_2)$ .

Now take  $\mathcal{E}_1$  to be the deformation condition with determinant  $\epsilon$  as follows:  $\mathcal{E}_1$  and  $\mathcal{E}_0$  differ only at  $\mathfrak{r}$ , and at  $\mathfrak{r}$ , the local component consists of deformations of the form  $(\begin{smallmatrix} \chi & * \\ 0 & 1 \end{smallmatrix}) (\epsilon/\chi)^{1/2}$  considered in Example 2.3. Here,  $(\epsilon/\chi)^{1/2}$  is the unramified character determined by taking the square-root of  $\epsilon(\text{Frob}_{\mathfrak{r}})\chi^{-1}(\text{Frob}_{\mathfrak{r}})$ . Since  $\text{Frob}_{\mathfrak{r}}$  lifts  $g$  we have  $\chi(\text{Frob}_{\mathfrak{r}}) \equiv -1 \pmod{p^n}$ , and consequently  $\mathcal{E}_1$  is a substantial deformation condition for  $\rho_n$ . The rest is identical to the proof of Proposition 4.2, [11]: The dual Selmer group for  $\mathcal{E}_1$  has dimension one less than that of the dual Selmer group for  $\mathcal{E}_0$ . (Of course  $\delta(\mathcal{E}_1) = \delta(\mathcal{E}_0) = \delta(\mathcal{D})$ .)  $\square$

We can now prove a general result for lifting a  $\pmod{p^n}$  representation to characteristic 0.

**Theorem 3.2.** *Let  $\mathcal{D}$  be a deformation condition for  $\rho_n$  with determinant  $\epsilon$ , and let  $S$  be a fixed finite set of primes of  $F$  including primes where  $\mathcal{D}$  is ramified and all the infinite primes. Suppose that each local component is substantial. We can then find a deformation condition  $\mathcal{E}$  for  $\rho_n$  with determinant  $\epsilon$  such that:*

- The local conditions  $\mathcal{E}_{\mathfrak{q}}$  and  $\mathcal{D}_{\mathfrak{q}}$  are the same at primes  $\mathfrak{q} \in S$ ;
- Each local component is a substantial deformation condition;
- The dual Selmer group  $H^1_{\{\mathfrak{t}_{\varepsilon_{\mathfrak{q}}}\}}(F, \text{ad}^0 \bar{\rho}(1))$  is trivial.

$\mathcal{E}$  is a smooth deformation condition and the universal deformation ring is a power series ring over  $W$  in  $\delta(\mathcal{D})$  variables. In particular, there is a representation  $\rho : G_F \rightarrow GL_2(W)$  of type  $\mathcal{E}$  lifting  $\rho_n$ .

*Proof.* The only verification required is to check that  $\delta(\mathcal{D}) \geq 0$  and that  $\dim_{\mathbf{k}} H^1_{\{\mathfrak{t}_{\varepsilon_{\mathfrak{q}}}\}}(F, \text{ad}^0 \bar{\rho}) = \delta(\mathcal{E}) = \delta(\mathcal{D})$ . This is done using Wiles' formula (cf [15, Theorem 8.6.20]).  $\square$

**3.2. Modular characteristic 0 lifts and proof of Main Theorem.** We now look at the question of producing characteristic zero liftings which are modular. Given a  $\pmod{p^n}$  Galois representation  $\rho_n : G_F \rightarrow GL_2(W/p^n)$  with  $\rho_n \pmod{p}$  modular, when can we guarantee the existence of a modular form  $f$  with  $\rho_{f,p} \pmod{p^n} \sim \rho_n$ ? Our answer is a modest attempt using Theorem 3.2 to produce a characteristic 0 lift and then invoking results of Skinner and Wiles [17] to prove that it is modular.

For the rest of this section,  $F$  is a totally real field and  $\psi : G_F \rightarrow W^\times$  is a finite order character of  $G_F$  unramified at primes dividing  $p$ .

**Proposition 3.3.** *Let  $\rho_n : G_F \rightarrow GL_2(W/p^n)$  be a continuous odd representation satisfying 3.1. Suppose  $\epsilon := \psi\chi^a, a \geq 1$  lifts the determinant of  $\rho_n$ . Assume that:*

- (i) *At a prime  $\mathfrak{q} \nmid p$  where  $\rho_n$  is ramified, the restriction  $\rho_n|_{G_{\mathfrak{q}}}$  is substantial there and that a substantial deformation condition  $\mathcal{D}_{\mathfrak{q}}$  is specified for  $\rho_n$ .*

(ii) At a prime  $\mathfrak{p}$  dividing  $p$ ,

$$\rho_n|_{G_{\mathfrak{p}}} \sim \begin{pmatrix} \chi^a \psi_{1\mathfrak{p}} & * \\ 0 & \psi_{2\mathfrak{p}} \end{pmatrix}$$

where  $\psi_{1\mathfrak{p}}, \psi_{2\mathfrak{p}}$  are unramified,  $\chi^a \psi_{1\mathfrak{p}} \not\equiv \psi_{2\mathfrak{p}} \pmod{p}$  and  $\chi^a \psi_{1\mathfrak{p}} \not\equiv \chi \psi_{2\mathfrak{p}} \pmod{p}$ .

(iii) There is an ordinary, parallel weight at least 2, modular form which is a  $(\psi_{2\mathfrak{p}} \pmod{p})$ -good lift of  $\rho_n \pmod{p}$ .

There is then a modular form  $f$  such that its associated  $p$ -adic representation  $\rho_{f,p} : G_F \rightarrow GL_2(W)$  lifts  $\rho_n$ , has determinant  $\psi \chi^a$ , is of type  $\mathcal{D}_{\mathfrak{q}}$  at primes  $\mathfrak{q} \nmid p$  where  $\rho_n$  is ramified, and

$$\rho_{f,p}|_{G_{\mathfrak{p}}} \sim \begin{pmatrix} \psi'_{1\mathfrak{p}} \chi^a & * \\ 0 & \psi'_{2\mathfrak{p}} \end{pmatrix}$$

at primes  $\mathfrak{p}|p$  with  $\psi'_{2\mathfrak{p}}$  an unramified lift of  $\psi_{2\mathfrak{p}} \pmod{p}$ .

*Proof.* At a prime  $\mathfrak{p}|p$  take  $\mathcal{D}_{\mathfrak{p}}$  to be the class of deformations of the form

$$\begin{pmatrix} \psi'_{1\mathfrak{p}} \chi^a & * \\ 0 & \psi'_{2\mathfrak{p}} \end{pmatrix}$$

where  $\psi'_{1\mathfrak{p}}$  (resp.  $\psi'_{2\mathfrak{p}}$ ) is an unramified lifting of  $\psi_{1\mathfrak{p}} \pmod{p}$  (resp.  $\psi_{2\mathfrak{p}} \pmod{p}$ ), and  $\psi_{1\mathfrak{p}} \psi_{2\mathfrak{p}} = \psi$ . This is a substantial deformation for  $\rho_n$  at  $\mathfrak{p}$  by Proposition 2.5. By Theorem 3.2, there is a smooth deformation condition  $\mathcal{E}$  for  $\rho_n$  which agrees with  $\mathcal{D}_{\mathfrak{p}}$  at primes above  $p$  and primes where  $\rho_n$  is ramified. Thus there is continuous representation  $\rho : G_F \rightarrow GL_2(W)$  with  $\rho \pmod{p^n} = \rho_n$ , unramified outside finitely many primes, determinant  $\psi \chi^a$  and  $\rho|_{I_{\mathfrak{p}}} \sim \begin{pmatrix} \chi^a & * \\ 0 & 1 \end{pmatrix}$  at primes  $\mathfrak{p}|p$ . The proposition now follows from Skinner-Wiles [17].  $\square$

*Proof of Main Theorem.* Let's recall the set up: We are given a Hilbert modular newform  $f$  of weight  $k \geq 2$  character  $\psi$  which is ordinary at  $p$  and whose reduction  $\pmod{p^n}$  gives  $\rho_{f,n} : G_F \rightarrow GL_2(W/p^n)$ . For each prime  $\mathfrak{p}$  of  $F$  over  $p$ , let  $\psi_{1\mathfrak{p}}, \psi_{2\mathfrak{p}}$  be the unramified characters such that

$$\rho_f|_{G_{\mathfrak{p}}} \sim \begin{pmatrix} \psi_{1\mathfrak{p}} \chi^{k-1} & * \\ 0 & \psi_{2\mathfrak{p}} \end{pmatrix}.$$

As  $\psi_{1\mathfrak{p}} \psi_{2\mathfrak{p}} = \psi$  and  $\psi_{2\mathfrak{p}}(\text{Frob}_{\mathfrak{p}}) = c(\mathfrak{p}, f)$ , hypothesis LC1 ensures  $\psi_{1\mathfrak{p}}, \psi_{2\mathfrak{p}}$  are distinct modulo  $\wp$ . From this, one deduces easily that if  $f$  has a weak companion form  $\pmod{p^n}$  then  $\rho_{f,n}$  splits at  $p$ . We now show that 'split at  $p$ ' implies the existence of a weak companion form.

Let  $\rho_n := \rho_{f,n} \otimes \chi^{1-k}$ , and set  $\bar{\rho} := \rho_n \pmod{p}$ . Recall that  $k' \geq 2$  is the smallest integer satisfying the congruence  $k + k' \equiv 2 \pmod{(p-1)p^{n-1}}$ . Define a global deformation condition  $\mathcal{D}$  for  $\bar{\rho} \otimes \chi^{1-k} \pmod{p}$  by the following requirements:

(a) Deformations are unramified outside primes dividing  $pn$  and have determinant  $\psi \chi^{k'-1}$ .

(b) At a prime  $\mathfrak{p} \mid p$ , the local condition  $\mathcal{D}_{\mathfrak{p}}$  consists of deformations of the form

$$\begin{pmatrix} \psi'_{2\mathfrak{p}}\chi^{k'-1} & * \\ 0 & \psi'_{1\mathfrak{p}} \end{pmatrix}$$

where  $\psi'_{1\mathfrak{p}}$  (resp.  $\psi'_{2\mathfrak{p}}$ ) is an unramified lifting of  $\psi_{1\mathfrak{p}} \pmod p$  (resp.  $\psi_{2\mathfrak{p}} \pmod p$ ), and  $\psi_{1\mathfrak{p}}\psi_{2\mathfrak{p}} = \psi$ .

(c) Let  $\mathfrak{q}$  be a prime dividing  $n$ , the level of  $f$ . We need to distinguish two cases:

- (i) If  $\mathfrak{q}$  does not divide the conductor of  $\psi$  then  $\bar{\rho}|_{G_{\mathfrak{q}}} \sim \begin{pmatrix} \bar{\chi} & * \\ 0 & 1 \end{pmatrix} \bar{\epsilon}$  for some character  $\bar{\epsilon}$ . Further, hypothesis LC2 ensures that if  $\bar{\rho}|_{G_{\mathfrak{q}}}$  is semisimple then  $\bar{\chi} \neq 1$ . We then take  $\mathcal{D}_{\mathfrak{q}}$  to be local liftings with determinant  $\psi\chi^{k'-1}$  of the type considered in Example 2.3.
- (ii) If  $\mathfrak{q}$  divides the conductor of  $\psi$  then  $\rho_f(I_{\mathfrak{q}}), \bar{\rho}(I_{\mathfrak{q}})$  are finite and have the same order. In this case we take  $\mathcal{D}_{\mathfrak{q}}$  as in Example 2.2 i.e. lifts with determinant  $\psi\chi^{k'-1}$  which factor through  $G_{\mathfrak{q}}/(I_{\mathfrak{q}} \cap \ker \bar{\rho})$ .

It then follows that  $\rho_n$  is a deformation of type  $\mathcal{D}$  and that at each prime  $\mathfrak{q} \nmid p$  where  $\rho_n$  is ramified the local deformation condition  $\mathcal{D}_{\mathfrak{q}}$  is substantial there. As  $p$  is unramified in  $F$ , the distinctness of  $\psi_{1\mathfrak{p}}, \psi_{2\mathfrak{p}}$  modulo  $p$  implies that  $\bar{\rho}$  satisfies hypothesis (ii) of Proposition 3.3. From the existence of  $\pmod p$  companion forms, ([5, Theorem 2.1]), it follows that  $\bar{\rho}$  has an ordinary modular lift which is  $(\psi_{1\mathfrak{p}} \pmod p)_{\mathfrak{p}|p}$ -good. The existence of a  $\pmod p^n$  weak companion form  $g$  for  $f$  of weight  $k'$  character  $\psi$  now follows from Proposition 3.3.  $\square$

#### 4. CHECKING LOCAL SPLITTING: A COMPUTATIONAL APPROACH

The lifting result of the previous section is not suitable for computational purposes in general because, except in the case when dual Selmer group was already trivial, we had no control of the level. There is, however, one case when we do have absolute control. We now describe this situation and go on to verify examples of local splitting.

**4.1. A special case.** Suppose  $\bar{\rho} : G_{\mathbb{Q}} \rightarrow GL_2(\mathbf{k})$  is absolutely irreducible and  $\mathcal{D}$  is a deformation condition for  $\bar{\rho}$  such that its tangent space is 0 dimensional. Then the universal deformation ring  $R_{\mathcal{D}}$  is a quotient of  $W(\mathbf{k})$ . If we also knew that there is a characteristic 0 lift of type  $\mathcal{D}$ , then we must have  $R_{\mathcal{D}} \simeq W$ . Consequently any  $\pmod p^n$  representation of type  $\mathcal{D}$  lifts to characteristic 0.

The question now is: How can one check if the tangent space is 0 dimensional? Observe that we must necessarily have exactly one characteristic 0 lift of type  $\mathcal{D}$ . This alone might not be enough though. For instance,  $R_{\mathcal{D}}$  might be  $W[X]/(X^2)$ .

To proceed further, and with the examples we have in mind, we shall assume that  $\bar{\rho} : G_{\mathbb{Q}} \rightarrow GL_2(\mathbf{k})$  is an absolutely irreducible representation with determinant  $\bar{\chi}$  such that

- $\bar{\rho}|_{G_p} \sim \begin{pmatrix} \bar{\chi}\psi^{-1} & * \\ 0 & \psi \end{pmatrix}$ , with  $\psi$  unramified and  $\psi \neq \psi^{-1}$ ,
- if  $q \nmid p$  then  $\#\bar{\rho}(I_{\mathfrak{q}}) \mid p$ .

By Lemma 3.24 of [3],  $\bar{\rho}|_{G_L}$  is absolutely irreducible where  $L = \mathbb{Q}(\sqrt{(-1)^{(p-1)/2}p})$ .

Let  $N$  be the Artin conductor of  $\bar{\rho}$ . For an integer  $k \geq 2$ , let  $S(k, N, \bar{\rho})$  be the (possibly empty) set of newforms of level  $N$  with  $\rho_f \pmod p \simeq \bar{\rho}$ .

With notation as in Theorem 3.42 of [3], we then have an isomorphism  $R_{\emptyset} \xrightarrow{\sim} \mathbb{T}_{\emptyset}$ , where  $R_{\emptyset}$  is the universal deformation ring for minimally ramified ordinary lifts and  $\mathbb{T}_{\emptyset}$  is the reduced Hecke

algebra generated by the Fourier coefficients of newforms in  $S(2, N, \bar{\rho})$ . In particular, the dimension of the tangent space in the minimally ramified case is 0 if and only if  $\#S(2, N, \bar{\rho}) = 1$ .

For  $n \geq 1$  set  $k_n := (p-1)p^{n-1} - (p-1) + 2$  and define a deformation condition  $\mathcal{D}_{k_n}$  for  $\bar{\rho}$  as follows: A lift  $\rho : G_{\mathbb{Q}} \rightarrow GL_2(A)$  is a deformation of type  $\mathcal{D}_{k_n}$  if

- $\det \rho = \chi^{k_n-1}$  and  $\rho$  is unramified outside primes dividing  $N$ ,
- at primes  $q|N$ ,  $\rho|_{G_q} \sim \begin{pmatrix} \chi & * \\ 0 & 1 \end{pmatrix}$  up to twist, and
- at  $p$ ,  $\rho|_{G_p} \sim \begin{pmatrix} \tilde{\psi}^{-1}\chi^{k_n-1} & * \\ 0 & \tilde{\psi} \end{pmatrix}$ , where  $\tilde{\psi}$  is an unramified lift of  $\psi$ .

Note that for  $n = 1$  the universal deformation ring  $R_{\mathcal{D}_{k_n}}$  is  $R_{\emptyset} \simeq \mathbb{T}_{\emptyset}$ . Clearly, the type  $\mathcal{D}_{k_n}$  deformations to  $\mathbf{k}[\epsilon]/(\epsilon^2)$  are in bijection with type  $\mathcal{D}_2$  deformations. Hence if the tangent space of  $\mathcal{D}_2$  has dimension 0 then so does  $\mathcal{D}_{k_n}$ . We conclude that  $R_{\mathcal{D}_{k_n}} \simeq W$ , corresponding to a unique newform in  $S(k_n, N, \bar{\rho})$ .

**Proposition 4.1.** *Let  $f$  be a newform of weight  $k \geq 2$ , level  $N$ , trivial character and ordinary at  $p$ , such that*

- $\bar{\rho}_f$  is absolutely irreducible,
- the conductor of  $\bar{\rho}_f$  is  $N$ ,
- $\bar{\rho}|_{G_p} \sim \begin{pmatrix} * & * \\ 0 & \psi \end{pmatrix}$  with  $\psi$  unramified and  $\psi^2 \neq 1$ ,
- if  $q \nmid p$  then  $\#\bar{\rho}_f(I_q)|p$ .

*Assume that  $p-1|k$  and that  $f$  has exactly one companion form mod  $p$ . Then  $\rho_f$  splits mod  $p^n$  iff  $f$  has a companion form mod  $p^n$ .*

*Proof.* The proposition is immediate from the preceding discussion by taking  $\bar{\rho}$  to be  $\bar{\rho}_f \otimes \bar{\chi}^{1-k}$ .  $\square$

**4.2. Greenberg's conjecture.** In this section we give some examples of the existence (or non-existence) of higher companion forms. We shall restrict ourselves to the setting of classical elliptic modular forms as we only give examples in this case.

Recall that a newform  $f$  is said to have **complex multiplication**, or just CM, by a quadratic character  $\phi : G_{\mathbb{Q}} \rightarrow \{\pm 1\}$  if  $T(q)f = \phi(\text{Frob}_q)c(q, f)f$  for almost all primes  $q$ . We will also refer to CM by the corresponding quadratic extension. It is well known that a modular form has CM if and only if its associated  $p$ -adic representation is induced from an algebraic Hecke character.

Let  $\mathcal{O}$  be a  $p$ -adic integer ring with residue field  $\mathbf{k}$ . Suppose the newform  $f$  is  $p$ -ordinary and  $\rho_f : G_{\mathbb{Q}} \rightarrow GL_2(\mathcal{O})$ . Then, as indicated in the introduction,  $\rho_f|_{G_p}$  can be assumed to be upper triangular with an unramified lower diagonal entry and this leads to the natural question of determining when  $\rho_f$  splits at  $p$ .

There is a well-known conjectural connection between  $\rho_f$  to be split at  $p$  and  $f$  to have CM. The antecedents are sketchy, but Hida [9] calls it Greenberg's local non-semisimplicity conjecture; we will simply refer to it as Greenberg's conjecture which asserts:

If  $f$  is ordinary at  $p$  and  $\rho_f$  splits at  $p$  then  $f$  has complex multiplication.

This is satisfactorily known for modular forms over  $\mathbb{Q}$  of weight 2. For higher weights, the question remains largely unresolved although some interesting results involving Hida families are shown in Ghate [6] which also has a survey of results for weight 2. The analogous problem for  $\Lambda$ -adic modular forms was resolved in Ghate-Vatsal [7] by using deformation theory but similar methods appear not to bear fruit in the classical case. Emerton [4] shows how this conjecture would follow from a  $p$ -adic version of the variational Hodge conjecture. Through the main theorem and proposition 4.1, higher congruence companions offer a slightly different perspective to the question of  $\rho_f$  splitting at  $p$ .

To describe this further, let  $N$  be the level of the  $p$ -ordinary newform  $f$ . Assume that  $f$  has trivial character and has weight  $p - 1$ . For each positive integer  $n$  we set  $k_n := p^{n-1}(p - 1) - (p - 1) + 2$ . We then proceed as follows:

- (a) Check that  $f$  has a companion form  $\pmod{p}$ . Check congruences to make sure that the residual representation  $\rho_f \pmod{\pi}$  is absolutely irreducible and that  $c(p, f) \not\equiv \pm 1 \pmod{\pi}$ . We can therefore write

$$\rho_f \pmod{\pi} = \begin{pmatrix} \bar{\chi}^{k-1} \bar{\psi} & 0 \\ 0 & \bar{\psi}^{-1} \end{pmatrix}$$

with  $\bar{\psi}^{-1} \neq \bar{\psi}$ .

- (b) In order to be able to check fewer cases, ensure that  $\rho_f$  is minimally ramified i.e. the Artin conductor of  $\rho_f \pmod{\pi}$  is  $N$ .
- (c) Set  $\bar{\rho} := \rho_f \otimes \chi^{1-k} \pmod{\pi}$ . For each  $n \geq 1$  let  $\mathcal{D}_{k_n}$  be the weight  $k_n$ , trivial character deformation condition as described in section 4.1. Check if the tangent spaces can be taken to be 0 dimensional. Thus we have to check if  $f$  has precisely one companion form of type  $\mathcal{D}_{k_1}$ . We then apply proposition 4.1 to deduce that  $\rho_f$  splits  $\pmod{p^n}$  if and only if  $f$  has a companion form  $\pmod{p^n}$  i.e. there is a newform  $g$  of level  $N$ , trivial character, weight  $k_n$  such that  $f \equiv g \otimes \chi^{k-1} \pmod{p^n}$

We check Greenberg's conjecture explicitly for two known non- $CM$  forms of weight 4. The computations were done on MAGMA. In both cases  $p = 5$ . We note that in these examples, taking  $N$  to be the level of  $f$ , one may check its companionship with a form  $g$  "by hand" by simply verifying the congruences  $c(f, m) \equiv m^3 c(g, m) \pmod{5^n}$  for  $(m, 5N) = 1$  up to the Sturm bound.

**Example 4.2.** Let  $f$  be the newform of weight 4, level 21 and trivial character with the following Fourier expansion:

$$g = q - 3q^2 - 3q^3 + q^4 - 18q^5 + 9q^6 + 7q^7 + 21q^8 + 9q^9 + 54q^{10} - 36q^{11} - 3q^{12} - 34q^{13} - 21q^{14} + 54q^{15} - 71q^{16} + 42q^{17} - 27q^{18} - 124q^{19} - 18q^{20} - 21q^{21} + 108q^{22} - 63q^{24} + 199q^{25} + 102q^{26} - 27q^{27} + 7q^{28} + 102q^{29} - 162q^{30} - 160q^{31} + 45q^{32} + 108q^{33} - 126q^{34} - 126q^{35} + 9q^{36} + 398q^{37} + 372q^{38} + 102q^{39} - 378q^{40} - 318q^{41} + 63q^{42} - 268q^{43} - 36q^{44} - 162q^{45} + 240q^{47} + 213q^{48} + 49q^{49} - 597q^{50} + \dots$$

MAGMA outputs modulo 5, a unique companion form  $g$  weight 2, level 21 and trivial character with the following Fourier expansion:

$$f = q - q^2 + q^3 - q^4 - 2q^5 - q^6 - q^7 + 3q^8 + q^9 + 2q^{10} + 4q^{11} - q^{12} - 2q^{13} + q^{14} - 2q^{15} - q^{16} - 6q^{17} - q^{18} + 4q^{19} + 2q^{20} - q^{21} - 4q^{22} + 3q^{24} - q^{25} + 2q^{26} + q^{27} + q^{28} - 2q^{29} + 2q^{30} - 5q^{32} + 4q^{33} + 6q^{34} + 2q^{35} - q^{36} + 6q^{37} - 4q^{38} - 2q^{39} - 6q^{40} + 2q^{41} + q^{42} - 4q^{43} - 4q^{44} - 2q^{45} - q^{48} + q^{49} + q^{50} + \dots$$

Clearly there are no companions of weight 2 and level 3 or 7. Modulo  $5^2$ ,  $f$  has no companion forms of weight 18, level dividing 21 and trivial character. Thus  $f$  does not split  $\pmod{5^2}$ .

**Example 4.3.** Let  $f$  be the newform of weight 4, level 57 and trivial character with Fourier expansion

$$f = q - q^2 + 3q^3 - 7q^4 - 12q^5 - 3q^6 - 20q^7 + 15q^8 + 9q^9 + 12q^{10} - 4q^{11} - 21q^{12} - 76q^{13} + 20q^{14} - 36q^{15} + 41q^{16} + 22q^{17} - 9q^{18} - 19q^{19} + 84q^{20} - 60q^{21} + 4q^{22} + 82q^{23} + 45q^{24} + 19q^{25} + 76q^{26} + 27q^{27} + 140q^{28} + 242q^{29} + 36q^{30} - 126q^{31} - 161q^{32} - 12q^{33} - 22q^{34} + 240q^{35} - 63q^{36} - 180q^{37} + 19q^{38} - 228q^{39} - 180q^{40} - 390q^{41} + 60q^{42} + 308q^{43} + 28q^{44} - 108q^{45} - 82q^{46} - 522q^{47} + 123q^{48} + 57q^{49} - 19q^{50} + \dots$$

It has a unique mod 5 companion form  $g$  of weight 2, level 57 and trivial character with Fourier expansion

$$g = q - 2q^2 - q^3 + 2q^4 - 3q^5 + 2q^6 - 5q^7 + q^9 + 6q^{10} + q^{11} - 2q^{12} + 2q^{13} + 10q^{14} + 3q^{15} - 4q^{16} - q^{17} - 2q^{18} - q^{19} - 6q^{20} + 5q^{21} - 2q^{22} - 4q^{23} + 4q^{25} - 4q^{26} - q^{27} - 10q^{28} - 2q^{29} - 6q^{30} - 6q^{31} + 8q^{32} - q^{33} + 2q^{34} + 15q^{35} + 2q^{36} + 2q^{38} - 2q^{39} - 10q^{42} - q^{43} + 2q^{44} - 3q^{45} + 8q^{46} - 9q^{47} + 4q^{48} + 18q^{49} - 8q^{50} + \dots$$

and no other companions of level dividing 57. Modulo  $5^2$ ,  $f$  has no companion forms of weight 18, level dividing 57 and trivial character.

#### ACKNOWLEDGEMENTS

The first author is grateful to Paul Mezo and Yuly Billig of Carleton University, Gabor Wiese of IEM, Essen and Universität Regensburg for their postdoctoral support that brought this work to fruition. The authors also thank Frazer Jarvis and Neil Dummigan for their useful comments and suggestions and Panagiotis Tsaknias for his valuable assistance with the computations.

#### REFERENCES

- [1] G. Böckle. On the isomorphism  $R_\emptyset \rightarrow T_\emptyset$ . Appendix to “On isomorphisms between deformation and Hecke rings” by C. Khare. *Invent. Math.*, 154(1):217–222, 2003.
- [2] G. Böckle. Presentations of universal deformation rings. In *L-functions and Galois representations*, volume 320 of *London Math. Soc. Lecture Note Ser.*, pages 24–58. Cambridge Univ. Press, Cambridge, 2007.
- [3] H. Darmon, F. Diamond and R. Taylor. Fermat’s Last Theorem. In *Current developments in mathematics*, 1995 Ed. R. Bott et al., International Press, Boston 1995
- [4] M. Emerton. A  $p$ -adic variational hodge conjecture and modular forms with complex multiplication. *preprint*.
- [5] T. Gee. Companion forms over totally real fields. II. *Duke Math. J.*, 136(2):275–284, 2007.
- [6] E. Ghate. On the local behaviour of ordinary modular galois representations. *Progress in Mathematics*, 224:105–224, 2004.
- [7] E. Ghate and V. Vatsal. On the local behaviour of ordinary  $\Lambda$ -adic representations. *Ann. Inst. Fourier (Grenoble)*, 54(7):2143–2162 (2005), 2004.
- [8] B. H. Gross. A tameness criterion for Galois representations associated to modular forms (mod  $p$ ). *Duke Math. J.*, 61(2):445–517, 1990.
- [9] H. Hida. CM components of the big Hecke algebra. A series of lectures at Hokkaido University.
- [10] C. Khare and J.-P. Wintenberger. Serre’s modularity conjecture. I. *Invent. Math.*, 178(3):485–504, 2009.
- [11] J. Manoharmayum. Lifting Galois representations of number fields. *J. Number Theory*, 129(5):1178–1190, 2009.
- [12] B. Mazur. Deforming Galois representations. In *Galois groups over  $\mathbf{Q}$  (Berkeley, CA, 1987)*, volume 16 of *Math. Sci. Res. Inst. Publ.*, pages 385–437. Springer, New York, 1989.
- [13] B. Mazur. Deformation theory of Galois representations. In *Modular Forms and Galois Representations*. Eds G. Cornell, J. H. Silverman, G. Stevens. Springer, New York, 1997.
- [14] B. Mazur and A. Wiles. On  $p$ -adic analytic families of Galois representations. *Compositio Math.*, 59(2):231–264, 1986.
- [15] J. Neukirch, A. Schmidt, and K. Wingberg. *Cohomology of number fields*, volume 323 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2008.
- [16] J.-P. Serre. *Abelian  $l$ -adic representations and elliptic curves*, volume 7 of *Research Notes in Mathematics*. A K Peters Ltd., Wellesley, MA, 1998. With the collaboration of Willem Kuyk and John Labute, Revised reprint of the 1968 original.
- [17] C. M. Skinner and A. J. Wiles. Nearly ordinary deformations of irreducible residual representations. *Ann. Fac. Sci. Toulouse Math. (6)*, 10(1):185–215, 2001.
- [18] A. Wiles. On ordinary  $\lambda$ -adic representations associated to modular forms. *Invent. Math.*, 94(3):529–573, 1988.
- [19] A. Wiles. Modular elliptic curves and Fermat’s last theorem. *Ann. of Math. (2)*, 141(3):443–551, 1995.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, 93053 REGENSBURG, GERMANY  
*E-mail address:* [rajender.adibhatla@mathematik.uni-regensburg.de](mailto:rajender.adibhatla@mathematik.uni-regensburg.de)

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SHEFFIELD, SHEFFIELD S3 7RH, UNITED KINGDOM  
*E-mail address:* [j.manoharmayum@sheffield.ac.uk](mailto:j.manoharmayum@sheffield.ac.uk)