

Metallocene Sulfide Chemistry of Niobium and Tantalum: New Insights Into the Formation of Niobium–Polysulfido Complexes and Synthesis, Structure, and Reactivity of $\text{Cp}'_2\text{Ta}(\eta^2\text{-S}_2)\text{H}$ ($\text{Cp}' = t\text{-BuC}_5\text{H}_4$)

Hans-Jürgen Bach, Henri Brunner, and Joachim Wachter*

Institut für Anorganische Chemie der Universität Regensburg, D-8400 Regensburg, Germany

Marek M. Kubicki, Jean-Claude Leblanc, Claude Moise, and Florence Volpatto

Laboratoire de Synthèse et d'Electrosynthèse Organométalliques, Université de Bourgogne, F-21100 Dijon, France


Bernd Nuber and Manfred L. Ziegler†

Institut für Anorganische Chemie der Universität, D-6900 Heidelberg, Germany

Received October 30, 1991


Summary: The previously described reaction of $\text{Cp}'_2\text{NbH}_3$ ($\text{Cp}' = t\text{-BuC}_5\text{H}_4$) with S_8 was investigated with regard to the kinetic stabilization of as yet elusive intermediates during the formation of the polysulfide complex $\text{Cp}'_4\text{Nb}_2\text{S}_9$ (1). Therefore, the permethylated complex $\text{Cp}^*_2\text{NbS}_3\text{H}$ (2; $\text{Cp}^* = \text{C}_5\text{Me}_5$) and the Ta complex $\text{Cp}'_2\text{TaS}_2\text{H}$ (3) were synthesized in a related reaction and structurally investigated. Despite the problematic localization of the hydride atom during X-ray crystallographic studies, the molecules can be described in a proper manner: Characteristic of 2 and 3 are $\text{M}(\eta^2\text{-S}_2)$ cores to which either an SH (2) or a H (3) ligand is attached. Complex 3 is the first known tantalocene sulfide. It readily loses sulfur upon reaction with PR_3 ($\text{R} = \text{C}_6\text{H}_5$, OCH_3) and CH_3I to form compounds 4 and 5, which both contain the $\text{Ta}=\text{S}$ moiety. Attempts to prepare "classical" metallocene sulfides by metathesis reaction of $\text{Cp}'_2\text{NbCl}_2$ and Li_2S_5 gave the polynuclear compounds $\text{Cp}'_3\text{Nb}_3\text{S}_{10}\text{O}$ (6) and $\text{Cp}'_4\text{Nb}_4\text{S}_{13}$ (7).

A new entry into niobium polysulfide chemistry has recently been found in the reaction of $\text{Cp}'_2\text{NbH}_3$ with elemental sulfur. Among very similar products $\text{Cp}'_4\text{Nb}_2\text{S}_9$ could be characterized unequivocally (eq 1). In this




complex two $\text{Cp}'_2\text{Nb}(\eta^2\text{-S}_2)$ units are linked by only one sulfur chain.¹ The high content of S-S bonds in 1 and related compounds gives rise to a very complex transformation into polynuclear compounds when their solutions are heated.² The question can also be raised of how such compounds, e.g. 1, are formed. This question gets still more important when we consider that during the reaction expressed by eq 1 no intermediates have been observed. From our investigations in cyclopentadienylmetal sulfides we are aware of the influence which is exerted by the substitution degree of the Cp ligands as well as by the employed metal on the nature of the products.³ Indeed, these observations were confirmed when $\text{Cp}^*_2\text{NbBH}_4$ ($\text{Cp}^* = \text{C}_5\text{Me}_5$), which is more easily accessible than $\text{Cp}'_2\text{NbH}_3$,^{4,5}

was reacted with sulfur (eq 2).¹




The two isomers A and B have been proposed for the mononuclear complex 2; however, an unambiguous structural attribution has not yet been possible. It is one goal



of this paper to elucidate the definite structure of 2 by means of an X-ray diffraction study.

Another goal is to study the consequence of the "metal effect" on the synthesis of group 5 metallocene chalcogenides by using $\text{Cp}'_2\text{TaH}_3$ instead of $\text{Cp}'_2\text{NbH}_3$. The structure of the resulting product as well as the first studies of its reactivity will be reported.

A final goal of this investigation was to find out whether Nb(IV) sulfides are able to play a role in the formation of 1. In this regard it was of interest to test synthetic routes which have been already successfully employed in vanadocene chalcogen chemistry (eq 3).^{6,7} By the method of eq 3 it was attempted to synthesize the NbS_2 unit⁸ directly by metathesis from $\text{Cp}'_2\text{NbCl}_2$ and Li_2S_2 or by thermal decomposition of the eventually formed $\text{Cp}'_2\text{NbS}_5$ to give dinuclear complexes.⁹



(4) Bell, R. A.; Cohen, S. A.; Doherty, N. M.; Threlkel, R. S.; Bercaw, J. E. *Organometallics* 1986, 5, 972.

(5) It has already been proved for the Cp' series that the trihydride reacts in the same manner with S_8 as the BH_4^- complex.¹

(6) Muller, K. G.; Petersen, J. L.; Dahl, L. F. *J. Organomet. Chem.* 1976, 111, 91. Petersen, J. L.; Dahl, L. F. *J. Am. Chem. Soc.* 1975, 97, 6416.

(7) Köpf, H.; Wirl, A.; Kahl, W. *Angew. Chem., Int. Ed. Engl.* 1971, 10, 137.

(8) Mononuclear niobocene complexes bearing the $\text{Nb}(\eta^2\text{-S}_2)$ moiety are as follows. (a) $\text{Cp}_2\text{NbS}_2\text{X}$ ($\text{Cp} = \text{C}_5\text{H}_5$; $\text{X} = \text{Cl}, \text{Br}, \text{I}, \text{SCN}$): Treichel, P. M.; Werber, G. P. *J. Am. Chem. Soc.* 1968, 90, 1753. Roder, R. M. Ph.D. Thesis, University of Wisconsin, 1973; *Chem. Abstr.* 1974, 81, 30727. (b) $\text{Cp}_2\text{NbS}_2[\text{S}_2\text{P}(\text{OR})_2]_2$: Viard, B.; Sala-Pala, J.; Amaudrut, J.; Guerchais, J. E.; Sanchez, C.; Livage, J. *Inorg. Chim. Acta* 1980, 39, 99. (c) $\text{Cp}_2\text{NbS}_2\text{Me}$: Amaudrut, J.; Guerchais, J. E.; Sala-Pala, J. *J. Organomet. Chem.* 1978, 157, C10.

*Deceased in April 1991.

(1) Brunner, H.; Klement, U.; Wachter, J.; Tsunoda, M.; Leblanc, J.-C.; Moise, C. *Inorg. Chem.* 1990, 29, 584.

(2) Brunner, H.; Meier, W.; Wachter, J.; Nuber, B.; Ziegler, M. L. J. *Organomet. Chem.* 1990, 381, C7.

(3) Wachter, J. *Angew. Chem., Int. Ed. Engl.* 1989, 28, 1613.

Experimental Section

All procedures were carried out under nitrogen with N₂-saturated, dry solvents. Elemental analyses were performed by the Mikroanalytisches Laboratorium, Universität Regensburg. IR spectra were obtained with a Beckman 4240 spectrophotometer and a Perkin-Elmer 580 B instrument. Field desorption mass spectra were run on a Finnigan MAT 311A instrument from toluene solutions. NMR spectra were recorded on Bruker WM 250 and 400 instruments. Complex 2 was prepared as described in ref 1 and recrystallized from toluene.

Preparation of Cp'₂TaS₂H (3). A mixture of 1000 mg (2.35 mmol) of (t-BuC₅H₄)₂TaH₃¹⁰ and 370 mg (1.44 mmol) of S₈ in 60 mL of toluene was stirred for 4 h at 25 °C. After chromatographic workup (SiO₂/toluene) 3 was isolated as a yellow band in 46% yield. Recrystallization from toluene/pentane gave bright yellow needles. Anal. Calcd for C₁₈H₂₇TaS₂: C, 44.26; H, 5.57; mol wt, 488.5. Found: C, 44.23; H, 5.66; mol wt, 487.9 (FD-MS). ¹H NMR (C₆D₆): δ 1.23 (s, 18), 2.79 (s, 1), 4.09 (q, 2), 4.38 (q, 2), 4.57 (q, 2), 6.04 (q, 2). IR (KBr): ν_{S-S} 525 cm⁻¹.

Preparation of Cp'₂Ta(S)H (4). To a solution of 700 mg of (t-BuC₅H₄)₂TaS₂H (1.43 mmol) in toluene (60 mL) was added 310 mg of P(OEt)₃ (1.89 mmol). The reaction mixture was refluxed for 4 h, and the solvent was then removed under vacuum. The residual solid was washed with pentane and taken to dryness, giving 425 mg (65% yield) of 4 as a pink solid, which was recrystallized from toluene/pentane. Anal. Calcd for C₁₈H₂₇TaS: C, 47.37; H, 5.96; mol wt, 456.4. Found: C, 46.73; H, 5.71; mol wt, 456.6 (FD-MS). ¹H NMR (C₆D₆): δ 1.34 (s, 18), 4.82 (q, 2), 5.24 (q, 2), 5.32 (q, 2), 5.67 (q, 2), 7.10 (s, 1). IR (CsI): ν_{Ta-H} 1859.0 cm⁻¹, ν_{Ta=S} 433.7 cm⁻¹.

Preparation of Cp'₂Ta(S)I (5). A solution of 270 mg (0.55 mmol) of (t-BuC₅H₄)₂TaS₂H in toluene (40 mL) was stirred at room temperature for 4 h with 230 mg (1.66 mmol) of iodomethane. The resulting brown solution was then evaporated, and the residual solid was chromatographed on silica gel. Toluene eluted compound 5 (34% yield) as a greenish band which yielded bright golden leaves after recrystallization from toluene. Anal. Calcd for C₁₈H₂₆TaSI: C, 37.12; H, 4.50; mol wt, 582.3. Found: C, 37.34; H, 4.22; mol wt, 582.2 (FD-MS). ¹H NMR (C₆D₆): δ 1.34 (s, 18), 4.73 (q, 2), 5.14 (q, 2), 6.08 (q, 2), 6.41 (q, 2). IR (CsI): ν_{Ta-S} 433.7 cm⁻¹.

Reaction of Cp'₂NbCl₂¹¹ with Li₂S₅. To a suspension of 1019 mg (2.47 mmol) of Cp'₂NbCl₂ in 80 mL of toluene was added 5 mL of 0.5 M Li₂S₅ in THF. The mixture was refluxed for 20 h and then concentrated to 10 mL. Chromatography on silanized SiO₂ (column 22 × 3.5 cm) gave a wine red band eluted with toluene, followed by a brown band eluted with THF. The latter consisted of a complex mixture which could not be separated into distinct components. The toluene band was first rechromatographed on SiO₂ (eluent toluene) and then on three connected Merck-Lobar columns (type B (310/25 mm); Lichroprep Si60 (40–63 μm)) in 4/1 petroleum ether (40–60 °C)/ether to give 6 and then 7. Both orange compounds were recrystallized from CH₂Cl₂. Analytical and spectroscopic data for 6 (yield 5.5%) were consistent with those for the completely characterized Cp'₃Nb₃S₁₀O₂.² Anal. Calcd for C₃₈H₅₂Nb₄S₁₃ (7): C, 34.00; H, 4.12; S, 32.76; mol wt, 1273.2. Found: C, 33.96; H, 4.05; S, 32.70; mol wt, 1272.2 (FD-MS). ¹H NMR (CDCl₃): δ 1.22 (s, 9), 1.29 (s, 18), 1.34 (s, 9), 5.26 (m, 2), 5.35 (m, 2), 5.42 (m, 4), 5.72 (m, 2), 5.85 (m, 2), 6.07 (m, 2), 6.20 (m, 2). IR (KBr): ν_{S-S} 544 w, 538 w, 522 s cm⁻¹.

Crystallographic Studies

Cp'₂NbS₃H. A red crystal, obtained from toluene, having the approximate dimensions 0.4 × 0.46 × 0.69 mm was mounted on a Syntex R3 diffractometer. The unit cell was determined and refined from 24 reflections (15 < 2θ < 31°). Relevant crystal and data collection parameters are given in Table I. The structure

Table I. Crystal Data and Data Collection Parameters for (C₅Me₅)₂Nb(S₂)SH (2) and (t-BuC₅H₄)₂Ta(S₂)H (3)

	2	3
mol formula	NbS ₃ C ₂₀ H ₃₁	TaS ₂ C ₁₈ H ₂₇
fw	460.57	488.49
cryst syst	monoclinic	monoclinic
space group	P2 ₁ /c (No. 14)	P2 ₁ /c (No. 14)
cell dimens		
a, Å	8.626 (5)	13.698 (1)
b, Å	14.82 (1)	10.084 (2)
c, Å	17.19 (1)	13.642 (8)
β, °	104.74 (5)	90.85 (2)
V, Å ³	2125.2	1884.2
Z	4	4
T, K	296	296
d _{calc} , g·cm ⁻³	1.44	1.722
linear abs, μ, cm ⁻¹	8.3	59.72
F(000)	960	960
diffractometer	Syntex R3	Enraf-Nonius CAD4
radiation, Å	λ(Mo Kα) = 0.71073	λ(Mo Kα) = 0.71073
scan type	θ-2θ	ω-2θ
rflns measd	h, 0-12; k, 0-21; l -24 to +24	h, 0-16; k, 0-12; l, -16 to +16
2θ range, deg	3-57.5	4-56
no. of rflns measd	6112	3587
abs cor; range of transmissn	empirical, 7 rflns; 0.87-1.00	empirical, DIFABS; 0.87-1.19
no. of unique obsd data	4143, I > 2.5σ(I)	2408, I > 3σ(I)
no. of parms (NV)	218	190
R(F)	0.056	0.043
R _w (F)	0.057	0.047
weighting scheme	w = 1/σ ² F _o	w ⁻¹ = [σ ² (I) + (0.07I) ²] ^{1/2}
GOF	3.97	3.608
ρ, residual, e·Å ⁻³	0.92	2.42

Table II. Atomic Coordinates and Equivalent Isotropic Displacement Parameters (Å²) for Cp'₂NbS₃H (2)

atom	x	y	z	U _{eq} ^a
Nb	0.2581 (1)	0.1954 (1)	0.1649 (1)	41 (1)
S1	0.3141 (3)	0.3197 (1)	0.2676 (1)	105 (1)
S2	-0.0176 (2)	0.1278 (2)	0.1076 (1)	95 (1)
S31	0.0699 (9)	0.2712 (6)	0.2264 (4)	119 (4)
S32	0.0113 (7)	0.2325 (5)	0.1990 (4)	94 (3)
C1	0.2537 (6)	0.0504 (4)	0.2420 (4)	59 (2)
C2	0.3419 (9)	0.1116 (4)	0.2957 (3)	67 (2)
C3	0.4801 (7)	0.1347 (4)	0.2717 (4)	66 (2)
C4	0.4788 (6)	0.0831 (3)	0.2031 (3)	53 (2)
C5	0.3360 (6)	0.0356 (3)	0.1815 (3)	49 (2)
C6	0.1124 (8)	-0.0009 (5)	0.2540 (5)	126 (4)
C7	0.3035 (13)	0.1365 (6)	0.3739 (4)	147 (6)
C8	0.6244 (10)	0.1824 (5)	0.3202 (5)	141 (5)
C9	0.6285 (8)	0.0649 (5)	0.1759 (5)	110 (4)
C10	0.2951 (9)	-0.0360 (4)	0.1187 (4)	97 (3)
C11	0.1514 (7)	0.2867 (6)	0.0404 (4)	82 (3)
C12	0.2202 (10)	0.2083 (4)	0.0181 (3)	74 (3)
C13	0.3790 (9)	0.2116 (4)	0.0483 (4)	66 (3)
C14	0.4185 (8)	0.2857 (5)	0.0925 (4)	73 (3)
C15	0.2815 (10)	0.3362 (4)	0.0873 (4)	74 (3)
C16	-0.0138 (10)	0.3205 (8)	0.0106 (6)	211 (7)
C17	0.1347 (15)	0.1443 (7)	-0.0443 (4)	226 (8)
C18	0.4899 (13)	0.1547 (7)	0.0138 (6)	199 (8)
C19	0.5864 (10)	0.3204 (6)	0.1286 (6)	169 (6)
C20	0.2663 (16)	0.4314 (5)	0.1094 (5)	231 (10)

^aEquivalent isotropic U defined as one-third of the trace of the orthogonalized U_{ij} tensor.

was solved by using standard Patterson methods, least-squares refinement, and Fourier techniques. All non-hydrogen atoms were refined anisotropically; the hydrogen atoms were included riding on calculated positions (HFIX program of SHELXTL¹²). All calculations were performed with the SHELXTL program;¹² scattering

(9) Bolinger, C. M.; Rauchfuss, T. B.; Rheingold, A. L. *Organometallics* 1982, 1, 1551.

(10) Green, M. L. H.; Joussemae, B. *J. Organomet. Chem.* 1980, 193, 339.

(11) Tebbe, F. N.; Parshall, G. W. *J. Am. Chem. Soc.* 1971, 93, 3793. Labinger, J. A. *Adv. Chem. Ser.* 1978, No. 167, 149.

(12) Sheldrick, G. M. SHELXTL Program, Universität Göttingen, Göttingen, Germany, 1983.




Figure 1. ORTEP plot of **2** (50% probability level). Two views are given, each of which contains the central S atom in a 50:50 statistical distribution. The H atom bound to S1 or S2 (see text) cannot be localized exactly and is therefore omitted.

factors were taken from ref 13.

Cp'₂TaS₂H. A yellow irregularly shaped crystal having the approximate dimensions $0.40 \times 0.16 \times 0.10$ mm was mounted on an Enraf-Nonius CAD4 diffractometer. The unit cell was determined and refined from 25 randomly selected reflections (CAD4 routines). Relevant crystal and data collection parameters are given in Table I. A preliminary search of diffraction maxima and systematic extinctions indicated the unique space group $P2_1/c$. All calculations were carried out by using the Enraf-Nonius SDP library¹⁴ with neutral-atom scattering factors. Intensities of the 3587 reflections measured were corrected for Lorentz and polarization effects and for linear decay; 2408 unique ones with $I > 3\sigma(I)$ were used in the structure solution and refinement. The structure was solved by conventional Patterson and difference Fourier methods, and no particular difficulty due to decomposition of the crystal was observed. The empirical absorption correction DIFABS¹⁵ was applied after isotropic refinement of the structure. In the final full-matrix least-squares refinements all non-hydrogen atoms were given anisotropic temperature factors and all hydrogen atoms of *t*-BuC₅H₄ ligands were placed in calculated positions (HYDRO program of SDP), riding on the carbon atoms bearing them, and included with B_{iso} varying from 5.0 to 13.0 Å². A final difference Fourier map revealed the three strongest peaks at 2.3, 2.2, and 2.1 e/Å³. The first peak was 2.11 and 1.21 Å away from the Ta and S2 atoms, respectively, and located in the TaS₂ plane. The attempt to refine it as a hydride was unsuccessful, because its positional parameters varied by more than 2 Å from one cycle to another. Thus, we considered this peak as an artifact due to high electron density in the TaS₂ plane. Other difference Fourier peaks are close to the tantalum atom (0.7–0.9 Å). Final atomic coordinates of non-hydrogen atoms are given in Table III.

Results and Discussion

Complex **2** was formed from $(C_5Me_5)_2NbBH_4$ and S_8 in moderate yields (eq 2).¹ Two isomeric forms were proposed for **2**, of which **2A** (involving a puckered niobathiacycle and a metal hydride) seemed to be more likely by means of IR and 1H NMR data.¹ We have now succeeded in obtaining single crystals from toluene, and thus an X-ray diffraction analysis was carried out.

Unfortunately, the result of this study (Table II, Figure 1) has been hampered by a disorder of the central sulfur atom S3 and the fact that the H atom could not be

Table III. Atomic Coordinates and Equivalent Isotropic Displacement Parameters (\AA^2) for $(t\text{-BuC}_5\text{H}_4)_2\text{TaS}_2\text{H}$ (3)

atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{eq} ^a
Ta	0.18050 (3)	0.14229 (4)	0.69670 (3)	34 (3)
S1	0.0999 (3)	0.1097 (4)	0.8585 (2)	75 (3)
S2	0.2445 (4)	0.0293 (4)	0.8374 (3)	94 (2)
C1	0.3028 (8)	0.313 (1)	0.6671 (8)	42 (6)
C2	0.2695 (9)	0.330 (1)	0.7650 (8)	48 (7)
C3	0.1703 (9)	0.366 (1)	0.7598 (8)	52 (6)
C4	0.1420 (8)	0.368 (1)	0.6604 (9)	51 (6)
C5	0.2212 (8)	0.335 (1)	0.6030 (8)	45 (6)
C6	0.4078 (9)	0.301 (1)	0.635 (1)	59 (8)
C7	0.445 (1)	0.444 (2)	0.624 (2)	111 (15)
C8	0.416 (1)	0.233 (2)	0.535 (1)	100 (10)
C9	0.471 (1)	0.229 (2)	0.711 (1)	114 (14)
C11	0.1589 (8)	-0.069 (1)	0.6035 (8)	43 (6)
C12	0.162 (1)	0.039 (1)	0.5389 (8)	56 (7)
C13	0.0814 (9)	0.120 (1)	0.5494 (9)	60 (7)
C14	0.0289 (8)	0.067 (1)	0.6290 (8)	48 (6)
C15	0.0769 (8)	-0.048 (1)	0.6615 (8)	47 (6)
C16	0.219 (1)	-0.197 (1)	0.600 (1)	67 (8)
C17	0.214 (1)	-0.274 (1)	0.697 (1)	112 (14)
C18	0.172 (1)	-0.280 (2)	0.518 (1)	104 (15)
C19	0.324 (1)	-0.171 (2)	0.582 (2)	145 (20)

^aEquivalent isotropic U defined as one-third of the trace of the orthogonalized U_{ii} tensor.

Table IV. Selected Distances and Angles for 2

Distances (Å)			
Nb-S1	2.511 (2)	Nb-C(1-5) _{av}	2.488 (6)
Nb-S2	2.538 (2)	Nb-C(11-15) _{av}	2.493 (8)
Nb-S31	2.427 (9)	S1-S31	2.167 (8)
Nb-S32	2.411 (7)	S2-S32	2.178 (8)
Angles (deg)			
S1-Nb-S2	123.8 (1)	S1-Nb-S32	71.7 (2)
S1-Nb-S31	52.0 (2)		

localized. Nevertheless, the planar NbS_3 core of the molecule now found allows a clear attribution of 2 to type B, which is characterized by the still rare assembly of a S_2^{2-} and a SH^- ligand at one metal center. The corresponding ν_{SH} absorption is obscured in the IR spectrum.¹⁶ Probably as a consequence of the disorder of the inner S atom the S-S distances (2.17 (1) Å) are rather long,¹⁷ but not too long when compared to S-S ligands in other co-ordination modes.¹⁸ The other Nb-S and all Nb-C bond

(13) *International Tables for X-Ray Crystallography*; Kynoch Press: Birmingham, U.K., 1974; Vol. 4.

(14) Frenz, B. A. The Enraf-Nonius CAD4-SDP. A Real Time System for Concurrent X-Ray Data Collection and Crystal Structure Determination. In *Computing in Crystallography*; Schenk, M., Olthof-Hazekamp, R., Van Koningsveld, M., Eds.; Delft University Press: Delft, Holland, 1978.

(15) Walker, N.; Stuart, D. *Acta Crystallogr., Sect. A* 1983, A39, 158.

(16) It is generally accepted that ν_{SH} absorptions are often very weak. See e.g.: Kwon, D.; Real, J.; Curtis, M. D.; Rheingold, A.; Haggarty, B. S. *Organometallics* 1991, 10, 143.

(17) Müller, A.; Diemann, E. *Adv. Inorg. Chem.* 1987, 31, 89.


Table V. Selected Distances and Angles for 3

	Distances (Å)	
Ta-S1	2.504 (4)	Ta-C(11-15) _{av}
Ta-S2	2.388 (5)	S1-S2
Ta-C(1-5) _{av}	2.42 (1)	
	Angles (deg)	
S1-Ta-S2	52.5 (2)	M _{Cp1} -Ta-M _{Cp2} ^a
		131.6 (2)

^a M_{Cp1} and M_{Cp2} are the midpoints of the Cp rings C1-C5 and C11-C15, respectively.

lengths (Table IV) do agree well with those already found in polynuclear cyclopentadienylniobium sulfides.^{1,2} Thus, 2 is the first niobocene complex bearing the NbS₂ unit which has allowed an exact structure determination.⁸

The reaction of Cp'₂TaH₃ with excess sulfur gives at ambient temperature the yellow complex 3 as the only product (the liberated gas consists of at least an important portion of H₂S). Reaction times longer than 4 h and elevated temperatures give rise to the formation of as yet uncharacterized polynuclear compounds. The new complex is readily soluble in common organic solvents; its solutions are air-sensitive. For the composition Cp'₂TaS₂H, which follows from analytical data, one may imagine the three isomeric forms A-C. As only one resonance for the



tert-butyl substituent is observed in the ¹H NMR spectrum, one may rule out isomer A, which would require two different environments for the Cp ligand. In addition, the resonances of the *t*-BuCp protons are close to the values observed in other Ta(V) complexes reported in this paper, whereas they are rather far from the chemical shifts reported for Ta(III) complexes such as Cp'₂Ta(CO)H.¹⁹

From spectroscopic data it is not possible to distinguish clearly between isomers B and C. Single crystals of 3 were obtained from toluene/pentane, and an X-ray structure analysis was carried out (Table III). Although an overall weakening of the diffraction pattern was observed during data collection and although the H atom could not be localized, the structure solution represents a sufficiently broad basis for the discussion of the structure. The core of the molecule is characterized by a TaS₂ triangle (Figure 2, Table V) with an angle of 52.5 (2)^o at Ta. This triangle is oriented in such a way that S2 lies only 0.17 Å outside the plane defined by the gravity centers of both Cp rings and Ta compared to -1.85 Å for S1. Such a geometry is only possible for isomer C, whereas in the case of B a π-bonding orbital would be localized in the plane bisecting the Cp rings with the corresponding S atoms in lateral positions. The formulation of a S₂ ligand is also consistent with the presence of a ν_{S-S} absorption at 525 cm⁻¹ in the IR spectrum.

It is striking that, as in 2, an expansion of the S-S bond (2.165 (7) Å) is observed. Therefore, one may try to explain this stretching, at least in part, by the strong acceptor properties of the S₂ ligand. The structural arrangement is further supported by the deshielded ¹H NMR resonance signal of the metal hydride at δ 2.79. A similarly low field shift has already been observed for complexes of the type





Figure 2. ORTEP plot of 3 (50% probability level). The H atom bound to Ta cannot be localized exactly and is therefore omitted.

Cp'₂Ta(*n*²-L)H (L = N₂CPh₂, CH(CN)NR),²⁰ which are isosteric with 3. Of speculative character is the assumption of a weak interaction of the metal hydride with the inner S atom of the S₂ ligand in the solid state as a quasi-intermediate between both isomeric forms A and C. It is not yet clear if such a behavior would explain the presence of two unequal Ta-S bonds and the missing ν_{M-H} absorption in the IR spectrum.

In contrast to the case for 2, complex 3 is available in quantities sufficient to allow us to study its chemical reactivity. Preliminary results which are in good agreement with structural type 3C are summarized in eq 4. If so-



lutions of 3 are treated with equimolar amounts of phosphites or phosphines, one S atom is abstracted to give the pink compound 4. An IR absorption at 1895 cm⁻¹ is indicative of a Ta-H moiety, whereas the only strong absorption below 600 cm⁻¹ is tentatively assigned as a ν_{Ta-S} absorption (433 cm⁻¹). The ν_{S-S} absorption of the starting material has disappeared. In the ¹H NMR spectrum only small changes are observed for the resonances of the ring protons when compared with those of 3. The metal hydride resonance of the latter now appears at δ +7.10, a fact which can only be explained by the presence of a Ta(V) center.²¹ A similar chalcogen abstraction with phosphines has recently been described for the structurally related complex Cp'₂Ta(*n*²-O₂)CH₃.²²

In another reaction 3 reacts with CH₃I to give complex 5, in which the ν_{Ta-H} and ν_{S-S} absorptions of 3 have disappeared. A new strong absorption at 433 cm⁻¹ is attributed to a ν_{Ta-S} frequency. Although the ¹H NMR spectrum is of less diagnostic value, it must be stated that all four quadruplets of the ring protons are shifted to lower

(20) Leboeuf, J. F.; Leblanc, J. C.; Moise, C. *J. Organomet. Chem.* 1989, 364, C22.

(21) Belmonte, P.; Schrock, R. R.; Churchill, M. R.; Youngs, W. J. *J. Am. Chem. Soc.* 1980, 102, 2858.


(22) von Assett, A.; Trimmer, M. S.; Henling, L. M.; Bercaw, J. E. *J. Am. Chem. Soc.* 1988, 110, 8254.

(18) Brunner, H.; Pfaunach, J.; Wachter, J.; Nuber, B.; Ziegler, M. L. *J. Organomet. Chem.* 1989, 359, 179.

(19) Reynoud, J. F.; Leblanc, J. C.; Moise, C. *Organometallics* 1985, 4, 1059.

field. As the first step in this reaction we assume that there is likely an attack of CH_3I at the nucleophilic S_2 ligand followed by the elimination of CH_3SH .

All compounds described in this paper exist formally in the oxidation state +5. The fact that in the related vanadium chalcogen chemistry oxidation state +4 is dominating^{6,7,9,23} prompted us to try the synthesis of Nb(IV) complexes due to eq 1. However, when $\text{Cp}'_2\text{NbCl}_2$ was reacted with Li_2S_2 or Li_2S_5 , the formation of the expected metallathiacycles could not be observed. Although their formation cannot generally be excluded, the reaction took place at such high temperatures (above 100 °C) that finally only multinuclear complexes were isolated in a poorly selective reaction in 6% (6) and 8% yield (7). The structure of 6, which is also one of the products of the thermal decomposition of 1, has already been solved. It is characterized by a set of five different S_2 ligands and a rather flat Nb_3O tetrahedron.² The structure of 7 may



be derived from a butterfly-like Nb_4 core spanned by a $\mu_4\text{-S}$ ligand; a Ta complex exhibiting such a structure is known.²⁴ Then the Nb centers would be coordinated by mono- and disulfide ligands in such a manner that the molecule contains a symmetry plane. This would be in agreement with the ^1H NMR spectrum (three CH_3 resonances for the *t*-Bu groups in the ratio 1:2:1).


Conclusion

The starting point for this investigation was the one-step synthesis of the polysulfide complex 1, which should proceed via distinct intermediates when regarding its rather complex structure. Now, the isolation and characterization of 2 and 3 allows a more detailed insight into

(23) Gambarotta, S.; Floriani, C.; Chiesi-Villa, A.; Guastini, C. *J. Chem. Soc., Chem. Commun.* 1983, 184.

(24) Fenske, D.; Mané, P. G. *Z. Naturforsch.* 1989, B44, 531.

the reaction of group 5 metallocene hydrides with sulfur, for the variation of the Cp substituents as well as the metal affords a kinetic stabilization of as yet elusive intermediate products. Thus, we propose a general reaction sequence in which the stability of the products depends on the π -ligands and the metal centers (eq 5). Disulfide complexes,



as represented by 3, still bearing a reactive metal hydride obviously are the first relatively stable intermediates after the attack of the nucleophilic metal center at the S_8 ring. Then one may imagine the insertion of sulfur into the M-H bond, leading in the case of 2 to an isolable product. As the final step oxidative coupling of two SH ligands of two different molecules by sulfur may lead to the bridging pentasulfide ligand being sufficiently stable in 1.²⁵ In this context we must cite one of the first observations in niobocene sulfide chemistry: $\text{Cp}_2\text{Nb}(\text{S}_2)\text{Cl}$ was found to react with H_2S to give a brown product of presumably oligomeric nature of composition $(\text{Cp}_2\text{NbS}_4)_n$.^{8a}

All these examples demonstrate that the metallocene sulfide chemistry of Nb and Ta is much more complicated than originally assumed.⁸ It also differs considerably from related group 5 chemistry²⁶ in that stable metallocene complexes are only formed in the +5 oxidation state and that no dinuclear compounds with a direct metal-metal interaction exist. Further investigations will be addressed to use 3 as a building block in the synthesis of polynuclear complexes and to employ the potential reactivity of its inorganic ligands in organic synthesis.

Registry No. 2, 125303-74-4; 3, 138783-57-0; 4, 138783-58-1; 5, 138783-59-2; 6, 127649-42-7; 7, 127649-43-8; (*t*-BuC₅H₄)₂TaH₃, 84749-48-4; Cp'₂NbCl₂, 61374-53-6; Li₂S₅, 66559-78-2.

Supplementary Material Available: Tables of bond lengths, bond angles, anisotropic displacement parameters, and H atom coordinates for 2 and 3 (8 pages); tables of observed and calculated structure factors (32 pages). Ordering information is given on any current masthead page.

(25) Further examples of related oxidative couplings of SH ligands are given in: Amarasekera, J.; Rauchfuss, T. B. *Inorg. Chem.* 1989, 28, 3875.

(26) Draganjac, M.; Rauchfuss, T. B. *Angew. Chem., Int. Ed. Engl.* 1985, 24, 742. Herberhold, M.; Kuhnlein, M.; Schrepfermann, M.; Ziegler, M. L.; Nuber, B. *J. Organomet. Chem.* 1990, 398, 259.