



## Epstein-Barr virus (EBV)-host cell interactions: an epigenetic dialogue?

Hans Helmut Niller, Kalman Szenthe and Janos Minarovits

Journal Name:	Frontiers in Genetics
ISSN:	1664-8021
Article type:	Mini Review Article
Received on:	13 Aug 2014
Accepted on:	02 Oct 2014
Provisional PDF published on:	02 Oct 2014
www.frontiersin.org:	<a href="http://www.frontiersin.org">www.frontiersin.org</a>
Citation:	Niller HH, Szenthe K and Minarovits J(2014) Epstein-Barr virus (EBV)-host cell interactions: an epigenetic dialogue?. <i>Front. Genet.</i> 5:367. doi:10.3389/fgene.2014.00367
Copyright statement:	© 2014 Niller, Szenthe and Minarovits. This is an open-access article distributed under the terms of the <a href="http://creativecommons.org/licenses/by/2.0/">Creative Commons Attribution License (CC BY)</a> . The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after rigorous peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

## **Epstein-Barr virus (EBV)-host cell interactions: an epigenetic dialogue?**

Hans Helmut Niller<sup>1\*</sup>, Kalman Szenthe<sup>2</sup>, and Janos Minarovits<sup>3</sup>

<sup>1</sup>Institute of Medical Microbiology and Hygiene, University of Regensburg, Franz-Josef-Strauss Allee 11, D-93053 Regensburg, Germany

<sup>2</sup>RT-Europe Nonprofit Research Ltd., Pozsonyi u. 88., H-9200 Mosonmagyaróvár, Hungary

<sup>3</sup>Department of Oral Biology and Experimental Dental Research, University of Szeged, Faculty of Dentistry, H-6720 Szeged, Tisza Lajos krt. 64. Hungary

\*Corresponding author

e-mail: Hans-Helmut.Niller@klinik.uni-regensburg.de

**Key words:** chromatin, CpG island, DNA methylation, viral latency, lytic cycle, methylome, oncoprotein, pioneer transcription factor, tumour suppressor gene, tumour virus

## **Abstract**

Here, we wish to highlight the genetic exchange and epigenetic interactions between EBV and its host. Epstein-Barr virus (EBV) is associated with diverse lymphoid and epithelial malignancies. Their molecular pathogenesis is accompanied by epigenetic alterations which are distinct for each of them. While lymphoblastoid cell lines (LCLs) derived from B cells transformed by EBV *in vitro* are characterized by a massive demethylation and euchromatinization of the viral and cellular genomes, the primarily malignant lymphoid tumour Burkitt's lymphoma (BL) and the epithelial tumours nasopharyngeal carcinoma (NPC) and EBV-associated gastric carcinoma (EBVaGC) are characterized by hypermethylation of a multitude of cellular tumour suppressor gene loci and of the viral genomes. In some cases, the viral latency and oncoproteins including the latent membrane proteins LMP1 and LMP2A and several nuclear antigens (EBNAs) affect the level of cellular DNA methyltransferases or interact with the histone modifying machinery. Specific molecular mechanisms of the epigenetic dialogue between virus and host cell remain to be elucidated.

### **Epstein-Barr virus – the first human tumour virus**

Epstein-Barr virus (EBV), the proto-typical gamma-herpesvirus infecting humans, has been discovered 50 years ago in cultured Burkitt's lymphoma (BL) cells (Epstein et al., 1964). EBV physiologically homes to memory B cells, a property which is reflected in the genus name *Lymphocryptovirus* for gamma-herpesviruses which "hide in lymphoid cells". EBV is the causative agent of infectious mononucleosis, and was the first known human tumour virus. In addition to BL, an endemic childhood tumour mainly of equatorial Africa, EBV plays a role in the origin or progression of other primarily malignant B cell tumours, such as Hodgkin lymphoma, and diverse AIDS-associated lymphomas including lymphomas of the central nervous system. A profound failure of T cell surveillance may allow early-onset post-transplant lymphoproliferative disorder (PTLD) which may be overcome upon a timely onset of T cell control, but may turn malignant if growing uncontrolled for too long a time span.

In the first decade of EBV research, EBV was considered a purely lymphotropic virus. Finding viral DNA in cellular DNA from biopsies of anaplastic carcinomas of the nasopharynx (NPC) by DNA hybridization did not change that general view, because NPC as a lymphoepithelial tumour contains a great many of infiltrating lymphocytes (zur Hausen et al., 1970). Localizing the virus specifically to the malignant epithelial cells first established EBV infection of non-lymphatic cells and paved the way for the novel concept of EBV as an epithelial tumour virus (Wolf et al., 1973). NPC is an endemic tumour with a strong preference for South East Asia, especially Guangdong and Hong Kong, with an incidence rate of 20 to 30 cases per 100.000 persons per year, and virtually 100% of nonkeratinizing and undifferentiated NPCs are EBV-associated. Almost two decades later on, the association of EBV with gastric carcinomas (EBVaGC) was established, too (Burke et al., 1990; Shibata et al., 1991). Contrary to the endemic tumour NPC, about 10% of the worldwide sporadic tumour gastric carcinoma (GC) are associated with EBV infection. Among gastric remnant carcinomas approximately 30% and among lymphoepithelioma-like GCs, approximately 80% are EBV-associated. Altogether, EBVaGC with an

estimated more than 80.000 cases per year is probably the most frequent EBV-associated malignancy worldwide (reviewed in Niller et al., 2014a;Niller et al., 2014b). Beyond the B cell lineage, EBV infection is currently associated with T cell lymphomas, epithelial tumours, and rarely with leiomyosarcoma, a neoplasm of mesodermal origin (McClain et al., 1995). Furthermore, the risk of autoimmune disease, including multiple sclerosis, is significantly increased after primary EBV-infection, and even more so after symptomatic primary infection, i.e. mononucleosis (Niller et al., 2008).

### **Pathogenesis of EBV-associated tumours**

The spectacular ability of EBV to transform and immortalize B cells dominated the first four decades of EBV research and tumour virology. General view was that the EBV-transformed cell was the origin of the endemic BL cell, too, although a fundamental difference of the epidemiology and pathogenesis between LCL-like tumours on one side (early onset PTLT), and of primarily malignant EBV-associated lymphomas on the other side (endemic BL) became evident. Early onset PTLT originate under conditions of severe immune suppression and depend on viral transforming functions, including Epstein-Barr virus nuclear antigens (EBNAs) and latent membrane proteins (LMPs) that are expressed both in PTLTs *in vivo* and in LCLs immortalized *in vitro*. On the other hand, BL and Hodgkin lymphoma originate under conditions of hyperstimulation of the lymphoid germinal center reaction, and they do not depend on EBNA2 which the vast majority of them do not express (Klein, 1987;Lenoir and Bornkamm, 1987), (Table 1). In this context, it is important to distinguish between morphological and oncogenic transformation (Niller et al., 2011). Our discovery of a binding site for the oncoprotein c-Myc in the central locus control region of the EBV genome suggested that the molecular pathogenesis of endemic BL does not depend on a previous EBNA2-transformed state of the B-cell, but mostly on a dysbalance of pro-and anti-apoptotic functions in consequence of myc-translocation, a molecular accident in a virus infected B cell undergoing the germinal center reaction (Niller et al., 2003). The

need to counter-balance the pro-apoptotic force of translocated *c-Myc* through anti-apoptotic functions, either encoded by the viral genome or induced by virus infection, in order for a BL to emerge has recently been re-emphasized (Mbulaiteye, 2013; Westhoff Smith and Sugden, 2013). Our differential pathogenesis model for EBV-associated lymphomas was controversial at first (Rossi and Bonetti, 2004; Thorley-Lawson, 2004; reviewed in Niller et al., 2012), but has now gained strong support by recent large-scale epigenomic analyses of LCLs and tumour cells (see below). Thus, finding the binding site for the oncoprotein *c-Myc* in the locus control region of EBV caused a conceptual shift away from the morphologically transformed cell and has turned out as a heuristic discovery (Niller et al., 2003).

### **Genetic exchange between herpesviruses and host cells**

Homology between herpesviral and human genes is now a common theme which was first highlighted by the finding of a gene for a functional thymidylate synthase (*TS*) in the genome of herpesvirus saimiri with an extremely high homology of 70% identical amino acids with the human *TS* gene. Various parameters suggested that the *TS* gene had been acquired in virus evolution by an ancestral herpesvirus from the cellular genome (Bodemer et al., 1986; Honess et al., 1986). The exchange of human and viral genes, and in the case of human herpesvirus 6 (HHV-6) the invasion of an entire herpesviral genome into the human germ line (Daibata et al., 1999) in about 0.8% of humans, must have happened on numerous occasions in evolutionary time. In the case of EBV, the intimate evolutionary relationship of virus and host cell is emphasized by the presence of several viral genes with sequence homology to cellular genes, i.e. *BHRF1* and *BALF1*, two anti-apoptotic *BCL2* homologues, *BILF1*, coding for a constitutively active G protein-coupled receptor (GPCR) homologue, and *BCRF1*, an *IL-10* (*interleukin 10*) gene homologue. The *BCRF1* protein appears to be a functional homologue of *IL-10*, an immune suppressive cytokine secreted by Th2 cells with a sequence identity of about 70% (Hsu et al., 1990). The sequence homologies of the other three viral peptides are of lower degree than that. *BHRF1* carries a 25% identity of an 150 amino acid C-terminal portion with

the anti-apoptotic cellular protein BCL2 (Cleary et al., 1986). BALF1 shows homology at a similar degree in functionally important domains to its cellular counterparts BCL2 and BCLX (Marshall et al., 1999). BILF1 carries a homology of around 20% with the human chemokine receptor CXCR3A (Davis-Poynter and Farrell, 1996;Beisser et al., 2005;Paulsen et al., 2005). EBV-associated carcinoma cells express BARF1, identified initially as a lytic cycle protein that binds to hCSF-1 (human colony stimulating factor 1) as a viral decoy receptor, although its crystal structure is most closely related to CD80, a co-stimulatory molecule of antigen presenting cells (Seto et al., 2005;Tarbouriech et al., 2006;Elegheert et al., 2012).

Herpesviral DNA binding proteins ICP8 of HSV1 and BALF2 of EBV, and additional homologous proteins of human herpesviruses which are required for viral replication belong to a class of “DDE/RNase H-like fold-family” nucleases, together with the recombination activating gene (RAG) 1 protein, essential for V(D)J recombination, and the Argonaute protein of the RNA induced silencing complex (RISC). For ICP8 of HSV, divalent cation binding of the DDE-site was actually shown to be functional and required for viral replication (Bryant et al., 2012). Furthermore, inhibitors of HIV integrase, another RNase H-fold protein, inhibited the replication of viruses from all herpesvirus genera, too (Yan et al., 2014). Based on a co-regulatory transcriptional network for both *RAG-1* and *RAG-2*, and the genes for herpesviral DNA-binding proteins, and based on signature sequence homologies between V(D)J recombination sites and the viral terminal repeats, an evolutionary relationship between the RAG recombinase and herpesviral DNA binding proteins was proposed. The *RAG* locus may have originally been introduced to host cells by a primordial herpesvirus (Dreyfus, 2009). We found a striking co-linearity of structural and functional elements between the cellular immunoglobulin gene loci and the left part of the EBV genome. Therefore, although speculative, we agree with the view that, in the case of the *RAG* genes, the appearance of the adaptive immune system may have been dependent on a primordial herpesvirus genome and, in the case of the B cell, may have developed further as consequence of an evolutionary ping-pong game between EBV and the host cell (Niller et al., 2004).

### **Epigenetic interactions between EBV and its host cells**

EBV infects both B lymphocytes and epithelial cells *in vivo*. It enters B cells after binding to CD21, a cell surface molecule absent from epithelial cells. However, EBV infected B cells are capable to transfer the virus to epithelial cells lacking the EBV receptor. Both B cells and epithelial cells can support productive (lytic) EBV replication when all of the proteins and nontranslated RNAs encoded by the viral genome are expressed in a sequential order. EBV also causes latent infections, typically in resting memory B cells and in various neoplastic cells that usually carry circular, double stranded viral genomes. During latency, only a restricted set of EBV promoters is active. The activity of latent EBV promoters depends on the phenotype of host cells, and it is controlled by the epigenetic regulatory machinery. Based on the epigenetic marks deposited on the viral chromatin by the cellular epigenetic machinery one can distinguish between viral epigenotypes that are associated with unique patterns of viral gene expression (reviewed in Minarovits, 2006). In parallel, certain latent EBV proteins characteristic for the major cell types carrying latent EBV genomes act as epigenetic regulators themselves: they alter the cellular epigenotype and gene expression pattern and may contribute to the development of malignant tumours. Thus, the situation is similar to an „epigenetic dialogue”, indeed. Typical examples of EBV latency types are summarized in Table 1, based on the nomenclature suggested by Klein et al., 2013 (Klein et al., 2013), see also (Laytragoon-Lewin et al., 1995;Niller et al., 2012).

### **Epigenetic control of EBV latency promoters**

The epigenetic regulatory mechanisms of host cells not only control the preservation of cell type specific gene expression patterns from cell generation to cell generation, but ensure the maintenance of host cell-dependent usage of latent EBV promoters as well. Epigenetic regulation is based on writing, reading and erasing epigenetic marks on chromatin as well as on protein-DNA interactions that are stable even in mitotic, highly condensed chromatin (reviewed in Gopalakrishnan et al.,

2008;Zaret et al., 2008;Sharma et al., 2010;Blomen and Boonstra, 2011). Euchromatic marks favour transcription whereas heterochromatic marks are associated with a more condensed chromatin structure that usually represses promoter activity.

DNA methylation is involved in silencing of most latent EBV promoters. It is well documented that the alternative promoters Wp and Cp, where transcripts coding for six Epstein-Barr virus nuclear antigens (EBNAs) are initiated can be switched off by CpG methylation. In addition, LMP1p and LMP2Ap, the promoters for latent membrane protein 1 and 2A transcripts are silenced by the activity of cellular DNA methyltransferases, too. DNA methylation does not play a role, however, in switching off Qp, an alternative promoter for EBNA1 transcripts (reviewed in Li and Minarovits, 2003;Niller et al., 2012). One may speculate, that EBNA2, the major transactivator protein encoded by the EBV genome that interacts with both histone acetyltransferases and histone deacetylases, (reviewed in Niller et al., 2009) may activate key cellular genes mediating genome-wide demethylation in LCLs. Acetylation of histone H3 and histone H4 molecules and di- or trimethylation of lysine 4 of histone H3 (H3K4me2 or H3K4me3) are euchromatic marks frequently associated with active Cp, Qp, LMP1p and LMP2Ap (reviewed in Niller et al., 2012;Arvey et al., 2013). In principle, complexes formed by Polycomb and Trithorax group proteins that modify histone tails could also play a role in the regulation of latent EBV promoters but there are no data supporting such a mechanism. In contrast, Polycomb group protein EZH2 was observed to leave a heterochromatic histone mark (H3k27me3) at early lytic promoters of EBV that are silent during latency, and chromatin immunoprecipitation proved the association of the EZH2 methyltransferase with this class of viral promoters in the BL cell line Raji (Woellmer et al., 2012). The immediate early promoter Zp, where transcripts for the lytic cycle initiating BZLF1 protein are initiated was also found to be repressed by H3k27me3 but also by H4K20me3 in Raji cells (Murata et al., 2012).

Pioneer transcription factors and variant histone molecules that bind to repressive chromatin areas and mark the genes to be activated have not been implicated in the control of EBV latency. Chromatin

loops formed by binding of insulator proteins to EBV episomes may play a role, however, in the epigenetic regulation of the latency promoters Cp and Qp (Tempera et al., 2011).

### **EBV-encoded oncoproteins as epigenetic regulators**

The methylation patterns of nasopharyngeal carcinoma cells and EBVaGC cells differ from their normal counterparts: these neoplastic cells regularly carry hypermethylated genomic regions with silenced cellular promoters (Iizasa et al., 2012; Lo et al., 2012). Focal hypermethylations frequently inactivate tumour suppressor genes and may contribute to carcinogenesis and tumour progression. Because the EBV encoded transmembrane proteins LMP1 and LMP2A are capable of upregulating the cellular DNA methyltransferases DNMT1, DNMT3A and DNMT3B, it was suggested that hypermethylation of CpG rich sequences, the so called CpG islands, is mediated by LMP1 or LMP2A in EBV-associated carcinomas (reviewed in Niller et al., 2012). LMP1 and LMP2A are expressed in EBV positive Hodgkin lymphomas as well. Thus, they may contribute to gene silencing in these neoplasms, too.

It is worthy to note that, similarly to the EBV-associated carcinomas, LCLs established by *in vitro* EBV-infection of B cells also express LMP1 and LMP2A that could potentially upregulate DNA methyltransferases. It was observed, however, that the typical epigenetic change in LCLs is a widespread demethylation of the B cell methylome affecting one third of all cellular genes and 2.18 GB of the genome (Hansen et al., 2014). The mechanism of demethylation remains to be established. Furthermore, the EBV episomes carried by LCLs are also hypomethylated, in contrast to the overall hypermethylation of EBV genomes in latency type I BL lines, BL biopsies and EBV-associated carcinomas (Minarovits et al., 1991; Fernandez et al., 2009). One may speculate, that EBNA2, the major transactivator protein encoded by the EBV genome that interacts with both histone acetyltransferases and histone deacetylases (reviewed in Niller et al., 2009) may activate key cellular genes mediating genome-wide demethylation of both cellular and viral DNA in LCLs. The viral oncoproteins EBNA3A and EBNA3C expressed during *in vitro* immortalization of B cells silence distinct

cellular tumour suppressor genes by depositing a heterochromatic histone mark via the Polycomb repressor complex PRC2 (reviewed in Allday, 2013). The dominant change in EBV immortalized B cells seems to be, however, a genome-wide decrease and redistribution of heterochromatic marks (Hernando et al., 2014). The EBV latency products eliciting the reprogramming of the host cell epigenome in LCLs remain to be elucidated. Contrary to LCLs, primarily malignant lymphomas, i.e. BL and other lymphomas, are characterized not by a massive and wide-spread hypomethylation, but by a local hypermethylation of selected genomic loci (Martin-Subero et al., 2009a; Martin-Subero et al., 2009b; Kreck et al., 2013; reviewed in Niller et al., 2014b).

In contrast to BL tumours, in which hypermethylated loci are strongly enriched for polycomb repressive complex (PRC) target genes of embryonic stem cells, hypermethylated genes in EBVaGC are not enriched for PRC targets (Martin-Subero et al., 2009a; Matsusaka et al., 2011). Contrary to *Helicobacter pylori*-associated GC, EBVaGC does not emerge from an “epigenetic field” in the gastric mucosa (Matsusaka et al., 2011; Niller et al., 2014a). Thus, EBV-associated epigenetic changes may quite quickly set the stage for malignancy (Au et al., 2005; Niller et al., 2014a). Notably, even transient EBV infection of epithelial cells leaves permanent epigenetic scars indicating past infection (Queen et al., 2013; Birdwell et al., 2014).

Although there are no data as to the interaction of host cell-encoded pioneer transcription factors and the EBV genome, certain viral proteins that bind to both viral and cellular DNA and remain associated with mitotic chromosomes may act as pioneer transcription factors. EBNA1, a nuclear protein expressed in all EBV latency types is a putative pioneer factor functionally resembling transcription factors of the FoxA family that act as epigenetic regulators controlling important developmental processes (Niller and Minarovits, 2012). EBNA1 was shown to bind to different sets of cellular promoters in cell lines of epithelial and lymphoid origin, and upregulation as well as downregulation of distinct gene batteries was observed (Canaan et al., 2009). In the BL cell line Raji, EBNA1 was shown to interact with a large number of cellular genes as well as LINE elements, and high

affinity EBNA binding sites were observed in a repetitive sequence in chromosome 11 (Lu et al., 2010).

BZLF1, an immediate-early EBV protein initiating productive viral replication that preferentially binds methylated DNA sequences was also suggested to act as a pioneer transcription factor (Woellmer et al., 2012). BZLF1 is expressed not only during the lytic cycle but transiently also during the establishment of latent infection of B cells *in vitro* (Kalla and Hammerschmidt, 2012). Because BZLF1 can bind to cellular promoters (Lan et al., 2013) and elicit epigenetic alterations (Woellmer et al., 2012), an “epigenetic dialogue” between the latent EBV genomes and the host cell genome may occur, indeed: transient BZLF1 expression may change the cellular epigenotype followed by silencing of the BZLF1 promoter through the cellular epigenetic machinery. In parallel, EBV encoded epigenetic regulators may leave their marks on the cellular epigenotype in a next phase of EBV-mediated B cell immortalization.

In conclusion, on an evolutionary time scale a genetic exchange between herpesviruses and their hosts is evident. Beginning with the early steps of viral infection, epigenetic interactions between virus and host cell are taking place. The multi-tiered epigenetic dialogue between EBV and its host needs to be elucidated in greater molecular detail in order to understand the diverse outcomes of infection.

**Table 1. Host cell-dependent expression of latent Epstein-Barr virus proteins.** Key viral latency proteins or so-called oncoproteins are differentially expressed in EBV-associated malignancies and transformed cell lines. Depending on the specific viral gene expression patterns, specific viral latency classes are defined.

Protein	Cell type and latency type			
	BL	NPC	CLL	LCL
	Type I	Type IIa	Type IIb	Type III
EBNA1	+	+	+	+
EBNA2	-	-	+	+
LMP1	-	+/-	-	+
LMP2A	-	+	+	+
BARF1	-	+	-	-

#### Abbreviations

BL: Burkitt's lymphoma, NPC: nasopharyngeal carcinoma, CLL: chronic lymphocytik leukaemia, LCL: lymphoblastoid cell line, EBNA: Epstein-Barr virus nuclear antigen, LMP: latent membrane protein, BARF1: *Bam*HI A rightward frame 1.

## References

- Allday, M.J. (2013). EBV finds a polycomb-mediated, epigenetic solution to the problem of oncogenic stress responses triggered by infection. *Front Genet* 4:212.
- Arvey, A., Tempera, I., and Lieberman, P.M. (2013). Interpreting the Epstein-Barr Virus (EBV) epigenome using high-throughput data. *Viruses* 5, 1042-1054.
- Au, W.Y., Pang, A., Chan, E.C., Chu, K.M., Shek, T.W., and Kwong, Y.L. (2005). Epstein-Barr virus-related gastric adenocarcinoma: an early secondary cancer post hemopoietic stem cell transplantation. *Gastroenterology* 129, 2058-2063.
- Beisser, P.S., Verzijl, D., Gruijthuijsen, Y.K., Beuken, E., Smit, M.J., Leurs, R., Bruggeman, C.A., and Vink, C. (2005). The Epstein-Barr virus BILF1 gene encodes a G protein-coupled receptor that inhibits phosphorylation of RNA-dependent protein kinase. *J Virol* 79, 441-449.
- Birdwell, C.E., Queen, K.J., Kilgore, P., Rollyson, P., Trutschl, M., Cvek, U., and Scott, R.S. (2014). Genome-wide DNA methylation as an epigenetic consequence of Epstein-Barr virus infection of immortalized keratinocytes. *J Virol*, pii: JVI.00972-14.
- Blomen, V.A., and Boonstra, J. (2011). Stable transmission of reversible modifications: maintenance of epigenetic information through the cell cycle. *Cell Mol Life Sci* 68, 27-44.
- Bodemer, W., Niller, H.H., Nitsche, N., Scholz, B., and Fleckenstein, B. (1986). Organization of the thymidylate synthase gene of herpesvirus saimiri. *J Virol* 60, 114-123.
- Bryant, K.F., Yan, Z., Dreyfus, D.H., and Knipe, D.M. (2012). Identification of a divalent metal cation binding site in herpes simplex virus 1 (HSV-1) ICP8 required for HSV replication. *J Virol* 86, 6825-6834.
- Burke, A.P., Yen, T.S., Shekitka, K.M., and Sobin, L.H. (1990). Lymphoepithelial carcinoma of the stomach with Epstein-Barr virus demonstrated by polymerase chain reaction. *Mod Pathol* 3, 377-380.
- Canaan, A., Haviv, I., Urban, A.E., Schulz, V.P., Hartman, S., Zhang, Z., Palejev, D., Deisseroth, A.B., Lacy, J., Snyder, M., Gerstein, M., and Weissman, S.M. (2009). EBNA1 regulates cellular gene expression by binding cellular promoters. *Proc Natl Acad Sci U S A* 106, 22421-22426.
- Cleary, M.L., Smith, S.D., and Sklar, J. (1986). Cloning and structural analysis of cDNAs for bcl-2 and a hybrid bcl-2/immunoglobulin transcript resulting from the t(14;18) translocation. *Cell* 47, 19-28.
- Daibata, M., Taguchi, T., Nemoto, Y., Taguchi, H., and Miyoshi, I. (1999). Inheritance of chromosomally integrated human herpesvirus 6 DNA. *Blood* 94, 1545-1549.
- Davis-Poynter, N.J., and Farrell, H.E. (1996). Masters of deception: a review of herpesvirus immune evasion strategies. *Immunol Cell Biol* 74, 513-522.
- Dreyfus, D.H. (2009). Paleo-immunology: evidence consistent with insertion of a primordial herpes virus-like element in the origins of acquired immunity. *PLoS One* 4, e5778.
- Elegheert, J., Bracke, N., Pouliot, P., Gutsche, I., Shkumatov, A.V., Tarbouriech, N., Verstraete, K., Bekaert, A., Burmeister, W.P., Svergun, D.I., Lambrecht, B.N., Vergauwen, B., and Savvides, S.N. (2012). Allosteric competitive inactivation of hematopoietic CSF-1 signaling by the viral decoy receptor BARF1. *Nat Struct Mol Biol* 19, 938-947.
- Epstein, M.A., Achong, B.G., and Barr, Y.M. (1964). Virus particles in cultured lymphoblasts from Burkitt's Lymphoma. *Lancet* 1, 702-703.
- Fernandez, A.F., Rosales, C., Lopez-Nieva, P., Grana, O., Ballestar, E., Roperio, S., Espada, J., Melo, S.A., Lujambio, A., Fraga, M.F., Pino, I., Javierre, B., Carmona, F.J., Acquadro, F., Steenbergen, R.D., Snijders, P.J., Meijer, C.J., Pineau, P., Dejean, A., Lloveras, B., Capella, G., Quer, J., Buti, M., Esteban, J.I., Allende, H., Rodriguez-Frias, F., Castellsague, X., Minarovits, J., Ponce, J., Capello, D., Gaidano, G., Cigudosa, J.C., Gomez-Lopez, G., Pisano, D.G., Valencia, A., Piris, M.A., Bosch, F.X., Cahir-Mcfarland, E., Kieff, E., and Esteller, M. (2009). The dynamic DNA methylomes of double-stranded DNA viruses associated with human cancer. *Genome Res* 19, 438-451.
- Gopalakrishnan, S., Van Emburgh, B.O., and Robertson, K.D. (2008). DNA methylation in development and human disease. *Mutat Res* 647, 30-38.

- Hansen, K.D., Sabunciyan, S., Langmead, B., Nagy, N., Curley, R., Klein, G., Klein, E., Salamon, D., and Feinberg, A.P. (2014). Large-scale hypomethylated blocks associated with Epstein-Barr virus-induced B-cell immortalization. *Genome Res* 24, 177-184.
- Hernando, H., Islam, A.B., Rodriguez-Ubreva, J., Forne, I., Ciudad, L., Imhof, A., Shannon-Lowe, C., and Ballestar, E. (2014). Epstein-Barr virus-mediated transformation of B cells induces global chromatin changes independent to the acquisition of proliferation. *Nucleic Acids Res* 42, 249-263.
- Honess, R.W., Bodemer, W., Cameron, K.R., Niller, H.H., Fleckenstein, B., and Randall, R.E. (1986). The A+T-rich genome of Herpesvirus saimiri contains a highly conserved gene for thymidylate synthase. *Proc Natl Acad Sci U S A* 83, 3604-3608.
- Hsu, D.H., De Waal Malefyt, R., Fiorentino, D.F., Dang, M.N., Vieira, P., De Vries, J., Spits, H., Mosmann, T.R., and Moore, K.W. (1990). Expression of interleukin-10 activity by Epstein-Barr virus protein BCRF1. *Science* 250, 830-832.
- Iizasa, H., Nanbo, A., Nishikawa, J., Jinushi, M., and Yoshiyama, H. (2012). Epstein-Barr Virus (EBV)-associated gastric carcinoma. *Viruses* 4, 3420-3439.
- Kalla, M., and Hammerschmidt, W. (2012). Human B cells on their route to latent infection--early but transient expression of lytic genes of Epstein-Barr virus. *Eur J Cell Biol* 91, 65-69.
- Klein, E., Nagy, N., and Rasul, A.E. (2013). EBV genome carrying B lymphocytes that express the nuclear protein EBNA-2 but not LMP-1: Type IIb latency. *Oncoimmunology* 2, e23035.
- Klein, G. (1987). "In defense of the "old" Burkitt Lymphoma scenario," in *Advances in Viral Oncology*, ed. G. Klein. (New York: Raven Press), 207-211.
- Kreck, B., Richter, J., Ammerpohl, O., Barann, M., Esser, D., Petersen, B.S., Vater, I., Murga Penas, E.M., Bormann Chung, C.A., Seisenberger, S., Boyd, V.L., Smallwood, S., Drexler, H.G., Macleod, R.A., Hummel, M., Krueger, F., Hasler, R., Schreiber, S., Rosenstiel, P., Franke, A., and Siebert, R. (2013). Base-pair resolution DNA methylome of the EBV-positive Endemic Burkitt lymphoma cell line DAUDI determined by SOLiD bisulfite-sequencing. *Leukemia* 27, 1751-1753.
- Lan, Y.Y., Yeh, T.H., Lin, W.H., Wu, S.Y., Lai, H.C., Chang, F.H., Takada, K., and Chang, Y. (2013). Epstein-Barr virus Zta upregulates matrix metalloproteinases 3 and 9 that synergistically promote cell invasion in vitro. *PLoS One* 8, e56121.
- Laytragoon-Lewin, N., Chen, F., Avila-Carino, J., Zou, J.Z., Mellstedt, H., Ernberg, I., and Klein, G. (1995). Epstein Barr virus (EBV)-carrying cells of a chronic lymphocytic leukemia (CLL) subpopulation express EBNA1 and LMPs but not EBNA2 in vivo. *Int J Cancer* 63, 486-490.
- Lenoir, G.M., and Bornkamm, G.W. (1987). "Burkitt's Lymphoma, a human cancer model for the study of the multistep development of cancer: proposal for a new scenario," in *Advances in Viral Oncology*, ed. G. Klein. (New York: Raven Press), 173-206.
- Li, H., and Minarovits, J. (2003). Host cell-dependent expression of latent Epstein-Barr virus genomes: regulation by DNA methylation. *Adv Cancer Res* 89, 133-156.
- Lo, K.W., Chung, G.T., and To, K.F. (2012). Deciphering the molecular genetic basis of NPC through molecular, cytogenetic, and epigenetic approaches. *Semin Cancer Biol* 22, 79-86.
- Lu, F., Wikramasinghe, P., Norseen, J., Tsai, K., Wang, P., Showe, L., Davuluri, R.V., and Lieberman, P.M. (2010). Genome-wide analysis of host-chromosome binding sites for Epstein-Barr Virus Nuclear Antigen 1 (EBNA1). *Virol J* 7:262.
- Marshall, W.L., Yim, C., Gustafson, E., Graf, T., Sage, D.R., Hanify, K., Williams, L., Fingerroth, J., and Finberg, R.W. (1999). Epstein-Barr virus encodes a novel homolog of the bcl-2 oncogene that inhibits apoptosis and associates with Bax and Bak. *J Virol* 73, 5181-5185.
- Martin-Subero, J.I., Ammerpohl, O., Bibikova, M., Wickham-Garcia, E., Agirre, X., Alvarez, S., Bruggemann, M., Bug, S., Calasanz, M.J., Deckert, M., Dreyling, M., Du, M.Q., Durig, J., Dyer, M.J., Fan, J.B., Gesk, S., Hansmann, M.L., Harder, L., Hartmann, S., Klapper, W., Kuppers, R., Montesinos-Rongen, M., Nagel, I., Pott, C., Richter, J., Roman-Gomez, J., Seifert, M., Stein, H., Suela, J., Trumper, L., Vater, I., Prosper, F., Haferlach, C., Cruz Cigudosa, J., and Siebert, R. (2009a). A comprehensive microarray-based DNA methylation study of 367 hematological neoplasms. *PLoS One* 4, e6986.

- Martin-Subero, J.I., Kreuz, M., Bibikova, M., Bentink, S., Ammerpohl, O., Wickham-Garcia, E., Rosolowski, M., Richter, J., Lopez-Serra, L., Ballestar, E., Berger, H., Agirre, X., Bernd, H.W., Calvanese, V., Cogliatti, S.B., Drexler, H.G., Fan, J.B., Fraga, M.F., Hansmann, M.L., Hummel, M., Klapper, W., Korn, B., Kuppers, R., Macleod, R.A., Moller, P., Ott, G., Pott, C., Prosper, F., Rosenwald, A., Schwaenen, C., Schubeler, D., Seifert, M., Sturzenhofecker, B., Weber, M., Wessendorf, S., Loeffler, M., Trumper, L., Stein, H., Spang, R., Esteller, M., Barker, D., Hasenclever, D., and Siebert, R. (2009b). New insights into the biology and origin of mature aggressive B-cell lymphomas by combined epigenomic, genomic, and transcriptional profiling. *Blood* 113, 2488-2497.
- Matsusaka, K., Kaneda, A., Nagae, G., Ushiku, T., Kikuchi, Y., Hino, R., Uozaki, H., Seto, Y., Takada, K., Aburatani, H., and Fukayama, M. (2011). Classification of Epstein-Barr virus-positive gastric cancers by definition of DNA methylation epigenotypes. *Cancer Res* 71, 7187-7197.
- Mbulaiteye, S.M. (2013). Burkitt Lymphoma: beyond discoveries. *Infect Agent Cancer* 8, 35.
- McClain, K.L., Leach, C.T., Jenson, H.B., Joshi, V.V., Pollock, B.H., Parmley, R.T., Dicarlo, F.J., Chadwick, E.G., and Murphy, S.B. (1995). Association of Epstein-Barr virus with leiomyosarcomas in children with AIDS. *N Engl J Med* 332, 12-18.
- Minarovits, J. (2006). Epigenotypes of latent herpesvirus genomes. *Curr Top Microbiol Immunol* 310, 61-80.
- Minarovits, J., Minarovits-Kormuta, S., Ehlin-Henriksson, B., Falk, K., Klein, G., and Ernberg, I. (1991). Host cell phenotype-dependent methylation patterns of Epstein-Barr virus DNA. *J Gen Virol* 72, 1591-1599.
- Murata, T., Kondo, Y., Sugimoto, A., Kawashima, D., Saito, S., Isomura, H., Kanda, T., and Tsurumi, T. (2012). Epigenetic histone modification of Epstein-Barr virus BZLF1 promoter during latency and reactivation in Raji cells. *J Virol* 86, 4752-4761.
- Niller, H.H., Banati, F., Ay, E., and Minarovits, J. (2012). "Epigenetic changes in virus-associated neoplasms," in *Patho-Epigenetics of Disease*, eds. J. Minarovits & H.H. Niller. (New York: Springer), 179-225.
- Niller, H.H., Banati, F., and Minarovits, J. (2014a). Epigenetic alterations in nasopharyngeal carcinoma and Epstein-Barr virus (EBV) associated gastric carcinoma: a lesson in contrasts. *J Nasopharyng Carcinoma* 1, e9.
- Niller, H.H., and Minarovits, J. (2012). Similarities between the Epstein-Barr Virus (EBV) Nuclear Protein EBNA1 and the Pioneer Transcription Factor FoxA: Is EBNA1 a "Bookmarking" Oncoprotein that Alters the Host Cell Epigenotype? *Pathogens* 1, 37-51.
- Niller, H.H., Salamon, D., Ilg, K., Koroknai, A., Banati, F., Bauml, G., Rucker, O., Schwarzmann, F., Wolf, H., and Minarovits, J. (2003). The in vivo binding site for oncoprotein c-Myc in the promoter for Epstein-Barr virus (EBV) encoding RNA (EBER) 1 suggests a specific role for EBV in lymphomagenesis. *Med Sci Monit* 9, HY1-HY9.
- Niller, H.H., Salamon, D., Rahmann, S., Ilg, K., Koroknai, A., Banati, F., Schwarzmann, F., Wolf, H., and Minarovits, J. (2004). A 30 kb region of the Epstein-Barr virus genome is colinear with the rearranged human immunoglobulin gene loci: implications for a "ping-pong evolution" model for persisting viruses and their hosts. A review. *Acta Microbiol Immunol Hung* 51, 469-484.
- Niller, H.H., Tarnai, Z., Decsi, G., Zsedenyi, A., Szenthe, K., Banati, F., and Minarovits, J. (2014b). The role of epigenetics in Epstein-Barr virus regulation and pathogenesis. *Future Microbiol* 9, 747-746.
- Niller, H.H., Wolf, H., and Minarovits, J. (2008). Regulation and dysregulation of Epstein-Barr virus latency: implications for the development of autoimmune diseases. *Autoimmunity* 41, 298-328.
- Niller, H.H., Wolf, H., and Minarovits, J. (2009). Epigenetic dysregulation of the host cell genome in Epstein-Barr virus-associated neoplasia. *Semin Cancer Biol* 19, 158-164.
- Niller, H.H., Wolf, H., and Minarovits, J. (2011). Viral hit and run-oncogenesis: Genetic and epigenetic scenarios. *Cancer Lett* 305, 200-217.

- Paulsen, S.J., Rosenkilde, M.M., Eugen-Olsen, J., and Kledal, T.N. (2005). Epstein-Barr virus-encoded BILF1 is a constitutively active G protein-coupled receptor. *J Virol* 79, 536-546.
- Queen, K.J., Shi, M., Zhang, F., Cvek, U., and Scott, R.S. (2013). Epstein-Barr virus-induced epigenetic alterations following transient infection. *Int J Cancer* 132, 2076-2086.
- Rossi, G., and Bonetti, F. (2004). EBV and Burkitt's lymphoma. *N Engl J Med* 350, 2621.
- Seto, E., Yang, L., Middeldorp, J., Sheen, T.S., Chen, J.Y., Fukayama, M., Eizuru, Y., Ooka, T., and Takada, K. (2005). Epstein-Barr virus (EBV)-encoded BARP1 gene is expressed in nasopharyngeal carcinoma and EBV-associated gastric carcinoma tissues in the absence of lytic gene expression. *J Med Virol* 76, 82-88.
- Sharma, S., Kelly, T.K., and Jones, P.A. (2010). Epigenetics in cancer. *Carcinogenesis* 31, 27-36.
- Shibata, D., Tokunaga, M., Uemura, Y., Sato, E., Tanaka, S., and Weiss, L.M. (1991). Association of Epstein-Barr virus with undifferentiated gastric carcinomas with intense lymphoid infiltration. Lymphoepithelioma-like carcinoma. *Am J Pathol* 139, 469-474.
- Tarbouriech, N., Ruggiero, F., De Turenne-Tessier, M., Ooka, T., and Burmeister, W.P. (2006). Structure of the Epstein-Barr virus oncogene BARP1. *J Mol Biol* 359, 667-678.
- Tempera, I., Klichinsky, M., and Lieberman, P.M. (2011). EBV latency types adopt alternative chromatin conformations. *PLoS Pathog* 7, e1002180.
- Thorley-Lawson, D.A. (2004). EBV and Burkitt's lymphoma. *N Engl J Med* 350, 2621.
- Westhoff Smith, D., and Sugden, B. (2013). Potential cellular functions of Epstein-Barr Nuclear Antigen 1 (EBNA1) of Epstein-Barr Virus. *Viruses* 5, 226-240.
- Woellmer, A., Arteaga-Salas, J.M., and Hammerschmidt, W. (2012). BZLF1 governs CpG-methylated chromatin of Epstein-Barr Virus reversing epigenetic repression. *PLoS Pathog* 8, e1002902.
- Wolf, H., Zur Hausen, H., and Becker, V. (1973). EB viral genomes in epithelial nasopharyngeal carcinoma cells. *Nat New Biol* 244, 245-247.
- Yan, Z., Bryant, K.F., Gregory, S.M., Angelova, M., Dreyfus, D.H., Zhao, X.Z., Coen, D.M., Burke, T.R., Jr., and Knipe, D.M. (2014). HIV Integrase Inhibitors Block Replication of Alpha-, Beta-, and Gammaherpesviruses. *MBio* 5, pii: e01318-14.
- Zaret, K.S., Watts, J., Xu, J., Wandzioch, E., Smale, S.T., and Sekiya, T. (2008). Pioneer factors, genetic competence, and inductive signaling: programming liver and pancreas progenitors from the endoderm. *Cold Spring Harb Symp Quant Biol* 73, 119-126.
- Zur Hausen, H., Schulte-Holthausen, H., Klein, G., Henle, W., Henle, G., Clifford, P., and Santesson, L. (1970). EBV DNA in biopsies of Burkitt tumours and anaplastic carcinomas of the nasopharynx. *Nature* 228, 1056-1058.