



Two-loop evolution equations for light-ray operators



V.M. Braun^{a,*}, A.N. Manashov^{a,b}

^a Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany

^b Department of Theoretical Physics, St.-Petersburg University, 199034, St.-Petersburg, Russia

ARTICLE INFO

Article history:

Received 4 April 2014

Accepted 7 May 2014

Available online 20 May 2014

Editor: A. Ringwald

ABSTRACT

QCD in non-integer $d = 4 - 2\epsilon$ space-time dimensions possesses a nontrivial critical point and enjoys *exact* scale and conformal invariance. This symmetry imposes nontrivial restrictions on the form of the renormalization group equations for composite operators in physical (integer) dimensions and allows to reconstruct full kernels from their eigenvalues (anomalous dimensions). We use this technique to derive two-loop evolution equations for flavor-nonsinglet quark–antiquark light-ray operators that encode the scale dependence of generalized hadron parton distributions and light-cone distribution amplitudes in the most compact form.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

1. Studies of hard exclusive reactions contribute significantly to the research program at all major existing and planned accelerator facilities. The relevant nonperturbative input in such processes involves operator matrix elements between states with different momenta, dubbed generalized parton distributions (GPDs), or vacuum-to-hadron matrix elements related to light-front hadron wave functions at small transverse separations, the distribution amplitudes (DAs). Scale dependence of these distributions is governed by the renormalization group (RG) equations for the corresponding (nonlocal) operators and has to be calculated to a sufficiently high order in perturbation theory in order to make the QCD description of exclusive reactions fully quantitative. At present, the evolution equations for GPDs (and DAs) are known to the two-loop accuracy [1–3], one order less compared to the corresponding “inclusive” distributions that involve forward matrix elements [4,5] and closing this gap is desirable. The direct calculation is very challenging, and also finding a suitable representation for the results may become a problem as the two-loop evolution equations for GPDs are already very cumbersome.

It has been known for some time [6] that conformal symmetry of the QCD Lagrangian allows one to restore full evolution kernels at given order of perturbation theory from the spectrum of anomalous dimensions at the same order, and the calculation of the special conformal anomaly at one order less. This result was used to calculate the complete two-loop mixing matrix for twist-two operators in QCD [7–9], and derive the two-loop evolution kernels in momentum space for the GPDs [1–3]. In Ref. [10] we have suggested an alternative technique, the difference being

that instead of studying conformal symmetry *breaking* in the physical theory [7–9] we make use of the *exact* conformal symmetry of a modified theory – QCD in $d = 4 - 2\epsilon$ dimensions at critical coupling. Exact conformal symmetry allows one to use algebraic group-theory methods to resolve the constraints on the operator mixing and also suggests the optimal representation for the results in terms of light-ray operators. In this way a delicate procedure of the restoration of the evolution kernels from the results for local operators is completely avoided. We expect that these features will become increasingly advantageous in higher orders.

This modified approach was illustrated in [10] on several examples to the two- and three-loop accuracy for scalar theories. Application to gauge theories, in particular QCD, involves several subtleties that are considered in this work. The main new result are the two-loop evolution equations for flavor-nonsinglet quark–antiquark light-ray operators that encode the scale dependence of generalized hadron parton distributions and light-cone distribution amplitudes in the most compact form.

2. Before going over to technical details, let us first describe the general structure of the approach and the results on a more qualitative level. In order to make use of the (approximate) conformal symmetry of QCD it is natural to use a coordinate-space representation in which the symmetry transformations have a simple form [11]. The relevant objects are light-ray operators that can be understood as generating functions for the renormalized leading-twist local operators:

$$\begin{aligned}
 [\mathcal{O}](x; z_1, z_2) &\equiv [\bar{q}(x + z_1 n) \not{n} q(x + z_2 n)] \\
 &\equiv \sum_{m,k} \frac{z_1^m z_2^k}{m!k!} [(D_n^m \bar{q})(x) \not{n} (D_n^k q)(x)].
 \end{aligned}
 \tag{1}$$

* Corresponding author.

Here $q(x)$ is a quark field, the Wilson line is implied between the quark fields on the light-cone, $D_n = n_\mu D^\mu$ is a covariant derivative, n^μ is an auxiliary light-like vector, $n^2 = 0$, that ensures symmetrization and subtraction of traces of local operators. The square brackets [...] stand for the renormalization using dimensional regularization and MS subtraction. We will tacitly assume that the quark and antiquark have different flavor so that there is no mixing with gluon operators. In most situations the overall coordinate x^μ is irrelevant and can be put to zero; we will often abbreviate $\mathcal{O}(z_1, z_2) \equiv \mathcal{O}(0; z_1, z_2)$.

Light-ray operators satisfy a renormalization-group equation of the form [12]

$$(M\partial_M + \beta(g)\partial_g + \mathbb{H})[\mathcal{O}(z_1, z_2)] = 0, \quad (2)$$

where \mathbb{H} is an integral operator acting on the light-cone coordinates of the fields. It can be written as

$$\mathbb{H}[\mathcal{O}](z_1, z_2) = \int_0^1 d\alpha \int_0^1 d\beta h(\alpha, \beta) [\mathcal{O}](z_{12}^\alpha, z_{21}^\beta), \quad (3)$$

where

$$z_{12}^\alpha \equiv z_1 \bar{\alpha} + z_2 \alpha, \quad \bar{\alpha} = 1 - \alpha, \quad (4)$$

and $h(\alpha, \beta)$ is a certain weight function (kernel).

One can show, see e.g. [10], that the powers $[\mathcal{O}](z_1, z_2) \mapsto (z_1 - z_2)^N$ are eigenfunctions of the evolution operator \mathbb{H} , and the corresponding eigenvalues

$$\gamma_N = \int d\alpha d\beta h(\alpha, \beta) (1 - \alpha - \beta)^{N-1} \quad (5)$$

are nothing else as the anomalous dimensions of local operators of spin N (with $N - 1$ derivatives).

In general the function $h(\alpha, \beta)$ is a function of two variables and therefore the knowledge of the anomalous dimensions γ_N is not sufficient to fix it. However, if the theory is conformally invariant then to the one loop accuracy \mathbb{H} must commute with the generators of the $SL(2)$ transformations $[\mathbb{H}, S_\alpha^{(0)}] = 0$, where

$$\begin{aligned} S_+^{(0)} &= z_1^2 \partial_{z_1} + z_2^2 \partial_{z_2} + 2(z_1 + z_2), \\ S_0^{(0)} &= z_1 \partial_{z_1} + z_2 \partial_{z_2} + 2, \quad S_-^{(0)} = -\partial_{z_1} - \partial_{z_2}. \end{aligned} \quad (6)$$

In this case it can be shown that the function $h(\alpha, \beta)$ (up to trivial terms $\sim \delta(\alpha)\delta(\beta)$ that correspond to the unit operator) takes the form [13]

$$h(\alpha, \beta) = \bar{h}(\tau), \quad \tau = \frac{\alpha\beta}{\bar{\alpha}\bar{\beta}} \quad (7)$$

and is effectively a function of one variable τ called the conformal ratio. This function can easily be reconstructed from its moments (5), alias from the anomalous dimensions.

Conformal symmetry of QCD is broken by quantum corrections which implies that the symmetry of the evolution equations is lost at the two-loop level. In other words, writing the evolution kernel as an expansion in the coupling constant

$$\begin{aligned} \mathbb{H} &= a_s \mathbb{H}^{(1)} + a_s^2 \mathbb{H}^{(2)} + \dots \\ \mapsto \quad h(\alpha, \beta) &= a_s h^{(1)}(\alpha, \beta) + a_s^2 h^{(2)}(\alpha, \beta) + \dots, \end{aligned} \quad (8)$$

where $a_s = \alpha_s/(4\pi)$, we expect that $h^{(1)}(\alpha, \beta)$ only depends on the conformal ratio whereas higher-order contributions remain to be nontrivial functions of two variables α and β .

This prediction is confirmed by the explicit calculation [12]:

$$\begin{aligned} \mathbb{H}^{(1)} f(z_1, z_2) &= 4C_F \left\{ \int_0^1 d\alpha \frac{\bar{\alpha}}{\alpha} [2f(z_1, z_2) - f(z_{12}^\alpha, z_2) - f(z_1, z_{21}^\beta)] \right. \\ &\quad \left. - \int_0^1 d\alpha \int_0^{\bar{\alpha}} d\beta f(z_{12}^\alpha, z_{21}^\beta) + \frac{1}{2} f(z_1, z_2) \right\}. \end{aligned} \quad (9)$$

The corresponding one-loop kernel $h^{(1)}(\alpha, \beta)$ can be written in the following, remarkably simple form [13]

$$h^{(1)}(\alpha, \beta) = -4C_F \left[\delta_+(\tau) + \theta(1 - \tau) - \frac{1}{2} \delta(\alpha)\delta(\beta) \right], \quad (10)$$

where the regularized δ -function, $\delta_+(\tau)$, is defined as

$$\begin{aligned} \int d\alpha d\beta \delta_+(\tau) f(z_{12}^\alpha, z_{21}^\beta) &= \int_0^1 d\alpha \int_0^1 d\beta \delta(\tau) [f(z_{12}^\alpha, z_{21}^\beta) - f(z_1, z_2)] \\ &= - \int_0^1 d\alpha \frac{\bar{\alpha}}{\alpha} [2f(z_1, z_2) - f(z_{12}^\alpha, z_2) - f(z_1, z_{21}^\alpha)]. \end{aligned} \quad (11)$$

Taking appropriate matrix elements and making a Fourier transformation to the momentum fraction space one can check that the expression in Eq. (10) reproduces all classical leading-order (LO) QCD evolution equations: DGLAP equation for parton distributions, ERBL equation for the meson light-cone DAs, and the general evolution equation for GPDs.

The two-loop kernel $h^{(2)}(\alpha, \beta)$ contains contributions of two color structures and a term proportional to the QCD beta function,

$$\begin{aligned} h^{(2)}(\alpha, \beta) &= 8C_F^2 h_1^{(2)}(\alpha, \beta) + 4C_F C_A h_2^{(2)}(\alpha, \beta) \\ &\quad + 4b_0 C_F h_3^{(2)}(\alpha, \beta). \end{aligned} \quad (12)$$

Let us explain how it can be calculated. The idea of Ref. [10] is to consider a modified theory, QCD in non-integer $d = 4 - 2\epsilon$ dimensions. In this theory the β -function has the form

$$\begin{aligned} \beta(a) &= M\partial_M a = 2a(-\epsilon - b_0 a + \mathcal{O}(a^2)), \\ b_0 &= \frac{11}{3} N_c - \frac{2}{3} n_f, \end{aligned} \quad (13)$$

and for a large number of flavors n_f there exists a critical coupling $a_s^* = \alpha_s^*/(4\pi) \sim \epsilon$ such that $\beta(a_s^*) = 0$. The theory thus enjoys exact scale invariance [14,15] and one can argue (see below) that full conformal invariance is also present.¹ As a consequence, the renormalization group equations are exactly conformally invariant: the evolution kernels commute with the generators of the conformal group. The generators are, however, modified by quantum corrections as compared to their canonical expressions (6):

$$S_\alpha = S_\alpha^{(0)} + a_s^* \Delta S_\alpha^{(1)} + (a_s^*)^2 \Delta S_\alpha^{(2)} + \dots \quad (14)$$

¹ Formally the gauge-fixed QCD Lagrangian contains two charges, the coupling and the gauge parameter. The corresponding β -function, $\beta_\xi = M\partial_M \xi$, vanishes in the Landau gauge, $\xi = 0$, so that all Green functions are scale-invariant at the critical point in this gauge; β_ξ also drops out of the RG equations for the correlation functions of gauge-invariant operators.

One can show that

$$\begin{aligned} S_- &= S_-^{(0)}, \\ S_0 &= S_0^{(0)} - \epsilon + \frac{1}{2} \mathbb{H}(a_s^*), \quad \mathbb{H}(a_s^*) = a_s^* \mathbb{H}^{(1)} + \dots \\ S_+ &= S_+^{(0)} + (z_1 + z_2) \left(-\epsilon + \frac{1}{2} a_s^* \mathbb{H}^{(1)} \right) + a_s^* (z_1 - z_2) \Delta_+ \\ &\quad + \mathcal{O}(\epsilon^2), \end{aligned} \quad (15)$$

where

$$\begin{aligned} \Delta_+[\mathcal{O}](z_1, z_2) \\ = -2C_F \int_0^1 d\alpha \left(\frac{\bar{\alpha}}{\alpha} + \ln \alpha \right) [\mathcal{O}(z_{12}^\alpha, z_2) - \mathcal{O}(z_1, z_{21}^\alpha)] \end{aligned} \quad (16)$$

i.e. the generator S_- is not modified, the deformation of S_0 can be calculated exactly in terms of the evolution operator (to all orders in perturbation theory) [10], whereas the deformation of S_+ is nontrivial and has to be calculated explicitly order by order, to the required accuracy. The one-loop expression shown in (15), (16) is derived below, it is a new result. From the pure technical point of view, this calculation replaces evaluation of the conformal anomaly in the theory with broken symmetry in integer dimensions via the conformal Ward identities (CWI) in the approach due to D. Müller [6].

Conformal symmetry of the modified QCD at the critical coupling implies that the generators (15) satisfy the usual $SL(2)$ commutation relations

$$[S_0, S_\pm] = \pm S_\pm, \quad [S_+, S_-] = 2S_0. \quad (17)$$

Expanding them in powers of the coupling a_s^* one obtains a nested set of commutator relations [10]

$$\begin{aligned} [S_+^{(0)}, \mathbb{H}^{(1)}] &= 0, \\ [S_+^{(0)}, \mathbb{H}^{(2)}] &= [\mathbb{H}^{(1)}, \Delta S_+^{(1)}], \\ [S_+^{(0)}, \mathbb{H}^{(3)}] &= [\mathbb{H}^{(1)}, \Delta S_+^{(2)}] + [\mathbb{H}^{(2)}, \Delta S_+^{(1)}], \end{aligned} \quad (18)$$

etc. Note that the commutator of the canonical generator $S_+^{(0)}$ with the evolution kernel at order k on the l.h.s. of each equation is given in terms of the evolution kernels $\mathbb{H}^{(k)}$ and the corrections to the generators $\Delta S_+^{(m)}$ at one order less, $m \leq k - 1$. The commutation relations Eq. (18) can be viewed as, essentially, inhomogeneous first-order differential equations on the evolution kernels. Their solution determines $\mathbb{H}^{(k)}$ up to an $SL(2)$ -invariant term (solution of a homogeneous equation $[\mathbb{H}_{inv}^{(k)}, S_+^{(0)}] = 0$), which can, again, be restored from the spectrum of the anomalous dimensions. This procedure is described in detail for scalar theories in Ref. [10].

Last but not least, in MS-like schemes the evolution kernels (anomalous dimensions) do not depend on the space–time dimension by construction. Indeed, the renormalization \mathbb{Z} factors relating the renormalized and bare light-ray operators $[\mathcal{O}](z_1, z_2) = \mathbb{Z} \mathcal{O}(z_1, z_2)$ are given by the expansion

$$\mathbb{Z} = 1 + \sum_{j=1}^{\infty} \epsilon^{-j} \sum_{k=j}^{\infty} a_s^k \mathbb{Z}_{jk}, \quad (19)$$

where \mathbb{Z}_{jk} have the integral representation similar to (3) in terms of functions of two variables, $Z_{jk}(\alpha, \beta)$ that do not depend on ϵ . Thus, eliminating the ϵ -dependence of the expressions derived in the d -dimensional (conformal) theory for the critical coupling by the expansion $\epsilon = -b_0 a_s^* + \mathcal{O}(a_s^{*2})$ allows one to restore the evolution kernels for the theory in integer dimensions for arbitrary coupling $a_s^* \rightarrow a_s$; this rewriting is simple and exact to all orders.

3. The statement of conformal invariance of QCD in d dimensions at the critical coupling is not trivial. It is believed that “physically reasonable” scale-invariant theories are also conformally invariant, see Ref. [16] for a discussion, however, to the best of our knowledge there is no proof of this statement for $d > 2$ dimensions (but there are no counterexamples as well). In non-gauge theories conformal invariance for the Green functions of basic fields can be checked in perturbative expansions [17]. In gauge theories conformal invariance does not hold for the correlators of basic fields and can be expected only for the Green functions of gauge-invariant operators. For local composite operators a proof of conformal invariance is based on the analysis of pair counterterms for the product of the trace of energy-momentum tensor and local operators [18]. This analysis is beyond the scope of this Letter; it becomes rather complicated in gauge theories due to mixing of gauge invariant operators with BRST variations and equation of motion (EOM) operators [19].

A short comment may, nevertheless, be relevant. Let \mathcal{O}_N be a gauge-invariant multiplicatively renormalizable operator

$$(M\partial_M + \beta(a)\partial_a + \gamma_N(a))[\mathcal{O}_N] = 0, \quad (20)$$

where $\gamma_N(a)$ is the anomalous dimension. As a consequence, it possesses a certain (critical) dimension for the fine-tuned value of the coupling (critical point) $a = a_*$, $\beta(a_*) = 0$:

$$i[\mathbf{D}, [\mathcal{O}_N](x)] = (x\partial_x + \Delta_N^*)[\mathcal{O}_N](x), \quad (21)$$

where \mathbf{D} is the operator of dilatations, Δ_N is the canonical dimension of the operator \mathcal{O}_N , and $\Delta_N^* = \Delta_N + \gamma_N^*$ is the scaling dimension, $\gamma_N^* = \gamma_N(a_*)$.

The statement that $[\mathcal{O}_N](x)$ becomes a conformal operator at the critical point, as widely expected, means that action of the generator of special conformal transformations on this operator takes the form

$$\begin{aligned} i[\mathbf{K}^\mu, [\mathcal{O}_N](x)] &= \left[2x^\mu(x\partial) - x^2\partial^\mu + 2\Delta_N^*x^\mu \right. \\ &\quad \left. + 2x^\nu \left(n^\mu \frac{\partial}{\partial n^\nu} - n_\nu \frac{\partial}{\partial n_\mu} \right) \right] [\mathcal{O}_N](x). \end{aligned} \quad (22)$$

Equivalently, a correlation function of such operators at the critical point must satisfy the Ward identity

$$(K_{x_1}^\mu + \dots + K_{x_n}^\mu) \langle [\mathcal{O}_1](x_1) \dots [\mathcal{O}_n](x_n) \rangle = 0, \quad (23)$$

where it is assumed that all space–time points x_i are different. Calculating the l.h.s. in perturbation theory (see Ref. [18]) and making use of the dilatation Ward identity produces the expression of the form

$$\begin{aligned} (K_{x_1}^\mu + \dots + K_{x_n}^\mu) \langle [\mathcal{O}_1](x_1) \dots [\mathcal{O}_n](x_n) \rangle \\ = \sum_{i=1}^n \langle [\mathcal{O}_1](x_1) \dots \tilde{\mathcal{O}}_i^\mu(x_i) \dots [\mathcal{O}_n](x_n) \rangle, \end{aligned} \quad (24)$$

where $\tilde{\mathcal{O}}_i^\mu(x_i)$ are local operator insertions that involve several contributions: EOM operators, operators representing a BRST variation of another operator, and gauge-invariant operators. The first two can be neglected as they do not contribute to the correlation function (assuming $x_j \neq x_k$, for all $j \neq k$). The last ones can further be expanded in terms of gauge invariant operators $[\mathcal{O}_{iq}^\mu](x_i)$ with a certain critical dimension,

$$\tilde{\mathcal{O}}_i^\mu(x_i) = \sum_q c_q(a) [\mathcal{O}_{iq}^\mu](x_i).$$

Dilatation invariance implies that the both sides of Eq. (24) must have the same scaling dimension at the critical point. Note that application of K^μ lowers the scaling dimension of an operator by one. As a consequence, for each contribution on the r.h.s., either $c_q(a_*)$ vanishes, or the scaling dimensions of $[\mathcal{O}_{iq}^\mu]$ and $[\mathcal{O}_i]$ must differ by one, $\dim \mathcal{O}_{iq}^\mu = \dim \mathcal{O}_i - 1$, to all orders of perturbation theory. If such operators do not exist, then all coefficient $c_q(a)$ have to vanish at the critical point that implies conformal invariance Eq. (23). For the leading-twist operators that are subject of this Letter, absence of operators with the same anomalous dimension and the canonical dimension less by one is easy to verify. For the general case, it would be quite unexpected that two different operators have the same anomalous dimension if they are not related by some exact symmetry, although existence of such pairs cannot be excluded.

4. The correction ΔS_+ , $S_+ = S_+^{(0)} + \Delta S_+$, to the generator of special conformal transformations at the critical point in QCD in d dimensions (in the light-ray operator representation) can be derived from the analysis of CWIs for suitable correlation functions. The standard procedure is to consider the Green functions of twist-two operators with quark fields, see Refs. [3,7] (for $d = 4$), which are not gauge invariant that complicates the analysis. This difficulty can be avoided by considering, instead, the correlator of two light-ray operators that depend on different auxiliary vectors n and \bar{n} :

$$G(x, z, w) = \langle [\mathcal{O}^{(n)}](0, z) [\mathcal{O}^{(\bar{n})}](x, w) \rangle \\ = \mathcal{N}^{-1} \int D\Phi e^{-S_R(\Phi)} [\mathcal{O}^{(n)}](0, z) [\mathcal{O}^{(\bar{n})}](x, w), \quad (25)$$

where $\Phi \equiv \{A, q, \bar{q}, c, \bar{c}\}$ is the set of fundamental fields, $z \equiv \{z_1, z_2\}$, $w \equiv \{w_1, w_2\}$ and we assume that the auxiliary light-like vectors are normalized as $(n \cdot \bar{n}) = 1$.

The QCD Lagrangian in $d = 4 - 2\epsilon$ dimensional Euclidean space-time in covariant gauge has the form²

$$\mathcal{L} = \bar{q}(\not{\partial} - ig\not{A})q + \frac{1}{4}F_{\mu\nu}^a F^{a,\mu\nu} + \partial_\mu \bar{c}^a (D^\mu c)^a + \frac{1}{2\xi}(\partial A^a)^2. \quad (26)$$

The renormalized action S_R is obtained from (26) by the replacement

$$\Phi \rightarrow \Phi_0 = Z_\Phi \Phi, \quad g \rightarrow g_0 = M^\epsilon Z_g g, \\ \xi \rightarrow \xi_0 = Z_\xi \xi, \quad (27)$$

i.e. $S_R(\Phi, g, \xi) = S(\Phi_0, g_0, \xi_0)$. Note that we do not send $\epsilon \rightarrow 0$ in the renormalized correlation functions so that they explicitly depend on ϵ .

The form of the CWI is simpler for the special choice $(n \cdot x) = (\bar{n} \cdot x) = 0$ that we accept for this calculation. For the local conformal operators defined with respect to the \bar{n} vector, $\mathcal{O}_N^{(\bar{n})}(x)$, cf. [11], it follows from Eq. (22) that

$$i[(\bar{n}\mathbf{K}), \mathcal{O}_N^{(\bar{n})}(x)] = -x^2(\bar{n}\partial_x) \mathcal{O}_N^{(\bar{n})}(x). \quad (28)$$

Going over to the light-ray operators one obtains, therefore

$$i[(\bar{n}\mathbf{K}), \mathcal{O}^{(\bar{n})}(x, w)] = -x^2(\bar{n}\partial_x) \mathcal{O}^{(\bar{n})}(x, w). \quad (29)$$

Thus conformal invariance of the correlation function (25) at the critical point implies the constraint

$$\frac{i}{2} \langle [(\bar{n}\mathbf{K}), \mathcal{O}^{(n)}(0, z)] \mathcal{O}^{(\bar{n})}(x, w) + \mathcal{O}^{(n)}(0, z) [(\bar{n}\mathbf{K}), \mathcal{O}^{(\bar{n})}(x, w)] \rangle \\ = \left(S_+^{(z)} - \frac{1}{2}x^2(\bar{n}\partial_x) \right) G(x, z, w) = 0, \quad (30)$$

where the superscript $S_+^{(z)}$ reminds that it is a differential operator acting on z_1, z_2 coordinates.

Explicit expression for $S_+^{(z)}$ can be derived from the CWI

$$0 = -\langle \delta_+ S_R [\mathcal{O}^{(n)}](z) [\mathcal{O}^{(\bar{n})}](x, w) \rangle + \langle \delta_+ [\mathcal{O}^{(n)}](z) [\mathcal{O}^{(\bar{n})}](x, w) \rangle \\ + \langle [\mathcal{O}^{(n)}](z) \delta_+ [\mathcal{O}^{(\bar{n})}](x, w) \rangle, \quad (31)$$

where $[\mathcal{O}](z_1, z_2) \equiv [\mathcal{O}](x=0; z_1, z_2)$, that follows from invariance of the correlation function (25) under a change of variables $\Phi \mapsto \Phi + \delta_+ \Phi$ in the path integral,

$$\delta_+ q(x) = \bar{n}^\mu \left((2x_\mu(x\partial) - x^2\partial_\mu + 2\Delta_q x_\mu) q(x) + \frac{1}{2}[\gamma_\mu, \not{x}] q(x) \right), \\ \delta_+ A_\rho(x) = \bar{n}^\mu \left((2x_\mu(x\partial) - x^2\partial_\mu + 2\Delta_A x_\mu) A_\rho(x) \right. \\ \left. + 2g_{\mu\rho}(xA) - 2x_\rho A_\mu(x) \right), \\ \delta_+ c(x) = \bar{n}^\mu (2x_\mu(x\partial) - x^2\partial_\mu + 2\Delta_c x_\mu) c(x), \\ \delta_+ \bar{c}(x) = \bar{n}^\mu (2x_\mu(x\partial) - x^2\partial_\mu + 2\Delta_{\bar{c}} x_\mu) \bar{c}(x). \quad (32)$$

The choice of the parameters Δ_Φ is a matter of convenience. They can be taken, e.g., equal to the canonical dimensions of the fields in d space-time dimensions, as in [10]. For QCD a different choice proves to be more convenient: $\Delta_q = 3/2 - \epsilon$, $\Delta_A = 1$, $\Delta_c = 2$ and $\Delta_{\bar{c}} = 0$. In this case the quark part of the Lagrangian is invariant and variation of the action takes the form

$$\delta_+ S_R = 4\epsilon \int d^d x (x\bar{n}) (\mathcal{L}_A + \mathcal{L}_\xi + \mathcal{L}_{ghost}) \\ + 2(d-2)\bar{n}^\mu \int d^d x \left(Z_c^2 \bar{c} D_\mu c - \frac{1}{\xi} A_\mu (\partial A) \right). \quad (33)$$

The reason for choosing different scaling dimensions for the ghost and anti-ghost fields is that in this case the last term $\sim (d-2)$ that does not vanish in four dimensions is a BRST variation [1]

$$\left(Z_c^2 \bar{c}^a D_\mu c^a - \frac{1}{\xi} A_\mu^a (\partial A^a) \right) = \delta_{BRST} (\bar{c}^a A_\mu^a). \quad (34)$$

Hence, it does not contribute to the variation of (25).

In this work we are interested in the one-loop correction to the generator S_+ . To this accuracy, obviously, the ghost Lagrangian \mathcal{L}_{ghost} and gluon self-interaction do not contribute i.e. we have to keep terms quadratic in gluon fields only. One obtains after some algebra

$$\delta_+ S_R = -2\epsilon \int d^d x \left[(x\bar{n}) A_\alpha^a K^{\alpha\beta} A_\beta^a + \left(1 + \frac{1}{\xi} \right) (\bar{n}A^a) (\partial A^a) \right] \\ + \dots, \quad (35)$$

where the ellipses stands for the terms that are irrelevant at one loop order and

$$K^{\alpha\beta} = g^{\alpha\beta} \partial^2 - \partial^\alpha \partial^\beta \left(1 - \frac{1}{\xi} \right) \quad (36)$$

is nothing but the inverse gluon propagator (with a minus sign). Moreover, as follows from Eq. (34), the last term $\sim (\bar{n}A)(\partial A)$ can be written as a combination of the BRST variation and the ghost term, so that it does not contribute to the one-loop accuracy as well. Thus, to our accuracy, the insertion of $\delta_+ S_R$ generates an effective vertex insertion $-2\epsilon (\bar{n}x) A_\alpha^a(x) K^{\alpha\beta} A_\beta^a(x)$ in a gluon line in one loop diagrams:

² Our notation follows closely Ref. [20]

$$\begin{aligned}
A_1(\alpha) &= \ln \alpha \ln \bar{\alpha} - \frac{3}{2} \bar{\alpha} - (2 - \alpha) \ln \alpha + \frac{2 + \alpha}{\alpha} \bar{\alpha} \ln \bar{\alpha}, \\
B_1(\alpha, \beta) &= (\beta - \alpha) \left[\frac{3}{2} - \ln(1 - \alpha - \beta) \right] \\
&\quad - \alpha \ln \alpha + \beta \ln \beta + \frac{1}{\alpha} \ln \bar{\alpha} - \frac{1}{\beta} \ln \bar{\beta}, \\
A_3(\alpha) &= \bar{\alpha}, \quad B_3(\alpha, \beta) = \alpha - \beta.
\end{aligned} \tag{46}$$

The simplest way to calculate the evolution kernels $h_1^{(2)}, h_3^{(2)}$ is to try the following ansatz:

$$\begin{aligned}
h_k^{(2)}(\alpha, \beta) &= -\delta_+(\tau) [\phi_k(\alpha) + \phi_k(\beta)] \\
&\quad + \varphi_k(\alpha, \beta) + c_k \delta(\alpha) \delta(\beta).
\end{aligned} \tag{47}$$

Calculating the commutator $[S_+^{(0)}, \mathbb{H}^{(2)}]$ one obtains first-order differential equations on the functions ϕ_k, φ_k (the terms in c_k drop out from the commutator)

$$\begin{aligned}
\bar{\alpha}^2 \partial_\alpha \phi_k(\alpha) &= -A_k(\alpha), \\
(\alpha \bar{\alpha} \partial_\alpha - \beta \bar{\beta} \partial_\beta) \varphi_k(\alpha, \beta) &= B_k(\alpha, \beta),
\end{aligned} \tag{48}$$

where $\partial_\alpha = d/d\alpha$, etc. The solutions can be chosen as

$$\begin{aligned}
\phi_1(\alpha) &= -\ln \bar{\alpha} \left[\frac{3}{2} - \ln \bar{\alpha} + \frac{1 + \bar{\alpha}}{\bar{\alpha}} \ln \alpha \right], \quad \phi_3(\alpha) = \ln \bar{\alpha}, \\
\varphi_1(\alpha, \beta) &= -\theta(1 - \tau) \left[\frac{1}{2} \ln^2(1 - \alpha - \beta) \right. \\
&\quad \left. + \frac{1}{2} \ln^2 \bar{\alpha} + \frac{1}{2} \ln^2 \bar{\beta} - \ln \alpha \ln \bar{\alpha} - \ln \beta \ln \bar{\beta} \right. \\
&\quad \left. - \frac{1}{2} \ln \alpha - \frac{1}{2} \ln \beta + \frac{\bar{\alpha}}{\alpha} \ln \bar{\alpha} + \frac{\bar{\beta}}{\beta} \ln \bar{\beta} \right], \\
\varphi_3(\alpha, \beta) &= -\ln(1 - \alpha - \beta) \theta(1 - \tau).
\end{aligned} \tag{49}$$

They are defined up to arbitrary solutions of the corresponding homogeneous equations: a constant for $\phi_{1,3}(\alpha)$ and a function of the conformal ratio for $\varphi_{1,3}(\alpha, \beta)$. These missing pieces and also the complete kernel $h_2^{(2)}(\tau)$ can be fixed from the known spectrum of the two-loop anomalous dimensions.

The well-known decomposition

$$\gamma_N^{(2)} = m_N^+ + (-1)^{N-1} m_N^-, \tag{50}$$

where m_N^\pm can be extended to analytic functions $m^\pm(N)$ with poles at negative real axis, corresponds for light-ray operators to the decomposition in integration regions where the quark and the antiquark retain their ordering on the light cone or are interchanged

$$h(\tau) = \theta(1 - \tau) h^+(\tau) + \theta(\tau - 1) h^-(\tau^{-1}), \tag{51}$$

corresponding to $\alpha + \beta < 1$ and $\alpha + \beta > 1$, respectively. The equations

$$m^\pm(N) = \int_0^1 d\alpha \int_0^{\bar{\alpha}} d\beta h^\pm(\tau) (1 - \alpha - \beta)^{N-1} \tag{52}$$

can be inverted as

$$h^\pm(\tau) = \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} dN (2N + 1) m^\pm(N) P_N \left(\frac{1 + \tau}{1 - \tau} \right), \tag{53}$$

where P_N is the Legendre function. The integration goes along the imaginary axis such that all poles of $m^\pm(N)$ lie to the left of the integration contour. In practice it turns out to be more efficient to

start from a certain “educated guess” for the kernels, calculate the moments and find the coefficients.

The final result reads

$$\begin{aligned}
h_1^{(2)}(\alpha, \beta) &= -\delta_+(\tau) (\phi_1(\alpha) + \phi_1(\beta)) + \varphi_1(\alpha, \beta) \\
&\quad + \theta(\bar{\tau}) \left[2 \text{Li}_2(\tau) + \ln^2 \bar{\tau} + \ln \tau - \frac{1 + \bar{\tau}}{\tau} \ln \bar{\tau} \right] \\
&\quad + \theta(-\bar{\tau}) \left[\ln^2(-\bar{\tau}/\tau) - \frac{2}{\tau} \ln(-\bar{\tau}/\tau) \right] \\
&\quad + \left(-6\zeta(3) + \frac{1}{3} \pi^2 + \frac{21}{8} \right) \delta(\alpha) \delta(\beta), \\
h_2^{(2)}(\alpha, \beta) &= \frac{1}{3} (\pi^2 - 4) \delta_+(\tau) - 2\theta(\bar{\tau}) \\
&\quad \times \left[\text{Li}_2(\tau) - \text{Li}_2(1) + \frac{1}{2} \ln^2 \bar{\tau} - \frac{1}{\tau} \ln \bar{\tau} + \frac{5}{3} \right] \\
&\quad - \theta(-\bar{\tau}) \left[\ln^2(-\bar{\tau}/\tau) - \frac{2}{\tau} \ln(-\bar{\tau}/\tau) \right] \\
&\quad + \left(6\zeta(3) - \frac{2}{3} \pi^2 + \frac{13}{6} \right) \delta(\alpha) \delta(\beta), \\
h_3^{(2)}(\alpha, \beta) &= -\delta_+(\tau) \left[\ln \bar{\alpha} + \ln \bar{\beta} + \frac{5}{3} \right] \\
&\quad - \theta(\bar{\tau}) \left[\ln(1 - \alpha - \beta) + \frac{11}{3} \right] + \frac{13}{12} \delta(\alpha) \delta(\beta),
\end{aligned} \tag{54}$$

where $\bar{\tau} = 1 - \tau$, and the functions $\phi_1(\alpha)$ and $\varphi_1(\alpha, \beta)$ are given in Eq. (49).

6. Our result for the two-loop evolution of flavor-nonsinglet light-ray operators in Eqs. (12), (54) is equivalent to the corresponding evolution equation for GPDs obtained in Ref. [1] but is more compact and has manifest $SL(2)$ symmetry properties. The latter feature presents the crucial advantage of the light-ray operator (alias position space) representation which makes this technique attractive for higher-order calculations. Exact conformal symmetry of QCD at the critical point proves to be very helpful as it provides one with algebraic group-theory methods to calculate the commutators of integral operators that appear in Eq. (18). Evolution equations for GPDs can be obtained from our expressions by a Fourier transformation which is rather straightforward, cf. [21].

Apart from the evolution kernels, another new result is the calculation of the generator of special conformal transformations S_+ to the one-loop accuracy, see Eq. (15). The QCD expression differs from the corresponding result in the scalar theory [10] but remains simple. As we have demonstrated, this result can be obtained from the gauge-invariant correlation function of two light-ray operators, thus bypassing the complications due to non-gauge-invariant contributions in the usual approach dealing with Green functions involving fundamental fields. We expect that the same technique can be used for the flavor-singlet light-ray operators and for the calculation of S_+ to the two-loop accuracy, which is the first step towards the NNLO evolution equations. This task, obviously, goes beyond the scope of this Letter.

Acknowledgements

We thank D. Müller for useful discussions. This work was supported by the DFG, grant BR2021/5-2.

References

- [1] A.V. Belitsky, A. Freund, D. Müller, Nucl. Phys. B 574 (2000) 347.

- [2] A.V. Belitsky, D. Müller, Nucl. Phys. B 537 (1999) 397.
- [3] A.V. Belitsky, D. Müller, Nucl. Phys. B 527 (1998) 207.
- [4] S. Moch, J.A.M. Vermaseren, A. Vogt, Nucl. Phys. B 688 (2004) 101.
- [5] A. Vogt, S. Moch, J.A.M. Vermaseren, Nucl. Phys. B 691 (2004) 129.
- [6] D. Müller, Z. Phys. C 49 (1991) 293.
- [7] D. Müller, Phys. Rev. D 49 (1994) 2525.
- [8] D. Müller, Phys. Rev. D 58 (1998) 054005.
- [9] A.V. Belitsky, D. Müller, Phys. Lett. B 417 (1998) 129.
- [10] V.M. Braun, A.N. Manashov, Eur. Phys. J. C 73 (2013) 2544.
- [11] V.M. Braun, G.P. Korchemsky, D. Mueller, Prog. Part. Nucl. Phys. 51 (2003) 311.
- [12] I.I. Balitsky, V.M. Braun, Nucl. Phys. B 311 (1989) 541.
- [13] V.M. Braun, S.E. Derkachov, G.P. Korchemsky, A.N. Manashov, Nucl. Phys. B 553 (1999) 355.
- [14] T. Banks, A. Zaks, Nucl. Phys. B 196 (1982) 189.
- [15] A. Hasenfratz, P. Hasenfratz, Phys. Lett. B 297 (1992) 166.
- [16] Y. Nakayama, A lecture note on scale invariance vs conformal invariance, arXiv: 1302.0884 [hep-th].
- [17] S.E. Derkachov, N.A. Kivel, A.S. Stepanenko, A.N. Vasiliev, Theor. Math. Phys. 92 (1992) 1047.
- [18] A.N. Vasil'ev, Quantum Field Renormalization Group in the Theory of Critical Behaviour and Stochastic Dynamics, PNPI Press, Sankt-Petersburg, 1998, 774 pp.
- [19] J.C. Collins, Renormalization, CUP, 1984.
- [20] M. Ciuchini, S.E. Derkachov, J.A. Gracey, A.N. Manashov, Phys. Lett. B 458 (1999) 117; M. Ciuchini, S.E. Derkachov, J.A. Gracey, A.N. Manashov, Nucl. Phys. B 579 (2000) 56.
- [21] V.M. Braun, A.N. Manashov, B. Pirnay, Phys. Rev. D 80 (2009) 114002; V.M. Braun, A.N. Manashov, B. Pirnay, Phys. Rev. D 86 (2012) 119902 (Erratum).