



The Chiral index of the fermionic
signature operator

Felix Finster

Preprint Nr. 06/2014

THE CHIRAL INDEX OF THE FERMIONIC SIGNATURE OPERATOR

FELIX FINSTER

APRIL 2014

ABSTRACT. We define an index of the fermionic signature operator on even-dimensional globally hyperbolic spin manifolds of finite lifetime. The invariance of the index under homotopies is studied. The definition is generalized to causal fermion systems with a chiral grading. We give examples of space-times and Dirac operators thereon for which our index is non-trivial.

CONTENTS

1. Introduction	1
2. Preliminaries	2
3. The Chiral Index	4
4. Generalization to the Setting of Causal Fermion Systems	5
5. The Chiral Index in the Massless Odd Case	7
6. Homotopy Invariance	7
7. Example: Shift Operators in the Setting of Causal Fermion Systems	9
8. Example: A Dirac Operator with $\text{ind}_0 S \neq 0$	10
9. Example: A Dirac Operator with $\text{ind} S \neq 0$	14
10. Examples Illustrating the Homotopy Invariance	18
11. Conclusion and Outlook	20
References	21

1. INTRODUCTION

In the recent papers [6, 7] the fermionic signature operator was introduced on globally hyperbolic Lorentzian spin manifolds. It is a bounded symmetric operator on the Hilbert space of solutions of the Dirac equation which depends on the global geometry of space-time. This raises the question how the geometry of space-time is related to spectral properties of the fermionic signature operator. The first step in developing the resulting “Lorentzian spectral geometry” is the paper [5] where the simplest situation of Lorentzian surfaces is considered. In the present paper, we proceed in a somewhat different direction and show that there is a nontrivial index associated to the fermionic signature operator. This is the first time that an index is defined for a geometric operator on a Lorentzian manifold.

We make essential use of the decomposition of spinors in even space-time dimension into left- and right-handed components (the “chiral grading”). The basic idea is to

Supported in part by the Deutsche Forschungsgemeinschaft.

decompose the fermionic signature operator \mathcal{S} using the chiral grading as

$$\mathcal{S} = \mathcal{S}_L + \mathcal{S}_R \quad \text{with} \quad \mathcal{S}_L^* = \mathcal{S}_R, \quad (1.1)$$

and to define the so-called *chiral index* of \mathcal{S} as the Noether index of \mathcal{S}_L . After providing the necessary preliminaries (Section 2), this definition will be given in Section 3 in space-times of finite lifetime. In order to work out the mathematical essence of our index, in Section 4 we also give its definition in the general setting of causal fermion systems (for an introduction to causal fermion systems see [3] or [4]). Section 5 is devoted to a variant of the chiral index which applies in the special case of the massless Dirac equation and a Dirac operator which is odd with respect to the chiral grading. In Section 6 we analyze the invariance properties of the chiral indices when space-time or the Dirac operator are deformed by a homotopy. In Sections 8–10 we construct examples of fermionic signature operators with a non-trivial index and illustrate the homotopy invariance. Finally, in Section 11 we discuss our results and give an outlook on potential extensions and applications, like the generalization to space-times of infinite lifetime.

We point out that the purpose of this paper is to define the chiral index, to study a few basic properties and to show in simple examples that it is in general non-trivial. But we do not work out any physical applications, nor do we make the connection to geometric or topological invariants. These shortcomings are mainly due to the fact that we only succeeded in computing the index explicitly in highly symmetric and rather artificial examples. Moreover, it does not seem easy to verify the conditions needed for the homotopy invariance. For these reasons, we leave physically interesting examples and geometric stability results as a subject of future research. All we can say for the moment is that the chiral index describes the “chiral asymmetry” of the Dirac operator in terms of an integer. This integer seems to depend on the geometry of the boundary of space-time and on the singular behavior of the potentials in the Dirac equation. Smooth potentials in the Dirac equation, however, tend to not affect the index.

2. PRELIMINARIES

We recall a few basic constructions from [6]. Let (\mathcal{M}, g) be a smooth, globally hyperbolic Lorentzian spin manifold of even dimension $k \geq 2$. For the signature of the metric we use the convention $(+, -, \dots, -)$. We denote the spinor bundle by $S\mathcal{M}$. Its fibers $S_x\mathcal{M}$ are endowed with an inner product $\langle \cdot | \cdot \rangle_x$ of signature (n, n) with $n = 2^{k/2-1}$ (for details see [1, 8]), which we refer to as the spin scalar product. Clifford multiplication is described by a mapping γ which satisfies the anti-commutation relations,

$$\gamma : T_x\mathcal{M} \rightarrow \text{L}(S_x\mathcal{M}) \quad \text{with} \quad \gamma(u)\gamma(v) + \gamma(v)\gamma(u) = 2g(u, v)\mathbf{1}_{S_x(\mathcal{M})}.$$

We write Clifford multiplication in components with the Dirac matrices γ^j and use the short notation with the Feynman dagger, $\gamma(u) \equiv u^j \gamma_j \equiv \not{u}$. The metric connections on the tangent bundle and the spinor bundle are denoted by ∇ .

In the even-dimensional situation under consideration, the spinor bundle has a decomposition into left- and right-handed components. We describe this chiral grading by an operator Γ (the “pseudoscalar operator,” in physics usually denoted by γ^5),

$$\Gamma : S_x\mathcal{M} \rightarrow S_x\mathcal{M},$$

having for all $u \in T_x \mathcal{M}$ the properties

$$\Gamma^* = -\Gamma, \quad \Gamma^2 = \mathbf{1}, \quad \Gamma \gamma(u) = -\gamma(u) \Gamma, \quad \nabla \Gamma = 0 \quad (2.1)$$

(where the star denotes the adjoint with respect to the spin scalar product). We denote the chiral projections to the left- and right-handed components by

$$\chi_L = \frac{1}{2}(\mathbf{1} - \Gamma) \quad \text{and} \quad \chi_R = \frac{1}{2}(\mathbf{1} + \Gamma). \quad (2.2)$$

The sections of the spinor bundle are also referred to as wave functions. We denote the smooth sections of the spinor bundle by $C^\infty(\mathcal{M}, S\mathcal{M})$. Similarly, $C_0^\infty(\mathcal{M}, S\mathcal{M})$ denotes the smooth sections with compact support. On the compactly supported wave functions, one can introduce the Lorentz invariant inner product

$$\langle \cdot | \cdot \rangle : C_0^\infty(\mathcal{M}, S\mathcal{M}) \times C_0^\infty(\mathcal{M}, S\mathcal{M}) \rightarrow \mathbb{C}, \quad (2.3)$$

$$\langle \psi | \phi \rangle := \int_{\mathcal{M}} \langle \psi | \phi \rangle_x d\mu_{\mathcal{M}}. \quad (2.4)$$

The Dirac operator \mathcal{D} is defined by

$$\mathcal{D} := i\gamma^j \nabla_j + \mathcal{B} : C^\infty(\mathcal{M}, S\mathcal{M}) \rightarrow C^\infty(\mathcal{M}, S\mathcal{M}),$$

where $\mathcal{B} \in L(S_x)$ (the ‘‘external potential’’) typically is a smooth multiplication operator which is symmetric with respect to the spin scalar product. In some of our examples, \mathcal{B} will be chosen more generally as a convolution operator which is symmetric with respect to the inner product (2.4). For a given real parameter $m \in \mathbb{R}$ (the ‘‘mass’’), the Dirac equation reads

$$(\mathcal{D} - m)\psi = 0.$$

We mainly consider solutions in the class $C_{sc}^\infty(\mathcal{M}, S\mathcal{M})$ of smooth sections with spatially compact support. On such solutions one has the scalar product

$$(\psi | \phi) = 2\pi \int_{\mathcal{N}} \langle \psi | \psi \phi \rangle_x d\mu_{\mathcal{N}}(x), \quad (2.5)$$

where \mathcal{N} denotes any Cauchy surface and ν its future-directed normal. Due to current conservation, the scalar product is independent of the choice of \mathcal{N} (for details see [6, Section 2]). Forming the completion gives the Hilbert space $(\mathcal{H}_m, (\cdot | \cdot))$.

For the construction of the fermionic signature operator, we need to extend the bilinear form (2.4) to the solution space of the Dirac equation. In order to ensure that the integral in (2.4) exists, we need to make the following assumption (for more details see [6, Section 3.2]).

Definition 2.1. *A globally hyperbolic space-time (\mathcal{M}, g) is said to be **m-finite** if there is a constant $c > 0$ such that for all $\phi, \psi \in \mathcal{H}_m \cap C_{sc}^\infty(\mathcal{M}, S\mathcal{M})$, the function $\langle \phi | \psi \rangle_x$ is integrable on \mathcal{M} and*

$$|\langle \phi | \psi \rangle| \leq c \|\phi\| \|\psi\|$$

(where $\|\cdot\| = (\cdot | \cdot)^{\frac{1}{2}}$ is the norm on \mathcal{H}_m).

Under this assumption, the space-time inner product is well-defined as a bounded bilinear form on \mathcal{H}_m ,

$$\langle \cdot | \cdot \rangle : \mathcal{H}_m \times \mathcal{H}_m \rightarrow \mathbb{C}.$$

Applying the Riesz representation theorem, we can uniquely represent this bilinear form with a signature operator \mathcal{S} ,

$$\mathcal{S} : \mathcal{H}_m \rightarrow \mathcal{H}_m \quad \text{with} \quad \langle \phi | \psi \rangle = (\phi | \mathcal{S} \psi). \quad (2.6)$$

We refer to \mathcal{S} as the **fermionic signature operator**. It is obviously a bounded symmetric operator on \mathcal{H}_m . We note that the construction of the fermionic signature operator is manifestly covariant and independent of the choice of a Cauchy surface.

3. THE CHIRAL INDEX

We now modify the construction of the fermionic signature operator by inserting the chiral projection operators into (2.4). We thus obtain the bilinear forms

$$\langle \psi | \phi \rangle_{L/R} = \int_{\mathcal{M}} \langle \psi | \chi_{L/R} \phi \rangle_x d\mu_{\mathcal{M}}. \quad (3.1)$$

For the space-time integrals to exist, we need the following assumption.

Definition 3.1. *A globally hyperbolic space-time (\mathcal{M}, g) is said to be Γ -finite if there is a constant $c > 0$ such that for all $\phi, \psi \in \mathcal{H}_m \cap C_{sc}^\infty(\mathcal{M}, S\mathcal{M})$, the function $\langle \phi | \Gamma \psi \rangle_x$ is integrable on \mathcal{M} and*

$$|\langle \phi | \Gamma \psi \rangle| \leq c \|\phi\| \|\psi\|.$$

There seems no simple relation between m -finiteness and Γ -finiteness. But both conditions are satisfied if we assume that the space-time (\mathcal{M}, g) has **finite lifetime** in the sense that it admits a foliation $(\mathcal{N}_t)_{t \in (t_0, t_1)}$ by Cauchy surfaces with $t_0, t_1 \in \mathbb{R}$ such that the function $\langle \nu, \partial_t \rangle$ is bounded on \mathcal{M} (see [6, Definition 3.4]). The following proposition is an immediate generalization of [6, Proposition 3.5].

Proposition 3.2. *Every globally hyperbolic manifold of finite lifetime is m -finite and Γ -finite.*

Proof. Let $\psi \in \mathcal{H}_m \cap C_{sc}^\infty(\mathcal{M}, S\mathcal{M})$ and $C(x)$ one of the operators $\mathbf{1}_{S_x}$ or $i\Gamma(x)$. Applying Fubini's theorem and decomposing the volume measure, we obtain

$$\langle \psi | C \psi \rangle = \int_{\mathcal{M}} \langle \psi | C \psi \rangle(x) d\mu_{\mathcal{M}}(x) = \int_{t_0}^{t_1} \int_{\mathcal{N}_t} \langle \psi | C \psi \rangle \langle \nu, \partial_t \rangle dt d\mu_{\mathcal{N}_t}$$

and thus

$$|\langle \psi | C \psi \rangle| \leq \sup_{\mathcal{M}} \langle \nu, \partial_t \rangle \int_{t_0}^{t_1} dt \int_{\mathcal{N}_t} |\langle \psi | C \psi \rangle| d\mu_{\mathcal{N}_t}.$$

Rewriting the integrand as

$$|\langle \psi | C \psi \rangle| = |\langle \psi | \not\psi (\not\psi C) \psi \rangle|,$$

the bilinear form $\langle \cdot | \not\psi \cdot \rangle$ is a scalar product. Moreover, the operator $\not\psi C$ is symmetric with respect to this scalar product. Using that

$$(\not\psi)^2 = \mathbf{1} = (i\not\psi\Gamma)^2,$$

we conclude that the sup-norm corresponding to the scalar product $\langle \cdot | \not\psi \cdot \rangle$ of the operator $\not\psi C$ is equal to one. Hence

$$\int_{\mathcal{N}_t} |\langle \psi | C \psi \rangle| d\mu_{\mathcal{N}_t} \leq \int_{\mathcal{N}_t} \langle \psi | \not\psi \psi \rangle d\mu_{\mathcal{N}_t} = (\psi | \psi),$$

and consequently

$$|\langle \psi | C\psi \rangle| \leq (t_1 - t_0) \sup_{\mathcal{M}} \langle \nu, \partial_t \rangle \|\psi\|^2.$$

Polarization and a denseness argument give the result. \square

Assuming that our space-time is m -finite and Γ -finite, the bilinear forms (3.1) are bounded on $\mathcal{H}_m \times \mathcal{H}_m$. Thus we may represent them with respect to the Hilbert space scalar product in terms of signature operators $\mathcal{S}_{L/R}$,

$$\mathcal{S}_{L/R} : \mathcal{H}_m \rightarrow \mathcal{H}_m \quad \text{with} \quad \langle \phi | \psi \rangle_{L/R} = (\phi | \mathcal{S}_{L/R} \psi). \quad (3.2)$$

We refer to $\mathcal{S}_{L/R}$ as the **chiral signature operators**. Taking the complex conjugate of the equation in (3.2) and using that $\chi_L^* = \chi_R$, we find that (1.1) holds, where the star denotes the adjoint in $L(\mathcal{H}_m)$.

We now define the chiral index as the Noether index of \mathcal{S}_L (sometimes called Fredholm index; for basics see for example [9, §27.1]).

Definition 3.3. *The fermionic signature operator is said to have finite chiral index if the operators of \mathcal{S}_L and \mathcal{S}_R both have a finite-dimensional kernel. The **chiral index** of the fermionic signature operator is defined by*

$$\text{ind } \mathcal{S} = \dim \ker \mathcal{S}_L - \dim \ker \mathcal{S}_R. \quad (3.3)$$

4. GENERALIZATION TO THE SETTING OF CAUSAL FERMION SYSTEMS

Our starting point is a causal fermion system as introduced in [3].

Definition 4.1. Given a complex Hilbert space $(\mathcal{H}, \langle \cdot | \cdot \rangle_{\mathcal{H}})$ (the “particle space”) and a parameter $n \in \mathbb{N}$ (the “spin dimension”), we let $\mathcal{F} \subset L(\mathcal{H})$ be the set of all self-adjoint operators on \mathcal{H} of finite rank, which (counting with multiplicities) have at most n positive and at most n negative eigenvalues. On \mathcal{F} we are given a positive measure ρ (defined on a σ -algebra of subsets of \mathcal{F}), the so-called *universal measure*. We refer to $(\mathcal{H}, \mathcal{F}, \rho)$ as a *causal fermion system*.

Starting from a Lorentzian spin manifold, one can construct a corresponding causal fermion system by choosing \mathcal{H} as a suitable subspace of the solution space of the Dirac equation, forming the local correlation operators (possibly introducing an ultraviolet regularization) and defining ρ as the push-forward of the volume measure on \mathcal{M} (see [6, Section 4] or the examples in [4]). The advantage of working with a causal fermion system is that the underlying space-time does not need to be a Lorentzian manifold, but it can be a more general “quantum space-time” (for more details see [2]).

We now recall a few basic notions from [3]. On \mathcal{F} we consider the topology induced by the operator norm $\|A\| := \sup\{\|Au\|_{\mathcal{H}} \mid \|u\|_{\mathcal{H}} = 1\}$. For every $x \in \mathcal{F}$ we define the *spin space* S_x by $S_x = x(\mathcal{H})$; it is a subspace of \mathcal{H} of dimension at most $2n$. On S_x we introduce the *spin scalar product* $\langle \cdot | \cdot \rangle_x$ by

$$\langle u | v \rangle_x = -\langle u | xu \rangle_{\mathcal{H}} \quad (\text{for all } u, v \in S_x); \quad (4.1)$$

it is an indefinite inner product of signature (p, q) with $p, q \leq n$. Moreover, we define *space-time* M as the support of the universal measure, $M = \text{supp } \rho$. It is a closed subset of \mathcal{F} .

In order to extend the chiral grading to causal fermion systems, we assume for every $x \in M$ an operator $\Gamma(x) \in L(\mathcal{H})$ with the properties

$$\Gamma(x)|_{S_x} : S_x \rightarrow S_x \quad \text{and} \quad x\Gamma(x) = -\Gamma(x)^*x. \quad (4.2)$$

We define the operators $\chi_{L/R}(x) \in \mathbf{L}(\mathcal{H})$ again by (2.2). In order to explain the equations (4.2), we first note that the right side of (4.2) obviously vanishes on the orthogonal complement of S_x . Using furthermore that, by definition of the spin space, the operator x is invertible on S_x , we infer that

$$\Gamma(x)|_{S_x^\perp} = 0.$$

Moreover, the computation

$$\begin{aligned} \prec \psi | \Gamma(x) \phi \succ_x &= -\langle \psi | x \Gamma(x) \phi \rangle_{\mathcal{H}} = -\langle \Gamma(x)^* x \psi | \phi \rangle_{\mathcal{H}} \\ &\stackrel{(4.2)}{=} \langle x \Gamma(x) \psi | \phi \rangle_{\mathcal{H}} = -\prec \Gamma(x) \psi | \phi \succ_x \end{aligned}$$

(with $\psi, \phi \in S_x$) shows that $\Gamma(x) \in \mathbf{L}(S_x)$ is antisymmetric with respect to the spin scalar product. Thus the first equation in (2.1) again holds. This implies that the adjoint of $\chi_L(x)$ with respect to $\prec \cdot | \cdot \succ_x$ equals $\chi_R(x)$. However, we point out that our assumptions (4.2) do not imply that $\Gamma(x)$ is idempotent (in the sense that $\Gamma(x)^2|_{S_x} = \mathbf{1}_{S_x}$). Hence the analog of the second equation in (2.1) does not need to hold on a causal fermion system. This property could be imposed in addition, but will not be needed here. The last two relations in (2.1) do not have an obvious correspondence on causal fermion systems, and they will also not be needed in what follows.

We now have all the structures needed for defining the fermionic signature operator and its chiral index. Namely, replacing the scalar product in (2.6) by the scalar product on the particle space $\langle \cdot | \cdot \rangle_{\mathcal{H}}$, we now demand in analogy to (2.4) and (2.6) that the relation

$$\langle u | \mathcal{S} v \rangle_{\mathcal{H}} = \int_M \prec u | v \succ_x d\rho(x)$$

should hold for all $u, v \in \mathcal{H}$. Using (4.1), we find that the fermionic signature operator is given by the integral

$$\mathcal{S} = - \int_M x d\rho(x).$$

Similarly, the left-handed signature operator can be introduced by

$$\mathcal{S}_L = - \int_M x \chi_L d\rho(x). \quad (4.3)$$

In the setting on a globally hyperbolic manifold, we had to assume that the manifold was m -finite and Γ -finite (see Definitions 2.1 and 3.1). Now we need to assume correspondingly that the integral (4.3) converges. For the sake of larger generality we prefer to work with weak convergence.

Definition 4.2. *The topological fermion system is \mathcal{S}_L -bounded if the integral in (4.3) converges weakly to a bounded operator, i.e. if there is an operator $\mathcal{S}_L \in \mathbf{L}(\mathcal{H})$ such that for all $u, v \in \mathcal{H}$,*

$$- \int_M \langle u | x \chi_L v \rangle_{\mathcal{H}} d\rho(x) = \langle u | \mathcal{S}_L v \rangle_{\mathcal{H}}.$$

Introducing the right-handed signature operator by $\mathcal{S}_R := \mathcal{S}_L^*$, we can define the **chiral index** again by (3.3).

5. THE CHIRAL INDEX IN THE MASSLESS ODD CASE

We return to the setting of Section 3 and consider the special case that the mass vanishes and that the Dirac operator is odd,

$$m = 0 \quad \text{and} \quad \Gamma \mathcal{D} = -\mathcal{D} \Gamma. \quad (5.1)$$

In this case, the solution space of the Dirac equation is obviously invariant under Γ ,

$$\Gamma : \mathcal{H}_0 \rightarrow \mathcal{H}_0.$$

Taking the adjoint with respect to the scalar product (2.5) and noting that Γ anti-commutes with ψ , one sees that Γ is symmetric on \mathcal{H}_0 . Hence χ_L and χ_R are orthogonal projection operators, giving rise to the orthogonal sum decomposition

$$\mathcal{H}_0 = \mathcal{H}_L \oplus \mathcal{H}_R \quad \text{with} \quad \mathcal{H}_{L/R} := \chi_{L/R} \mathcal{H}_0. \quad (5.2)$$

Moreover, the computation

$$\begin{aligned} \langle \chi_L \psi | \chi_{c'} \phi \rangle_c &= \int_{\mathcal{M}} \langle \chi_L \psi | \chi_c \chi_{c'} \phi \rangle_x d\mu_{\mathcal{M}} \\ &= \int_{\mathcal{M}} \langle \psi | \chi_R \chi_c \chi_{c'} \phi \rangle_x d\mu_{\mathcal{M}} = \delta_{Rc} \delta_{cc'} \langle \psi | \phi \rangle_c \end{aligned}$$

with $c, c' \in \{L, R\}$ (and similarly for L replaced by R) shows that \mathcal{S} maps the right-handed component to the left-handed component and vice versa. Moreover, in a block matrix notation corresponding to the decomposition (5.2), the operators \mathcal{S}_L and \mathcal{S}_R have the simple form

$$\mathcal{S}_L = \begin{pmatrix} 0 & 0 \\ A & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{S}_R = \begin{pmatrix} 0 & A^* \\ 0 & 0 \end{pmatrix}$$

with a bounded operator $A : \mathcal{H}_L \rightarrow \mathcal{H}_R$. As a consequence, both \mathcal{S}_L and \mathcal{S}_R have an infinite-dimensional kernel, so that the index cannot be defined by (3.3). This problem can easily be cured by restricting the operators to the respective subspace \mathcal{H}_L and \mathcal{H}_R .

Definition 5.1. *In the massless odd case (5.1), the fermionic signature operator is said to have finite chiral index if the operators $\mathcal{S}_L|_{\mathcal{H}_L}$ and $\mathcal{S}_R|_{\mathcal{H}_R}$ both have a finite-dimensional kernel. We define the index $\text{ind}_0 \mathcal{S}$ by*

$$\text{ind}_0 \mathcal{S} = \dim \ker(\mathcal{S}_L)|_{\mathcal{H}_L} - \dim \ker(\mathcal{S}_R)|_{\mathcal{H}_R}.$$

6. HOMOTOPY INVARIANCE

We first recall Dieudonné's general theorem on the homotopy invariance of the Noether index (see for example [9, Theorem 27.1.5]).

Theorem 6.1. *Let $T(t) : U \rightarrow V$, $0 \leq t \leq 1$, be a one-parameter family of bounded linear operators between Banach spaces U and V which is continuous in the norm topology. If for every $t \in [0, 1]$ the vector spaces*

$$\ker T \text{ and } V/T(\mathcal{H}) \text{ are both finite-dimensional,} \quad (6.1)$$

then

$$\text{ind } T(0) = \text{ind } T(1),$$

where $\text{ind } T := \dim \ker(T) - \dim V/T(\mathcal{H})$.

In most applications of this theorem, one knows from general arguments that the index of T remains finite under homotopies (for example, in the prominent example of the Atiyah-Singer index, this follows from elliptic estimates on a compact manifold). For our chiral index, however, there is no general reason why the chiral index of \mathcal{S} should remain finite. Indeed, the fermionic signature operator is bounded and typically has many eigenvalues near zero. It may well happen that for a certain value of t , an infinite number of these eigenvalues becomes zero (for an explicit example see Example 10.1 below).

Another complication when applying Theorem 6.1 to the fermionic signature operator is that the image of \mathcal{S}_L does not need to be a closed subspace of our Hilbert space. To explain the difficulty, we first consider the chiral index of Definition 3.3. Using that $\ker \mathcal{S}_R = \ker \mathcal{S}_L^* = \mathcal{S}_L(\mathcal{H}_m)^\perp$, the assumption that the fermionic signature operator has finite chiral index can be restated that the vector spaces $\ker \mathcal{S}_L$ and $\mathcal{S}_L(\mathcal{H}_m)^\perp$ are finite-dimensional subspaces of \mathcal{H}_m . Since

$$\dim \mathcal{S}_L(\mathcal{H}_m)^\perp = \dim \mathcal{H}_m / (\overline{\mathcal{S}_L(\mathcal{H}_m)}),$$

this implies that the *closure* of the image of \mathcal{S}_L has finite co-dimension. If the image of \mathcal{S}_L were closed in \mathcal{H}_m , the finiteness of the chiral index would imply that the conditions (6.1) hold if we set $T = \mathcal{S}_L$ and $U = V = \mathcal{H}_m$. However, the image of \mathcal{S}_L will in general *not* be a closed subspace of \mathcal{H}_m , and in this case it is possible that the condition (6.1) is violated for $T = \mathcal{S}_L$ and $U = V = \mathcal{H}_m$, although \mathcal{S} has finite chiral index (according to Definition 3.3). In the massless odd case, the analogous problem occurs if we choose $T = \mathcal{S}_L$, $U = \mathcal{H}_L$ and $V = \mathcal{H}_R$ (see Definition 5.1).

Our method for making Theorem 6.1 applicable is to endow a subspace of the Hilbert space with a finer topology, such that the image of \mathcal{S}_L lies in this subspace and is closed in this topology.

Theorem 6.2. *Let $\mathcal{S}(t) : \mathcal{H}_m \rightarrow \mathcal{H}_m$, $t \in [0, 1]$, be a family of fermionic signature operators with finite chiral index. Let E be a Banach space together with an embedding $\iota : E \hookrightarrow \mathcal{H}_m$ with the following properties:*

- (i) *For every $t \in [0, 1]$, the image of $\mathcal{S}_L(t)$ lies in $\iota(E)$, giving rise to the mapping*

$$\mathcal{S}_L(t) : \mathcal{H}_m \rightarrow E. \quad (6.2)$$

- (ii) *For every $t \in [0, 1]$, the image of the operator \mathcal{S}_L , (6.2), is a closed subspace of E .*

- (iii) *The family $\mathcal{S}_L(t) : \mathcal{H}_m \rightarrow E$ is continuous in the norm topology.*

Then the chiral index is a homotopy invariant,

$$\text{ind } \mathcal{S}(0) = \text{ind } \mathcal{S}(1).$$

In the chiral odd case, the analogous result is stated as follows.

Theorem 6.3. *Let $\mathcal{S}(t) : \mathcal{H}_0 \rightarrow \mathcal{H}_0$, $t \in [0, 1]$, be a family of fermionic signature operators of finite chiral index in the massless odd case (see (5.1)). Moreover, let E be a Banach space together with an embedding $\iota : E \hookrightarrow \mathcal{H}_R$ such that the operator $\mathcal{S}_L|_{\mathcal{H}_L} : \mathcal{H}_L \rightarrow \mathcal{H}_R$ has the following properties:*

- (i) *For every $t \in [0, 1]$, the image of $\mathcal{S}_L(t)$ lies in $\iota(E)$, giving rise to the mapping*

$$\mathcal{S}_L(t) : \mathcal{H}_L \rightarrow E. \quad (6.3)$$

- (ii) *For every $t \in [0, 1]$, the image of the operator \mathcal{S}_L , (6.3), is a closed subspace of E .*

(iii) *The family $\mathcal{S}_L(t) : \mathcal{H}_L \rightarrow E$ is continuous in the norm topology.*

Then the chiral index in the massless odd case is a homotopy invariant,

$$\text{ind}_0 \mathcal{S}(0) = \text{ind}_0 \mathcal{S}(1).$$

In Example 10.2 below, it will be explained how these theorems can be applied.

7. EXAMPLE: SHIFT OPERATORS IN THE SETTING OF CAUSAL FERMION SYSTEMS

In the remainder of this paper we illustrate the previous constructions in several examples. The simplest examples for fermionic signature operators with a non-trivial chiral index can be given in the setting of causal fermion systems. We let $\mathcal{H} = \ell^2(\mathbb{N})$ be the square-summable sequences with the scalar product

$$\langle u|v \rangle_{\mathcal{H}} = \sum_{l=1}^{\infty} \overline{u_l} v_l.$$

For any $k \in \mathbb{N}$ we define the operators x_k by

$$(x_k u)_k = -u_{k+1}, \quad (x_k u)_{k+1} = -u_k,$$

and all other components of $x_k u$ vanish. Thus, writing the series in components,

$$x_k u = \left(\underbrace{0, \dots, 0}_{k-1 \text{ entries}}, -u_{k+1}, -u_k, 0, \dots \right). \quad (7.1)$$

Every operator x_k obviously has rank two with the non-trivial eigenvalues ± 1 . We let μ be the counting measure on \mathbb{N} and $\rho = x_*(\mu)$ the push-forward measure of the mapping $x : k \mapsto x_k \in \mathcal{F} \subset L(\mathcal{H})$. We thus obtain a causal fermion system $(\mathcal{H}, \mathcal{F}, \rho)$ of spin dimension one.

Next, we introduce the pseudoscalar operators $\Gamma(x_k)$ by

$$\Gamma(x_k) u = \left(\underbrace{0, \dots, 0}_{k-1 \text{ entries}}, u_k, -u_{k+1}, 0, \dots \right). \quad (7.2)$$

Obviously, these operators have the properties (4.2). Moreover,

$$\begin{aligned} x \chi_L(x_k) u &= (0, \dots, 0, -u_{k+1}, 0, 0, \dots) \\ x \chi_R(x_k) u &= (0, \dots, 0, 0, -u_k, 0, \dots). \end{aligned}$$

Consequently, the operators

$$\mathcal{S}_{L/R} = - \sum_{k=1}^{\infty} x \chi_L(x_k) \quad (7.3)$$

take the form

$$\mathcal{S}_L u = (u_2, u_3, u_4, \dots), \quad \mathcal{S}_R u = (0, u_1, u_2, \dots)$$

(note that the series in (7.3) converges weakly; in fact it even converges strongly in the sense that the series $\sum_k (x_k \chi_L u)$ converges in \mathcal{H} for every $u \in \mathcal{H}$). These are the usual shift operators, implying that

$$\text{ind } \mathcal{S} = 1.$$

We finally remark that a general index $p \in \mathbb{N}$ can be arranged by modifying (7.1) and (7.2) to

$$\begin{aligned} x_k u &= (\underbrace{0, \dots, 0}_{k-1 \text{ entries}}, -u_{k+p}, \underbrace{0, \dots, 0}_{p-1 \text{ entries}}, -u_k, 0, \dots) \\ \Gamma(x_k) u &= (\underbrace{0, \dots, 0}, u_k, \underbrace{0, \dots, 0}, -u_{k+p}, 0, \dots) . \end{aligned}$$

Moreover, a negative index can be arranged by exchanging the left- and right-handed components.

8. EXAMPLE: A DIRAC OPERATOR WITH $\text{ind}_0 \mathcal{S} \neq 0$

We now construct a two-dimensional space-time (\mathcal{M}, g) together with an odd Dirac operator \mathcal{D} such that the resulting fermionic signature operator in the massless case has a non-trivial chiral index ind_0 (see Definition 5.1). We choose $\mathcal{M} = (0, 2\pi) \times S^1$ with coordinates $t \in (0, 2\pi)$ and $\varphi \in [0, 2\pi)$. We begin with the flat Lorentzian metric

$$ds^2 = dt^2 - d\varphi^2 . \quad (8.1)$$

We consider two-component complex spinors, with the spin scalar product

$$\langle \psi | \phi \rangle = \langle \psi | \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \phi \rangle_{\mathbb{C}^2} . \quad (8.2)$$

We choose the pseudoscalar matrix as

$$\Gamma = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} , \quad (8.3)$$

so that

$$\chi_L = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} , \quad \chi_R = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} . \quad (8.4)$$

The space-time inner product (2.4) becomes

$$\langle \psi | \phi \rangle = \int_0^{2\pi} \int_0^{2\pi} \langle \psi(t, \varphi) | \phi(t, \varphi) \rangle_{\mathbb{C}^2} d\varphi dt . \quad (8.5)$$

The Dirac operator \mathcal{D} should be chosen to be odd (see the right equation in (5.1)). This means that \mathcal{D} has the matrix representation

$$\mathcal{D} = \begin{pmatrix} 0 & \mathcal{D}_R \\ \mathcal{D}_L & 0 \end{pmatrix} \quad (8.6)$$

with suitable operators \mathcal{D}_L and \mathcal{D}_R . In order for current conservation to hold, the Dirac operator should be symmetric with respect to the inner product (8.5). This implies that the operators \mathcal{D}_L and \mathcal{D}_R must both be symmetric,


$$\mathcal{D}_L^* = \mathcal{D}_L , \quad \mathcal{D}_R^* = \mathcal{D}_R , \quad (8.7)$$

where the star denotes the formal adjoint with respect to the scalar product on the Hilbert space $L^2(\mathcal{M}, \mathbb{C})$. We consider the massless Dirac equation

$$\mathcal{D}\psi = 0 . \quad (8.8)$$

The scalar product (2.5) on the solutions takes the form

$$(\psi | \phi) = 2\pi \int_0^{2\pi} \langle \psi(t, \varphi) | \phi(t, \varphi) \rangle_{\mathbb{C}^2} d\varphi , \quad (8.9)$$

FIGURE 1. The eigenvalues $\omega_{k,L/R}$ in the case $p = 2$.

giving rise to the Hilbert space $(\mathcal{H}_0, (\cdot|\cdot))$. As a consequence of current conservation, this scalar product is independent of the choice of t .

We assume that the system is invariant under time translations and is a first order differential operator in time. More precisely, we assume that

$$\mathcal{D}_{L/R} = i\partial_t - H_{L/R} \quad (8.10)$$

with purely spatial operators $H_{L/R}$, referred to as the left- and right-handed Hamiltonians. Moreover, we assume that these Hamiltonians are homogeneous. This implies that they can be diagonalized by plane waves,

$$\mathcal{D}_c e^{ik\varphi} = \omega_{k,c} e^{ik\varphi} \quad \text{with } k \in \mathbb{Z} \text{ and } c \in \{L, R\}.$$

As a consequence, the Dirac equation (8.8) can be solved by the plane waves

$$\mathbf{e}_{k,L} = \frac{1}{2\pi} e^{-i\omega_{k,L}t + ik\varphi} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \mathbf{e}_{k,R} = \frac{1}{2\pi} e^{-i\omega_{k,R}t + ik\varphi} \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (8.11)$$

The vectors $(\mathbf{e}_{k,c})_{k \in \mathbb{Z}, c \in \{L,R\}}$ form an orthonormal basis of the Hilbert space \mathcal{H}_0 . We remark that the Dirac operator of the Minkowski vacuum is obtained by choosing

$$H_L = i\partial_\varphi, \quad H_R = -i\partial_\varphi$$

(see for example [5] or [4, Section 7.2]). In this case, $\omega_{k,L/R} = \mp k$. More generally, choosing \mathcal{D}_c as a homogeneous differential operator of first order, the eigenvalues $\omega_{k,c}$ are linear in k . Here we do not want to assume that the operators \mathcal{D}_c are differential operators. Then the eigenvalues $\omega_{k,L}$ and $\omega_{k,R}$ can be chosen arbitrarily and independently, except for the constraint coming from the symmetry (8.7) that these eigenvalues must be real.

More specifically, for a given parameter $p \in \mathbb{N}$ we choose

$$\omega_{k,L} = -k \quad \text{and} \quad \omega_{k,R} = \begin{cases} k & \text{if } k \leq 0 \\ k + p & \text{if } k > 0 \end{cases} \quad (8.12)$$

(see Figure 1). Then the space-time inner product of the basis vectors $(\mathbf{e}_{k,c})_{k \in \mathbb{Z}, c \in \{L,R\}}$ is computed by

$$\begin{aligned} \langle \mathbf{e}_{k,L} | \mathbf{e}_{k',L} \rangle &= 0 = \langle \mathbf{e}_{k,R} | \mathbf{e}_{k',R} \rangle \\ \langle \mathbf{e}_{k,R} | \mathbf{e}_{k',L} \rangle &= \frac{1}{2\pi} \delta_{k,k'} \int_0^{2\pi} e^{i(\omega_{k,R} - \omega_{k,L})t} dt = \delta_{k,k'} \delta_{\omega_{k,R}, \omega_{k,L}} = \delta_{k,0} \delta_{k',0}. \end{aligned}$$

We conclude \mathcal{S} does not have finite chiral index.

In order to obtain a non-trivial index, we need to modify our example. The idea is to change the space-time inner product in such a way that the inner product between two different plane-wave solutions with the same frequencies becomes non-zero. As a consequence, the corresponding pair of plane-wave solutions will disappear from the kernel. The only vectors which remain in the kernel are those which do not have a partner for pairing, so that

$$\ker \mathcal{S}_L|_{\mathcal{H}_L} = \text{span}(\mathbf{e}_{-1,L}, \dots, \mathbf{e}_{-p,L}), \quad \ker \mathcal{S}_R|_{\mathcal{H}_R} = \{0\}.$$

(see again Figure 1, where the pairs are indicated by horizontal dashed lines, whereas the vectors in the kernel correspond to the circled dots). Generally speaking, the method to modify the space-time inner product for states with the same frequency is to insert a potential into the Dirac equation which is time-independent but has a non-trivial spatial dependence. It is most convenient to work with a *conformal transformation*. Thus we go over from the Minkowski metric (8.1) to the conformally flat metric

$$d\tilde{s}^2 = f(\varphi)^2 (dt^2 - d\varphi^2), \quad (8.13)$$

where $f \in C^\infty(\mathbb{R}/(2\pi\mathbb{Z}))$ is a non-negative, smooth, 2π -periodic function. The conformal invariance of the Dirac equation (for details see for example [4, Section 8.1] and the references therein) states in our situation that the Dirac operator transforms as

$$\tilde{\mathcal{D}} = f^{-\frac{3}{2}} \mathcal{D} f^{\frac{1}{2}}, \quad (8.14)$$

so that

$$\tilde{\mathcal{D}} = \begin{pmatrix} 0 & \tilde{\mathcal{D}}_R \\ \tilde{\mathcal{D}}_L & 0 \end{pmatrix} \quad \text{with} \quad \tilde{\mathcal{D}}_{L/R} = f^{-\frac{3}{2}} \mathcal{D}_{L/R} f^{\frac{1}{2}}.$$

The solutions of the massless Dirac equation are modified simply by a conformal factor,

$$\tilde{\psi} = f^{-\frac{1}{2}} \psi. \quad (8.15)$$

The space-time inner product (8.5) and the scalar product (8.9) transform to

$$\begin{aligned} \langle \tilde{\psi} | \tilde{\phi} \rangle &= \int_0^{2\pi} \int_0^{2\pi} \langle \tilde{\psi}(t, \varphi) | \tilde{\phi}(t, \varphi) \rangle f(\varphi)^2 d\varphi dt \\ &= \int_0^{2\pi} \int_0^{2\pi} \langle \psi(t, \varphi) | \phi(t, \varphi) \rangle f(\varphi) d\varphi dt \end{aligned} \quad (8.16)$$

$$\begin{aligned} (\tilde{\psi} | \tilde{\phi}) &= 2\pi \int_0^{2\pi} \langle \tilde{\psi}(t, \varphi) | \tilde{\phi}(t, \varphi) \rangle_{\mathbb{C}^2} f(\varphi) d\varphi \\ &= 2\pi \int_0^{2\pi} \langle \psi(t, \varphi) | \phi(t, \varphi) \rangle_{\mathbb{C}^2} d\varphi = (\psi | \phi). \end{aligned} \quad (8.17)$$

To understand these transformation laws, one should keep in mind that the spin scalar product remains unchanged under conformal transformations. The same is true for the integrand $\langle \psi | \psi \phi \rangle_x$ of the scalar product (2.5), because the operator ψ is normalized by $\psi^2 = \mathbf{1}$.

From (8.17) we conclude that the scalar product does not change under conformal transformations. In particular, the conformally transformed plane-wave solutions

$$\tilde{\mathbf{e}}_{k,L/R} = f(\varphi)^{-\frac{1}{2}} \mathbf{e}_{k,L/R} \quad (8.18)$$

are an orthonormal basis of $\tilde{\mathcal{H}}_0$. The space-time inner product (8.16), however, involves a conformal factor $f(\varphi)$. As a consequence, the space-time inner product of the basis vectors $(\tilde{\mathbf{e}}_{k,c})_{k \in \mathbb{Z}, c \in \{L,R\}}$ can be computed by

$$\begin{aligned} \langle \tilde{\mathbf{e}}_{k,L} | \tilde{\mathbf{e}}_{k',L} \rangle &= 0 = \langle \tilde{\mathbf{e}}_{k,R} | \tilde{\mathbf{e}}_{k',R} \rangle \\ \langle \tilde{\mathbf{e}}_{k,R} | \tilde{\mathbf{e}}_{k',L} \rangle &= \int_0^{2\pi} dt \int_0^{2\pi} f(\varphi) d\varphi \langle \tilde{\mathbf{e}}_{k,R}(t, \varphi) | \tilde{\mathbf{e}}_{k',L}(t, \varphi) \rangle \\ &= \frac{1}{2\pi} \delta_{\omega_{k,R}, \omega_{k',L}} \int_0^{2\pi} f(\varphi) e^{-i(k-k')\varphi} d\varphi = \frac{1}{2\pi} \delta_{\omega_{k,R}, \omega_{k',L}} \hat{f}_{k-k'}, \end{aligned}$$

where \hat{f}_k is the k^{th} Fourier coefficient of f ,

$$f(\varphi) = \frac{1}{2\pi} \sum_{k \in \mathbb{Z}} \hat{f}_k e^{ik\varphi}.$$

Using the explicit form of the frequencies (8.12), we obtain the following invariant subspaces and corresponding matrix representations of \mathcal{S} ,

$$\begin{aligned} \hat{\mathcal{S}}|_{\text{span}(\tilde{\mathbf{e}}_{-k,L}, \tilde{\mathbf{e}}_{k,R})} &= \frac{1}{2\pi} \begin{pmatrix} 0 & \overline{\hat{f}_{2k}} \\ \hat{f}_{2k} & 0 \end{pmatrix} && \text{if } k \leq 0 \\ \hat{\mathcal{S}}|_{\text{span}(\tilde{\mathbf{e}}_{-k-p,L}, \tilde{\mathbf{e}}_{k,R})} &= \frac{1}{2\pi} \begin{pmatrix} 0 & \overline{\hat{f}_{2k+p}} \\ \hat{f}_{2k+p} & 0 \end{pmatrix} && \text{if } k > p \\ \hat{\mathcal{S}}|_{\text{span}(\tilde{\mathbf{e}}_{-1,L}, \tilde{\mathbf{e}}_{-p,L})} &= 0. \end{aligned}$$

In particular, we can read off the chiral index:

Proposition 8.1. *Assume that almost all Fourier coefficients \hat{f}_k of the conformal function in (8.13) are non-zero. Then the fermionic signature operator in the massless odd case has finite chiral index (see Definition 5.1) and $\text{ind}_0 \mathcal{S} = p$.*

We finally compute the Dirac operator in position space. The dispersion relations in (8.12) are realized by the operators

$$\begin{aligned} \mathcal{D}_L &= i(\partial_t - \partial_\varphi) \\ \mathcal{D}_R &= i(\partial_t + \partial_\varphi) + \mathcal{B}, \end{aligned}$$

where \mathcal{B} is the spatial integral operator

$$(\mathcal{B}\psi)(t, \varphi) = \int_0^{2\pi} \mathcal{B}(\varphi, \varphi') \psi(t, \varphi') d\varphi'$$

with the distributional integral kernel

$$\mathcal{B}(\varphi, \varphi') = -\frac{p}{2\pi} \sum_{k=1}^{\infty} e^{ik(\varphi-\varphi')} = -\frac{p}{2} \delta(\varphi - \varphi') - \frac{p}{2\pi} \frac{\text{PP}}{e^{-i(\varphi-\varphi')} - 1}.$$

Hence, choosing the Dirac matrices as

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (8.19)$$

and using (8.4), we obtain

$$\mathcal{D} = i\gamma^0 \partial_t + i\gamma^1 \partial_\varphi + \gamma^1 \chi_R \mathcal{B}. \quad (8.20)$$

Performing the conformal transformation (8.14), we finally obtain

$$(\tilde{\mathcal{D}}\psi)(t, \varphi) = \frac{i}{f(\varphi)} \left(\gamma^0 \partial_t + \gamma^1 \partial_\varphi + \frac{f'(\varphi)}{2f(\varphi)} - \frac{p}{2} \gamma^1 \chi_R \right) \psi(t, \varphi) \quad (8.21)$$

$$- \frac{p}{2\pi} \frac{\gamma^1 \chi_R}{f(\varphi)^{\frac{3}{2}}} \int_0^{2\pi} \frac{\text{PP}}{e^{-i(\varphi-\varphi')} - 1} \psi(t, \varphi') \sqrt{f(\varphi')} d\varphi'. \quad (8.22)$$

Thus (8.21) is the Dirac operator in the Lorentzian metric (8.13) with a constant right-handed potential. Moreover, the summand (8.22) is a nonlocal integral operator involving a singular integral kernel.

This example shows that the index of Proposition 8.1 in general does not encode the topology of space-time, because for a fixed space-time topology the index can take any integer value. The way we understand the index is that it gives topological information on the singular behavior of the potential in the Dirac operator.

9. EXAMPLE: A DIRAC OPERATOR WITH $\text{ind } \mathcal{S} \neq 0$

We now construct an example of a fermionic signature operator for which the index $\text{ind } \mathcal{S}$ of Definition 3.3 is non-trivial. To this end, we want to modify the example of the previous section. The major difference to the previous setting is that the Hilbert space \mathcal{H}_m does not have a decomposition into two subspaces \mathcal{H}_L and \mathcal{H}_R , making it necessary to consider the operators \mathcal{S}_L and \mathcal{S}_R as operators on the whole solution space \mathcal{H}_m . Our first task is to remove the infinite-dimensional kernels of the operators \mathcal{S}_L and \mathcal{S}_R . This can typically be achieved by perturbing the Dirac operator, for example by introducing a rest mass. The second and more substantial modification is to arrange that the operators \mathcal{S}_L and \mathcal{S}_R have *infinite-dimensional invariant subspaces*. This is needed for the following reason: In the example of the previous section, the operator $\mathcal{S}_L|_{\mathcal{H}_L} : \mathcal{H}_L \rightarrow \mathcal{H}_R$ mapped one Hilbert space to another Hilbert space. Therefore, we obtained a non-trivial index simply by arranging that the operator $\mathcal{S}_L|_{\mathcal{H}_L}$ gives a non-trivial “pairing” of vectors of \mathcal{H}_L with vectors of \mathcal{H}_R (as indicated in Figure 1 by the horizontal dashed lines). In particular, if considered as an operator on \mathcal{H}_0 , the operator \mathcal{S}_L had at most two-dimensional invariant subspaces. For the chiral index of Definition 3.3, however, we have only one Hilbert space \mathcal{H}_m to our disposal, so that the operator $\mathcal{S}_L : \mathcal{H}_m \rightarrow \mathcal{H}_m$ is an endomorphism of \mathcal{H}_m . As a consequence, the chiral index is trivial whenever \mathcal{H}_m splits into a direct sum of finite-dimensional subspaces which are invariant under \mathcal{S}_L (because on each invariant subspace, the index is trivial due to the rank-nullity theorem of linear algebra).

The following example is designed with the aim of showing in explicit detail that the index is non-zero. Our starting point are the plane-wave solutions (8.11) with the frequencies according to (8.12) with $p \in \mathbb{N}$. In Figure 2 the transformed plane-wave solutions $\tilde{\epsilon}_{k,c}$ (where the transformation from $\epsilon_{k,c}$ to $\tilde{\epsilon}_{k,c}$ will be explained below) are arranged according to their frequencies and momenta on a lattice. We shall construct the operator \mathcal{S}_L in such a way that these plane-wave solutions are mapped to each other as indicated by the arrows. Thus similar to a shift operator, \mathcal{S}_L maps the basis vectors to each other “spiraling in,” implying that the vector $\tilde{\epsilon}_{-1,L}$ (depicted with the circled dot) is in the kernel of \mathcal{S}_L . Likewise, the operator \mathcal{S}_R acts like a “spiraling out” shift vector, so that it is injective. In this way, we arrange that $\text{ind } \mathcal{S} = 1$. Similarly, in the case $p > 1$ we shall obtain p spirals, so that $\text{ind } \mathcal{S} = p$.




FIGURE 2. The action of S_L on the transformed plane-wave solutions in the case $p = 1$.

Before entering the detailed construction, we point out that our method is driven by the wish that the example should be explicit and that the kernels of the chiral signature operators should be given in closed form. This makes it necessary to introduce a Dirac operator which seems somewhat artificial. In particular, instead of introducing a rest mass, we arrange a mixing of the left- and right-handed components using a time-dependent vectorial gauge transformation. Moreover, we again work with a conformal transformation with a carefully adjusted spatial and time dependence. We consider these special features merely as a requirement needed in order to make the computations as simple as possible. In view of the stability result of Theorem 6.2, we expect that the index is also non-trivial in more realistic examples involving a rest mass and less fine-tuned potentials. But probably, this goes at the expense of longer computations or less explicit arguments.

We begin on the cylinder $\mathcal{M} = (0, 6\pi) \times S^1$, again with the Minkowski metric (8.1) and two-component spinors endowed with the spin scalar product (8.2). The space-time inner product (2.4) becomes


$$\langle \psi | \phi \rangle = \int_0^{6\pi} \int_0^{2\pi} \langle \psi(t, \varphi) | \phi(t, \varphi) \rangle d\varphi dt, \quad (9.1)$$

whereas the scalar product on solutions of the Dirac equation is again given by (8.9). We again consider the massless Dirac equation (8.8) with the Dirac operator (8.6) and the left- and right-handed operators according to (8.10). Moreover, we again assume that the operators $\mathcal{D}_{L/R}$ have the plane-wave solutions (8.11) with frequencies (8.12). For a fixed real parameter $\nu \neq 0$, we consider the transformation

$$U(t) = \frac{\mathbb{1} + i\nu\gamma^0 \cos t/3}{\sqrt{1 + \nu^2 \cos^2 t/3}} = \frac{1}{\sqrt{1 + \nu^2 \cos^2 t/3}} \begin{pmatrix} 1 & i\nu \cos t/3 \\ i\nu \cos t/3 & 1 \end{pmatrix}. \quad (9.2)$$

Obviously, $U(t) \in U(2)$ is a unitary matrix. Moreover, it commutes with γ^0 , implying that it is also unitary with respect to the spin scalar product. As a consequence, the transformation $U(t)$ is unitary both on the Hilbert space \mathcal{H}_0 and with respect to the inner product (9.1). Next, we again consider a conformal transformation (8.14) and (8.15), but now with a conformal function $f(t, \varphi)$ which depends on space and time. Thus we set

$$\tilde{\mathcal{D}} = f^{-\frac{3}{2}} U \mathcal{D} U^* f^{\frac{1}{2}} \quad \text{and} \quad \tilde{\psi} = f^{-\frac{1}{2}} U \psi. \quad (9.3)$$

FIGURE 3. The transformation V in momentum space.

Similar to (8.16) and (8.17), the inner products transform to

$$\langle \tilde{\psi} | \tilde{\phi} \rangle = \int_0^{6\pi} \int_0^{2\pi} \langle \psi(t, \varphi) | \phi(t, \varphi) \rangle f(t, \varphi) d\varphi dt \quad \text{and} \quad (\tilde{\psi} | \tilde{\phi}) = (\psi | \phi).$$

In particular, the transformed plane wave solutions $\tilde{e}_{k,c}$ are an orthonormal basis of \mathcal{H}_0 . Keeping in mind that the chiral projectors in (3.1) do *not* commute with U , we obtain

$$\langle \tilde{\psi} | \tilde{\phi} \rangle_L = \int_0^{6\pi} \int_0^{2\pi} \langle U(t) \psi(t, \varphi) | \chi_L U(t) \phi(t, \varphi) \rangle f(t, \varphi) d\varphi dt$$

and thus, in view of (3.2),

$$(\tilde{\mathbf{e}}_{k,c} | \mathcal{S}_L \tilde{\mathbf{e}}_{k',c'}) = \int_0^{6\pi} \int_0^{2\pi} \langle U \mathbf{e}_{k,c} | \chi_L U \mathbf{e}_{k',c'} \rangle f(t, \varphi) d\varphi dt. \quad (9.4)$$

In order to get rid of the square roots in (9.2), it is most convenient to set

$$V(t) = \begin{pmatrix} 1 & i\nu \cos t/3 \\ i\nu \cos t/3 & 1 \end{pmatrix} \quad \text{and} \quad \mu(t, \varphi) = \frac{f(t, \varphi)}{1 + \nu^2 \cos^2 t/3}. \quad (9.5)$$

Then (9.4) simplifies to

$$(\tilde{\mathbf{e}}_{k,c} | \mathcal{S}_L \tilde{\mathbf{e}}_{k',c'}) = \int_0^{6\pi} \int_0^{2\pi} \langle V \mathbf{e}_{k,c} | \chi_L V \mathbf{e}_{k',c'} \rangle \mu(t, \varphi) d\varphi dt. \quad (9.6)$$

Let us first discuss the effect of the transformation V . A left-handed spinor is mapped to

$$V \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \frac{i}{2} e^{it/3} \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \frac{i}{2} e^{-it/3} \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Thus two right-handed contributions are generated, whose frequency differ from the frequency of the left-handed component by $\pm 1/3$. Similarly, a right-handed spinor is mapped to

$$V \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \frac{i}{2} e^{it/3} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \frac{i}{2} e^{-it/3} \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

generating two left-handed components with frequencies shifted by $\pm 1/3$. Again plotting the frequencies vertically, we depict the transformation V as in Figure 3. The same notation is also used in Figure 2 for the transformed plane-wave solutions.

The inner product $\langle \cdot | \chi_L \cdot \rangle$ in (9.6) only gives a contribution if the arguments on the left and right have the opposite chirality. Since the transformed plane-wave solutions $V e_{k,c}$ have a fixed chirality at every lattice point, one sees in particular that (9.6) vanishes if μ is chosen as a constant. By adding to the constant $\mu = 1$ contributions with different momenta, we can connect the different lattice points in Figure 2. This leads us to the ansatz

$$\mu(t, \varphi) = 1 + \mu_{\text{hor}}(t, \varphi) + \mu_{\text{vert}}(t, \varphi), \quad (9.7)$$

where the last two summands should describe the horizontal respectively vertical arrows in Figure 2. For the horizontal arrows we can work similar to (8.18) with a spatially-dependent conformal transformation. However, in order to make sure that the left-handed component generated by V (corresponding to the two L s at the very right of Figure (2)) are not connected horizontally, we include two Fourier modes which shift the frequency by $\pm 2/3$,

$$\mu_{\text{hor}}(t, \varphi) = a(\varphi) \left(1 - e^{\frac{2it}{3}} - e^{-\frac{2it}{3}} \right), \quad (9.8)$$

where a has the Fourier decomposition

$$a(\varphi) = \sum_{k=1}^{\infty} \left(a_k e^{ik\varphi} + \overline{a_{-k}} e^{-ik\varphi} \right). \quad (9.9)$$

For the vertical arrows we must be careful that the left-handed contribution of $V \mathbf{e}_{k,L}$ is not connected to the right-handed component of $V \mathbf{e}_{k,R}$, because then the arrow would have the wrong direction. To this end, we avoid integer frequencies. Instead, we work with the frequencies in $\mathbb{Z} \pm 1/3$, because they connect the left-handed component $V \mathbf{e}_{k,R}$ to the right-handed component of $V \mathbf{e}_{k,L}$. This leads us to the ansatz

$$\mu_{\text{vert}}(t, \varphi) = \mu_{\text{vert}}(t) = \sum_{n \in \mathbb{Z}} e^{int} \left(b_n e^{\frac{it}{3}} + \overline{b_{-n}} e^{-\frac{it}{3}} \right). \quad (9.10)$$

The ansätze (9.9) and (9.10) ensure that μ is real-valued. Moreover, by choosing the Fourier coefficients sufficiently small, one can clearly arrange that the first summand in (9.7) dominates, so that μ is strictly positive. We thus obtain the following result.

Proposition 9.1. *Assume that the Fourier coefficients a_k and b_n in (9.9) and (9.10) are sufficiently small and that almost all Fourier coefficients are non-zero. Then the function μ defined by (9.8) and (9.7), is strictly positive. Consider the Dirac operator (9.3) with U and f according to (9.2) (for some fixed $\nu \in \mathbb{R} \setminus \{0\}$) and (9.5). Then the chiral index of the fermionic signature operator (see Definition 3.3) is finite and $\text{ind } \mathcal{S} = p$.*

We finally discuss the form of the Dirac operator in position space. Substituting (8.20) into (9.3) and using the above form of U and f , the Dirac operator \tilde{D} can be computed in closed form. Similar as discussed in the previous section, the Dirac operator contains a nonlocal integral operator with a singular potential. Moreover, the transformation U modifies the Dirac matrix γ^1 to

$$\gamma^1 \rightarrow U \gamma^1 U^* = \frac{1}{1 + \nu^2 \cos^2(t/3)} \left((1 - \nu^2 \cos^2(t/3)) \gamma^1 - 2\nu \cos^2(t/3) \gamma^2 \right),$$

where

$$\gamma^2 := \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}.$$

Thus the representation of the Dirac matrices becomes time-dependent; this is the main effect of the vectorial transformation U . This transformation changes the first order terms in the Dirac equation. Moreover, the conformal transformation also changes the first order terms just as in (8.21) by a prefactor $1/f$.

10. EXAMPLES ILLUSTRATING THE HOMOTOPY INVARIANCE

We now give two examples to illustrate our considerations on the homotopy invariance of the chiral index. We begin with an example which shows that the dimension of the kernel of \mathcal{S}_L does not need to be constant for deformations which are continuous in $L(\mathcal{H}_0)$. It may even become infinite-dimensional.

Example 10.1. We consider the space-time $\mathcal{M} = (0, T) \times S^1$ with coordinates $t \in (0, T)$ and $\varphi \in [0, 2\pi)$ endowed with the Minkowski metric

$$ds^2 = dt^2 - d\varphi^2.$$

We again choose two-component complex spinors with the spin scalar product (8.2). The Dirac operator is chosen as

$$\mathcal{D} = i\gamma^0 \partial_t + i\gamma^1 \partial_\varphi,$$

where the Dirac matrices are again given by (8.19). The pseudoscalar matrix and the chiral projectors are again chosen according to (8.3) and (8.4).

We consider the massless Dirac equation

$$\mathcal{D}\psi = 0.$$

This equation (8.8) can be solved by plane wave solutions, which we write as

$$\mathbf{e}_{k,L}(\zeta) = \frac{1}{2\pi} e^{+ikt+ik\varphi} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \mathbf{e}_{k,R}(\zeta) = \frac{1}{2\pi} e^{-ikt+ik\varphi} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (10.1)$$

where $k \in \mathbb{Z}$ (the indices L and R denote the left- and right-handed components; at the same time they propagate to the left respectively right). By direct computation, one verifies that $(\mathbf{e}_{k,c})_{k \in \mathbb{Z}, c \in \{L,R\}}$ is an orthonormal basis of \mathcal{H}_0 .

We next compute the space-time inner product (2.4),

$$\begin{aligned} \langle \mathbf{e}_{k,R} | \mathbf{e}_{0,L} \rangle &= \int_0^T dt \int_0^{2\pi} d\varphi \langle \mathbf{e}_{k,R}(t, \varphi) | \mathbf{e}_{0,L}(t, \varphi) \rangle = \frac{1}{2\pi} \int_0^T \delta_{k,0} dt = \frac{T}{2\pi} \delta_{k,0} \\ \langle \mathbf{e}_{k,R} | \mathbf{e}_{k',L} \rangle &= \frac{1}{2\pi} \delta_{k,k'} \int_0^T e^{2ikt} dt = \frac{e^{2ikT} - 1}{4\pi ik} \quad (k' \neq 0) \\ \langle \mathbf{e}_{k,L} | \mathbf{e}_{k',L} \rangle &= 0 = \langle \mathbf{e}_{k,R} | \mathbf{e}_{k',R} \rangle. \end{aligned}$$

Thus the fermionic signature operator \mathcal{S} is invariant on the subspaces $\mathcal{H}_0^{(k)}$ generated by the basis vectors $\mathbf{e}_{k,L}$ and $\mathbf{e}_{k,R}$. Moreover, in these bases it has the matrix representations

$$\mathcal{S}|_{\mathcal{H}_0^{(0)}} = \frac{T}{2\pi} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{S}|_{\mathcal{H}_0^{(k)}} = \frac{1}{4\pi ik} \begin{pmatrix} 0 & e^{2ikT} - 1 \\ e^{-2ikT} - 1 & 0 \end{pmatrix} \quad (k \neq 0).$$

If $T \notin \pi\mathbb{Q}$, the matrix entries $e^{\pm 2ikT} - 1$ are all non-zero. As a consequence, the operators $\mathcal{S}_L|_{\mathcal{H}_L}$ and $\mathcal{S}_R|_{\mathcal{H}_R}$ are both injective. Thus \mathcal{S} has finite chiral index in the massless odd case (see Definition 5.1). If $T \in \pi\mathbb{Q}$, however, the chiral index vanishes for all k for which $2kT$ is a multiple of 2π . As a consequence, the operators $\mathcal{S}_L|_{\mathcal{H}_L}$ and $\mathcal{S}_R|_{\mathcal{H}_R}$ both have an infinite-dimensional kernel, so that \mathcal{S} does not have a finite chiral index. \diamond

This example also explains why we need additional assumptions like those in Theorems 6.2 and 6.3. In particular, when considering homotopies of space-time or of the

Dirac operator, one must be careful to ensure that the chiral index remains finite along the chosen path.

We next want to construct examples of homotopies to which the stability result of Theorem 6.3 applies. To this end, it is convenient to work similar to (8.13) with a conformal transformation.

Example 10.2. As in Example 10.1 we consider the space-time $(0, T) \times S^1$, but now with the conformally transformed metric

$$d\tilde{s}^2 = f(t)^2 (dt^2 - d\varphi^2)$$

where f is a non-negative C^2 -function with

$$\text{supp } f \subset (-T, T) \quad \text{and} \quad f(0) > 0.$$

Similar to (8.18) and the computation thereafter, transforming the plane-wave solutions (10.1) conformally to $\tilde{\mathbf{e}}_{k,L/R} = f(t)^{-\frac{1}{2}} \mathbf{e}_{k,L/R}$, we again obtain an orthonormal basis of \mathcal{H}_0 and

$$\begin{aligned} \langle \tilde{\mathbf{e}}_{k,R} | \tilde{\mathbf{e}}_{k',L} \rangle &= \frac{1}{2\pi} \delta_{k,k'} \int_0^T f(t) e^{2ikt} dt \\ \langle \tilde{\mathbf{e}}_{k,L} | \tilde{\mathbf{e}}_{k',L} \rangle &= 0 = \langle \tilde{\mathbf{e}}_{k,R} | \tilde{\mathbf{e}}_{k',R} \rangle \end{aligned}$$

for all $k, k' \in \mathbb{Z}$.

The integration-by-parts argument

$$\begin{aligned} \int_0^T f(t) e^{2ikt} dt &= \frac{1}{2ik} \int_0^T f(t) \frac{d}{dt} e^{2ikt} dt = -\frac{f(0)}{2ik} - \frac{1}{2ik} \int_0^T f'(t) e^{2ikt} dt \\ &= -\frac{f(0)}{2ik} - \frac{f'(0)}{4k^2} - \frac{1}{4k^2} \int_0^T f''(t) e^{2ikt} dt \end{aligned}$$

shows that the space-time inner products have a simple explicit asymptotics for large k given by

$$\langle \tilde{\mathbf{e}}_{k,R} | \tilde{\mathbf{e}}_{k',L} \rangle = \frac{f(0)}{4\pi ik} \delta_{k,k'} + \mathcal{O}\left(\frac{1}{k^2}\right).$$

Hence the operator \mathcal{S}_L has the form

$$\mathcal{S}_L \mathbf{e}_{k,L} = c_k \mathbf{e}_{k,R}$$

with coefficients c_k having the asymptotics

$$c_k = \frac{f(0)}{4\pi ik} + \mathcal{O}\left(\frac{1}{k^2}\right).$$

From this asymptotics we can read off the following facts. First, it is obvious that $\mathcal{S}_L|_{\mathcal{H}_L}$ has a finite-dimensional kernel. Exchanging the chirality, the same is true for $\mathcal{S}_R|_{\mathcal{H}_R}$, implying that S has a finite chiral index (according to Definition 5.1). Next, the vectors in the image of \mathcal{S}_L are in the Sobolev space $W^{1,2}$,

$$\mathcal{S}_L|_{\mathcal{H}_L} : \mathcal{H}_L \rightarrow \mathcal{H}_R \cap W^{1,2}(S^1, \mathbb{C}^2).$$

Moreover, the image of this operator is closed (in the $W^{1,2}$ -norm). Finally, our partial integration argument also yields that

$$\|\mathcal{S}_L \psi\|_{W^{1,2}} \leq |f|_{C^2} \|\psi\|_{\mathcal{H}_0},$$

showing that the family of signature operators is norm continuous for a C^2 -homotopy of functions f .

Having verified the assumptions of Theorem 6.3, we conclude that the chiral index in the massless odd case is invariant under C^2 -homotopies of the conformal function f , provided that $f(0)$ stays away from zero. \diamond

11. CONCLUSION AND OUTLOOK

Our analysis shows that the chiral index of a fermionic signature operator is well-defined and in general non-trivial. Moreover, it is a homotopy invariant provided that the additional conditions stated in Theorems 6.2 and 6.3 are satisfied. As already mentioned at the end of the introduction, the physical and geometric meaning of this index is yet to be explored.

We now outline how our definition of the chiral index could be generalized or extended other situations. First, our constructions also apply in the *Riemannian setting* by working instead of causal fermion systems with so-called Riemannian fermion systems or general *topological fermion systems* as introduced in [4]. In this situation, one again imposes a pseudoscalar operators $\Gamma(x) \in L(\mathcal{H})$ with the properties (4.2). Then all constructions in Section 4 go through. Starting on an even-dimensional Riemannian spin manifold, one can proceed as explained in [4] and first construct a corresponding topological fermion system. For this construction, one must choose a particle space, typically of eigensolutions of the Dirac equation. Once the topological fermion system is constructed, one can again work with the index of Section 4. If the Dirac operator anti-commutes with the pseudoscalar operator, one can choose the particle space \mathcal{H} to be invariant under the action of Γ . This gives a decomposition of the particle space into two chiral subspaces, $\mathcal{H} = \mathcal{H}_L \oplus \mathcal{H}_R$. Just as explained in Section 5, this makes it possible to introduce other indices by restricting the chiral signature operators to \mathcal{H}_L or \mathcal{H}_R . Moreover, one could compose the operators from the left with the projection operators onto the subspaces $\mathcal{H}_{L/R}$ and consider the Noether indices of the resulting operators.

Another generalization concerns space-times of *infinite lifetime*. Using the constructions in [7], in such space-times one can still introduce the fermionic signature operator \mathcal{S}_m provided that the space-time satisfies the so-called mass oscillation property. By inserting chiral projection operators, one can again define chiral signature operators $\mathcal{S}_m^{L/R}$ and define the chiral index as their Noether index. Also the stability results of Theorems 6.2 and 6.3 again apply. It is unknown whether the resulting indices have a geometric meaning. Since \mathcal{S}_m depends essentially on the asymptotic form of the Dirac solutions near infinity, the corresponding chiral indices should encode information on the metric and the external potential in the asymptotic ends.

We finally remark that the fermionic signature operator could be *localized* by restricting the space-time integrals to a measurable subset $\Omega \subset \mathcal{M}$. For example, one can introduce a chiral signature operator $\mathcal{S}_L(\Omega)$ similar to (3.1) and (3.2) by

$$(\phi | \mathcal{S}_L(\Omega) \psi) = \int_{\Omega} \langle \psi | \chi_L \phi \rangle_x d\mu_{\mathcal{M}} .$$

Likewise, in the setting of causal fermion systems, one can modify (4.3) to

$$\mathcal{S}_L(\Omega) = - \int_{\Omega} x \chi_L d\rho(x) .$$

The corresponding indices should encode information on the behavior of the Dirac solutions in the space-time region Ω .

Acknowledgments: I would like to thank Niky Kamran and Hermann Schulz-Baldes for helpful discussions.

REFERENCES

- [1] H. Baum, *Spinor structures and Dirac operators on pseudo-Riemannian manifolds*, Bull. Polish Acad. Sci. Math. **33** (1985), no. 3-4, 165–171.
- [2] F. Finster and A. Grotz, *A Lorentzian quantum geometry*, arXiv:1107.2026 [math-ph], Adv. Theor. Math. Phys. **16** (2012), no. 4, 1197–1290.
- [3] F. Finster, A. Grotz, and D. Schiefeneder, *Causal fermion systems: A quantum space-time emerging from an action principle*, arXiv:1102.2585 [math-ph], Quantum Field Theory and Gravity (F. Finster, O. Müller, M. Nardmann, J. Tolksdorf, and E. Zeidler, eds.), Birkhäuser Verlag, Basel, 2012, pp. 157–182.
- [4] F. Finster and N. Kamran, *Spinors on singular spaces and the topology of causal fermion systems*, arXiv:1403.7885 [math-ph] (2014).
- [5] F. Finster and O. Müller, *Lorentzian spectral geometry for globally hyperbolic surfaces*, arXiv:1411.3578 [math-ph] (2014).
- [6] F. Finster and M. Reintjes, *A non-perturbative construction of the fermionic projector on globally hyperbolic manifolds I – Space-times of finite lifetime*, arXiv:1301.5420 [math-ph], to appear in Adv. Theor. Math. Phys. (2015).
- [7] ———, *A non-perturbative construction of the fermionic projector on globally hyperbolic manifolds II – Space-times of infinite lifetime*, arXiv:1312.7209 [math-ph] (2013).
- [8] H.B. Lawson, Jr. and M.-L. Michelsohn, *Spin Geometry*, Princeton Mathematical Series, vol. 38, Princeton University Press, Princeton, NJ, 1989.
- [9] P.D. Lax, *Functional Analysis*, Pure and Applied Mathematics (New York), Wiley-Interscience [John Wiley & Sons], New York, 2002.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, D-93040 REGENSBURG, GERMANY
E-mail address: finster@ur.de