

Links not concordant to the Hopf link

BY STEFAN FRIEDL

Fakultät für Mathematik, Universität Regensburg, Germany.

e-mail: sfriedl@gmail.com

AND MARK POWELL

Department of Mathematics, Indiana University, Bloomington, Indiana, U.S.A.

e-mail: macp@indiana.edu

(Received 16 May 2012; revised 1 August 2012)

Abstract

We give new Casson–Gordon style obstructions for a two–component link to be topologically concordant to the Hopf link.

1. Introduction

1.1. Definitions and previous results

An m –component (oriented) link is an embedded locally flat ordered collection of m disjoint (oriented) circles in S^3 . We say that two m –component oriented links $L = L_1 \cup \dots \cup L_m$ and $L' = L'_1 \cup \dots \cup L'_m$ are *concordant* if there exists a collection of m disjoint, embedded, locally flat, oriented annuli A_1, \dots, A_m in $S^3 \times [0, 1]$ such that $\partial A_i = L_i \times \{0\} \cup -L'_i \times \{1\}$, $i = 1, \dots, m$.

Throughout the paper we denote by H the Hopf link equipped with the orientation such that the linking number is 1, i.e. H consists of two generic fibres of the Hopf map

$$h: S^3 \longrightarrow S^2 \\ (z_1, z_2) \longmapsto [z_1, z_2],$$

where $S^3 \subset \mathbb{C}^2$ is the set of $(z_1, z_2) \in \mathbb{C}^2$ with $|z_1|^2 + |z_2|^2 = 1$ and $S^2 \cong \mathbb{C}P^1$.

There is a rich literature on obstructions for two links to be concordant. Roughly speaking the obstructions fall into the following categories:

- (i) obstructions to the components (viewed as knots) being concordant (the landmark papers being the work of Levine [Lev69], Casson and Gordon [CG86] and Cochran, Orr and Teichner [COT03]);
- (ii) Milnor’s $\bar{\mu}$ –invariants (see Milnor [Mil57]), in particular the linking numbers;
- (iii) obstructions from the multivariable Alexander polynomial and the Levine–Tristram signatures (see Murasugi [Mur67], Kawauchi [Kaw77][Kaw78, theorem B], Nakagawa [Nak78] and Tristram [Tri69]);
- (iv) twisted Alexander polynomials, twisted signature invariants and L^2 –signature invariants, where the corresponding representations factor through a nilpotent quotient (e.g. a p –group) or the algebraic closure of the link group (see Cha and Ko

[CK99b, CK06], Cha and Friedl [CF10], Friedl [Fri05], Harvey [Har08], Levine [Lev94] and Smolinsky [Smo89]); furthermore see Cha and Orr [CO09] for a closely related approach;

- (v) twisted Alexander polynomials and twisted signature invariants of boundary links, where the corresponding representations factor through the canonical epimorphism onto the free group (see Levine [Lev07]);
- (vi) by considering a prime power branched cover over one component of a link, one can view the preimages of the other components as a link in a rational homology sphere; using this reduction in the number of components one can reduce problems in link concordance to the study of ‘rational knot concordance’ (see Cochran–Orr [CO90, CO93], Cha–Kim [CK08], Cha–Livingston–Ruberman [CLR08] and Cha–Kim–Ruberman–Strle [CKRS10]);
- (vii) by doing a Dehn filling on one component one can also reduce problems in link concordance to the study of knot concordance in integral or rational homology spheres. This approach, which for example appears in Cha–Ko [CK99a] and the aforementioned [CKRS10], is often referred to as ‘blowing down along one component’.

Davis [Dav06] proved a partial converse to the above obstructions: if L is a 2-component link with linking number one and such that the multivariable Alexander polynomial $\Delta_L \in \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$ equals one, then the link is concordant to the Hopf link. This result generalised earlier results by Freedman ([Fre84, FQ90]) on knot concordance and by Hillman [Hil02].

In this paper we will give new obstructions to a link being concordant to the Hopf link by using signature invariants corresponding to appropriately chosen metabelian representations. This approach is inspired by the classic work of Casson and Gordon on knot concordance [CG86]. Generalising their work to the setting of link concordance requires overcoming several delicate technical issues. One particularly difficult step (Theorem 2.1 in this paper) appeared in an earlier paper by the authors [FP12].

1.2. *The linking pairing for 2-component links with linking number 1*

In the following we let L be a 2-component link of linking number 1. Throughout this paper we write $X = X_L = S^3 \setminus \nu L$, where νL is a tubular neighbourhood of L . The boundary ∂X_L can be written as the union of two tori ∂X_L^1 and ∂X_L^2 which correspond to the two link components. Moreover, for each i , we can decompose $\partial X_L^i \cong S^1 \times D^1 \cup_{S^1 \times S^0} S^1 \times D^1$, where each cross section $S^1 \times \{pt\}$ is a meridian of the link component L_i . We denote this splitting by

$$\partial X_L^i \cong Y_a^i \cup_{S^1 \times S^0} Y_b^i,$$

for $i = 1, 2$. Define $Y_a = Y_a^1 \sqcup Y_a^2$ and $Y_b = Y_b^1 \sqcup Y_b^2$.

Note that we have a canonical isomorphism $H_1(X_L; \mathbb{Z}) \xrightarrow{\cong} \mathbb{Z}^2$ which we will use to identify the groups. Now let $\varphi: H_1(X_L; \mathbb{Z}) = \mathbb{Z}^2 \rightarrow A$ be an epimorphism onto a finite abelian group. In the following we denote by $p: X_L^\varphi \rightarrow X$ the corresponding covering space and we write $Y_a^\varphi = p^{-1}(Y_a)$ and $Y_b^\varphi = p^{-1}(Y_b)$. In Section 2.2 we will see that there exists a non-singular linking form

$$\lambda: TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \times TH_1(X^\varphi, Y_b^\varphi; \mathbb{Z}) \longrightarrow \mathbb{Q}/\mathbb{Z},$$

where, given an abelian group G , we write $TG := \text{Tor}_{\mathbb{Z}}(G)$.

Now suppose that L is concordant to the Hopf link via a concordance $C \cong S^1 \times I \subset S^3 \times I$. Denote the exterior of the concordance by E_C . The inclusion maps $X_L \rightarrow E_C$ and

$X_H \rightarrow E_C$ are \mathbb{Z} -homology equivalences. Note that φ canonically extends to a homomorphism $H_1(E_C; \mathbb{Z}) \rightarrow A$ which we also denote by φ . In the following we denote by $p: E_C^\varphi \rightarrow E_C$ the corresponding covering space.

Before we state our first result we need to introduce one more definition. We say that a homomorphism $\varphi: \mathbb{Z}^2 \rightarrow A$ to a finite abelian group is *admissible* if φ is given by the direct sum of two projections on each summand of \mathbb{Z}^2 , i.e. φ has to be of the form $\mathbb{Z}^2 = \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z}_k \oplus \mathbb{Z}_l$. We now have the following proposition.

PROPOSITION 1.1. *Let L be an oriented 2-component link with linking number 1 which is concordant to the Hopf link via a concordance C . Then for any admissible epimorphism $\varphi: \mathbb{Z}^2 \rightarrow A$ to a group of prime power order,*

$$P := \ker(TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \rightarrow TH_1(E_C^\varphi, Y_a^\varphi; \mathbb{Z})),$$

is a metaboliser of λ , i.e. $P = P^\perp$.

Following the philosophy of [CG86], we will use this proposition to show that certain metabelian representations will extend over a concordance exterior.

1.3. *Definition of the invariant $\tau(L, \chi)$*

We define the closed manifold

$$[M_L := X_L \cup_{\partial X_L = \partial X_H} X_H :]$$

an ordering and an orientation of the link components of L and H determines a canonical identification of ∂X_L with ∂X_H which identifies the corresponding meridians and longitudes. We say that a homomorphism $\varphi: H_1(M_L; \mathbb{Z}) \rightarrow A$ is admissible if it restricts to an admissible homomorphism on

$$H_1(X_L; \mathbb{Z}) \subseteq H_1(M_L; \mathbb{Z}) \cong H_1(X_L; \mathbb{Z}) \oplus \mathbb{Z}.$$

We denote the corresponding finite cover by M_L^φ . Note that M_H^φ is a 3-torus. We denote the 3-torus $S^1 \times S^1 \times S^1$ by T . Let

$$\phi': \pi_1(M_L) \rightarrow \mathcal{H}' := H_1(M_L; \mathbb{Z}) \cong \mathbb{Z}^3$$

be the Hurewicz map. Let

$$\phi: \pi_1(M_L^\varphi) \subseteq \pi_1(M_L) \rightarrow \mathcal{H}'$$

be the restriction of ϕ' , and define $\mathcal{H} := \text{im } \phi$. Note that also $\mathcal{H} \cong \mathbb{Z}^3$. We pick an isomorphism

$$\psi: \pi_1(T) \xrightarrow{\cong} H_1(T; \mathbb{Z}) \xrightarrow{\cong} \mathcal{H}.$$

Finally, let

$$\chi: H_1(M_L^\varphi; \mathbb{Z}) \rightarrow S^1 \subset \mathbb{C}$$

be a character which factors through \mathbb{Z}_{q^i} for some prime q , and denote the trivial character by tr .

Following Casson–Gordon, in Section 3.2, we define a Witt group obstruction

$$\tau(L, \chi) \in L^0(\mathbb{C}(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q],$$

in terms of the twisted intersection form of a 4-manifold whose boundary is a finite number of copies of $(M_L^\varphi \sqcup -T, \chi \times \phi \sqcup \text{tr} \times \psi)$.

1.4. *Statement of the main theorem and examples*

The following is our main theorem.

THEOREM 1.2. *Suppose that L is concordant to the Hopf link. Then for any admissible homomorphism $\varphi: H_1(M_L) \rightarrow A$ to a finite abelian p -group A , there exists a metaboliser $P = P^\perp$ for the linking form*

$$TH_1(X_L^\varphi, Y_a^\varphi) \times TH_1(X_L^\varphi, Y_a^\varphi) \longrightarrow \mathbb{Q}/\mathbb{Z}$$

with the following property: for any character of prime power order $\chi: H_1(M_L^\varphi) \rightarrow \mathbb{Z}_{q^k} \rightarrow S^1$, which satisfies that $\chi|_{H_1(X_L^\varphi)}$ factors through

$$\chi|_{H_1(X_L^\varphi)}: H_1(X_L^\varphi) \longrightarrow H_1(X_L^\varphi, Y_a^\varphi) \xrightarrow{\delta} \mathbb{Z}_{q^k} \hookrightarrow S^1$$

and that δ vanishes on P , we have that

$$\tau(L, \chi) = 0 \in L^0(\mathbb{C}(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q].$$

This theorem is reminiscent of ‘Casson–Gordon type’ obstruction theorems in the context of knot concordance, especially of the main theorem of [CG86]. The proof of Theorem 1.2 is indeed guided by the knot theoretic case coupled with the main result of [FP12]. In particular we will show that if a link L is concordant to H and if χ satisfies the conditions of the theorem, then there exists a 4-manifold $W_C^\varphi = E_C^\varphi \cup (X_H^\varphi \times I)$ whose boundary is $M_L^\varphi \sqcup -M_H^\varphi = M_L^\varphi \sqcup -T$, and which has trivial (twisted) intersection form.

In Section 4 we will discuss an example of a link for which our main theorem detects that it is not concordant to the Hopf link. More precisely, we start out with a 2–component link with trivial components which is concordant to the Hopf link and we then take an appropriate satellite. We then show that our obstructions detect that the resulting link is not concordant to the Hopf link, but we will see that the methods (1) to (6) listed above do not detect this fact. It is likely though that our example gets detected by first blowing down and then considering the Casson–Gordon invariants of the resulting knot.

Remark 1.3.

- (i) When we exhibit our example, we make use of a Maple program to calculate the homology of a finite cover of the link complement explicitly. We also use Maple to identify a curve in a link diagram, for use in a satellite construction, which lifts to a generator of the torsion part of homology.
- (ii) One theme of this paper is that many arguments which work in the case of knots, for constructing concordance obstructions which involve representations that factor through derived series quotients, can be extended to the case of two component links of linking number one. Extending these arguments to the case of more components or other linking numbers seems much harder.
- (iii) It is an interesting question whether there also exists a ‘Cochran–Orr–Teichner type’ obstruction to a 2–component link being concordant to a Hopf link. Added in proof: Jae Choon Cha has recently answered this question in the affirmative: see [Cha12].
- (iv) We conjecture that Theorem 1.2 obstructs L from being ‘height (3.5)-Whitney tower/Grope concordant’ to the Hopf link in $S^3 \times I$ (compare [COT03, sections 8 and 9], [CST12]). Added in proof: Min Hoon Kim informs us that he has confirmed the conjecture: the obstructions of this paper indeed vanish for links which are height (3.5)-Whitney tower/Grope concordant to the Hopf link. At the time of writing his paper is in preparation.

- (v) Blowing down is in practice a remarkably effective method for studying link concordance. Even though we do not think that one can recover Theorem 1.2 using that approach, it seems that it will be difficult to find an example of a link that Theorem 1.2 can show is not concordant to the Hopf link but for which the blowing down approach fails.
- (vi) Whereas blowing down is rather effective as a way of obstructing links from being concordant to the Hopf link, it cannot obstruct the existence of height (3.5) Whitney tower or Grope concordances. From this point of view the conjecture in (iv) would show that the obstructions of this paper are stronger than blowing down.

1.5. Organization of the paper

The paper is organised as follows. In Section 2 we define linking forms on the torsion part of the first homology of finite covers of the link exterior, and use them to control the representations which extend over the exterior of a concordance. In Section 3 we define the obstruction $\tau(L, \chi)$ and prove Theorem 1.2. Finally, in Section 4 we construct examples of links which are not concordant to the Hopf link, and discuss the vanishing of many previously known obstructions. We conclude with a short appendix in which we prove a formula relating the homology of finite abelian covers of a 2-component link with linking number one to values of the multivariable Alexander polynomial.

2. Finite covers and linking forms

2.1. Injectivity result

Below we will quote a theorem from the authors' previous paper [FP12]. We will make use of this at several points in the construction of the obstruction theory. The special case that $\mathcal{H} \cong \mathbb{Z}$ was essentially proved by Letsche in [Let00]. Our paper makes use of the case that $\mathcal{H} \cong \mathbb{Z}^3$. Given a prime q , recall that a q -group is a finite group of q -power order. Let π be a group, let Q be a field and let \mathcal{H} be a free abelian group. Representations $\phi: \pi \rightarrow \mathcal{H}$ and $\alpha: \pi \rightarrow \text{GL}(k, Q)$ give rise to a right $\mathbb{Z}[\pi]$ -module structure on $Q^k \otimes_Q Q[\mathcal{H}] = Q[\mathcal{H}]^k$ as follows:

$$\begin{aligned} Q^k \otimes_Q Q[\mathcal{H}] \times \mathbb{Z}[\pi] &\longrightarrow Q^k \otimes_Q Q[\mathcal{H}] \\ (v \otimes p, g) &\longmapsto (v \cdot \alpha(g) \otimes p \cdot \phi(g)). \end{aligned}$$

THEOREM 2.1. *Let π be a group, let q be a prime and let $f: M \rightarrow N$ be a morphism of projective left $\mathbb{Z}[\pi]$ -modules such that*

$$\text{Id} \otimes f: \mathbb{Z}_q \otimes_{\mathbb{Z}[\pi]} M \longrightarrow \mathbb{Z}_q \otimes_{\mathbb{Z}[\pi]} N$$

is injective. Let $\phi: \pi \rightarrow \mathcal{H}$ be a homomorphism to a torsion-free abelian group and let $\alpha: \pi \rightarrow \text{GL}(k, Q)$ be a representation, with Q a field of characteristic zero. If $\alpha|_{\ker(\phi)}$ factors through a q -group, then

$$\text{Id} \otimes f: Q[\mathcal{H}]^k \otimes_{\mathbb{Z}[\pi]} M \longrightarrow Q[\mathcal{H}]^k \otimes_{\mathbb{Z}[\pi]} N$$

is also injective.

A special case of this theorem applies to prove the following lemma, which allows us to control the rank of the \mathbb{Q} -homology of finite covers of the link exterior.

LEMMA 2.2. *Suppose that $f : X \rightarrow E$ is a map of finite CW complexes which induces a \mathbb{Z} -homology equivalence*

$$f_* : H_*(X; \mathbb{Z}) \xrightarrow{\cong} H_*(E; \mathbb{Z}).$$

Let $\varphi : \pi_1(E) \rightarrow A$ be an epimorphism to a finite abelian p -group A . Let E^φ be the induced cover and let X^φ be the corresponding pull-back cover. Then

$$f_* : H_*(X^\varphi; \mathbb{Q}) \xrightarrow{\cong} H_*(E^\varphi; \mathbb{Q}).$$

is an isomorphism.

Proof. We refer to [COT03, proposition 2.10] for the details of a standard chain homotopy lifting argument, which should be applied to the relative chain complex

$$C_* := C_*(E, X; \mathbb{Q}[A]).$$

This argument was also given in the proof of [FP12, proposition 4.1]. We insert into the argument, at the point where injectivity is required, an application of Theorem 2.1. We use the special case that $\mathcal{H} \cong \{0\}$, $q = p$, $Q = \mathbb{Q}$, and $\alpha : \pi_1(E) \rightarrow A \rightarrow GL(k, \mathbb{Q})$ is given by $k = |A|$ and the regular representation $A \rightarrow \text{Aut}_Q(\mathbb{Q}[A])$. This special case was of course well-known before our contribution.

2.2. *Definition of linking forms*

Let L be an oriented 2-component link with linking number 1. We write $X = X_L$. Let $\varphi : \mathbb{Z}^2 \rightarrow A$ be any epimorphism onto a finite abelian group. We will now define a non-singular linking form:

$$\lambda = \lambda_L : TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \times TH_1(X^\varphi, Y_b^\varphi; \mathbb{Z}) \longrightarrow \mathbb{Q}/\mathbb{Z}.$$

First, we have the Poincaré duality isomorphism.

$$TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \xrightarrow{\cong} TH^2(X^\varphi, Y_b^\varphi; \mathbb{Z}).$$

By the universal coefficient theorem, since torsion in the cohomology $H^2(X^\varphi, Y_b^\varphi; \mathbb{Z})$ maps to zero in $\text{Hom}_{\mathbb{Z}}(H_2(X^\varphi, Y_b^\varphi; \mathbb{Z}), \mathbb{Z})$, there is a canonical isomorphism:

$$TH^2(X^\varphi, Y_b^\varphi; \mathbb{Z}) \xrightarrow{\cong} \text{Ext}_{\mathbb{Z}}^1(TH_1(X^\varphi, Y_b^\varphi; \mathbb{Z}), \mathbb{Z}).$$

For the reader’s convenience we now recall the proof of the following well-known claim:

Claim. Given any finite abelian group G there exists a canonical isomorphism

$$\text{Ext}_{\mathbb{Z}}^1(G, \mathbb{Z}) \xrightarrow{\cong} \text{Hom}(G, \mathbb{Q}/\mathbb{Z}).$$

Let G be any finite abelian group. The short exact sequence of coefficients

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Q} \longrightarrow \mathbb{Q}/\mathbb{Z} \longrightarrow 0$$

gives rise to a long exact sequence of cohomology

$$\text{Ext}_{\mathbb{Z}}^0(G, \mathbb{Q}) \longrightarrow \text{Ext}_{\mathbb{Z}}^0(G, \mathbb{Q}/\mathbb{Z}) \longrightarrow \text{Ext}_{\mathbb{Z}}^1(G, \mathbb{Z}) \longrightarrow \text{Ext}_{\mathbb{Z}}^1(G, \mathbb{Q}).$$

It is a simple calculation from the definition to see that $\text{Ext}_{\mathbb{Z}}^*(G, \mathbb{Q}) = 0$; we thus obtain an isomorphism $\text{Ext}_{\mathbb{Z}}^0(G, \mathbb{Q}/\mathbb{Z}) \xrightarrow{\cong} \text{Ext}_{\mathbb{Z}}^1(G, \mathbb{Z})$. Combining the inverse of this isomorphism with the canonical isomorphism $\text{Ext}_{\mathbb{Z}}^0(G, \mathbb{Q}/\mathbb{Z}) \xrightarrow{\cong} \text{Hom}(G, \mathbb{Q}/\mathbb{Z})$ shows that we have a

canonical isomorphism $\text{Ext}_{\mathbb{Z}}^1(G, \mathbb{Z}) \xrightarrow{\cong} \text{Hom}(G, \mathbb{Q}/\mathbb{Z})$. This concludes the proof of the claim.

Claim. There exists a canonical isomorphism of relative homology groups:

$$TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \xrightarrow{\cong} TH_1(X^\varphi, Y_b^\varphi; \mathbb{Z}).$$

First note that the two inclusion maps

$$\bigsqcup_2 S^1 \times D^1 \cong Y_a \hookrightarrow X_L$$

and

$$\bigsqcup_2 S^1 \times D^1 \cong Y_b \hookrightarrow X_L$$

are isotopic. We thus get an isomorphism of relative homology groups:

$$TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \xrightarrow{\cong} TH_1(X^\varphi, Y_b^\varphi; \mathbb{Z}).$$

To see that this isomorphism is canonical, consider the following \mathbb{Z} -coefficient commutative diagram.

$$\begin{array}{ccccccccc} H_1(Y_a^\varphi) & \longrightarrow & H_1(X_L^\varphi) & \longrightarrow & H_1(X_L^\varphi, Y_a^\varphi) & \longrightarrow & H_0(Y_a^\varphi) & \longrightarrow & H_0(X_L^\varphi) \\ \cong \downarrow r & & \cong \downarrow f & & \downarrow & & \cong \downarrow r & & \cong \downarrow f \\ H_1(Y_b^\varphi) & \longrightarrow & H_1(X_L^\varphi) & \longrightarrow & H_1(X_L^\varphi, Y_b^\varphi) & \longrightarrow & H_0(Y_b^\varphi) & \longrightarrow & H_0(X_L^\varphi) \end{array}$$

The maps labelled r are induced by a rotation of the two boundary tori which sends Y_a^φ to Y_b^φ . It is a homeomorphism so induces isomorphisms on homology. The maps labelled f are induced by an isotopy of homeomorphisms between this rotation of ∂X_L^φ and the identity map of ∂X_L^φ ,

$$i: \partial X_L^\varphi \times I \longrightarrow \partial X_L^\varphi.$$

We map a neighbourhood $\partial X_L^\varphi \times I$ of the boundary to itself using

$$\begin{aligned} f: \partial X_L^\varphi \times I &\longrightarrow \partial X_L^\varphi \times I \\ (x, t) &\longmapsto (i(x, t), t). \end{aligned}$$

Extending f by the identity to the rest of X_L^φ we obtain a homeomorphism, which induces the isomorphisms f on homology. The map of relative groups is also induced by the map f : note that f maps Y_a^φ to Y_b^φ . By the 5-lemma, the map of relative groups is an isomorphism. Since $H_0(Y_a^\varphi)$ is torsion-free, an element in $TH_1(X_L^\varphi, Y_a^\varphi)$ maps to zero in $H_0(Y_a^\varphi)$, and so lies in the image of $H_1(X_L^\varphi)$. This of course also holds with a replaced by b . Any element of $H_1(X_L^\varphi)$ can be taken to be disjoint from a collar neighbourhood of the boundary, which implies that the map $TH_1(X^\varphi, Y_a^\varphi) \xrightarrow{\cong} TH_1(X^\varphi, Y_b^\varphi)$ is canonical: it does not depend on choices of r or i , since their effect is restricted to a neighbourhood of the boundary of X_L^φ . This concludes the proof of the claim.

Combining the above isomorphisms yields

$$\begin{aligned} TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) &\xrightarrow{\cong} \text{Hom}_{\mathbb{Z}}(TH_1(X^\varphi, Y_b^\varphi; \mathbb{Z}), \mathbb{Q}/\mathbb{Z}) \\ &\xrightarrow{\cong} \text{Hom}_{\mathbb{Z}}(TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}), \mathbb{Q}/\mathbb{Z}), \end{aligned}$$

which defines the linking form $\lambda = \lambda_L$ and shows that it is non-singular. Standard arguments show that this linking form is symmetric.

2.3. *Linking forms and concordance*

PROPOSITION 2.3. *Let L be an oriented 2-component link which is concordant to the Hopf link via a concordance C . We write $X = X_L$. Let $\varphi: \mathbb{Z}^2 \rightarrow A$ be an admissible epimorphism onto a group of prime power order. Then*

$$P := \ker(TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \longrightarrow TH_1(E_C^\varphi, Y_a^\varphi; \mathbb{Z}))$$

is a metaboliser of the linking form λ_L .

Given an abelian group G we write $G^\wedge := \text{Hom}_{\mathbb{Z}}(G, \mathbb{Q}/\mathbb{Z})$. Note that λ_L defines a map $TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \rightarrow TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z})^\wedge$. The main technical ingredient in the proof of Proposition 2.3 is the following lemma:

LEMMA 2.4. *There are isomorphisms*

$$\vartheta_1: TH_1(E_C^\varphi, Y_a^\varphi; \mathbb{Z}) \longrightarrow TH_2(E_C^\varphi, X^\varphi; \mathbb{Z})^\wedge$$

and

$$\vartheta_2: TH_2(E_C^\varphi, X^\varphi; \mathbb{Z}) \longrightarrow TH_1(E_C^\varphi, Y_a^\varphi; \mathbb{Z})^\wedge$$

such that the following diagram commutes:

$$\begin{array}{ccccc} TH_2(E_C^\varphi, X^\varphi) & \xrightarrow{\partial} & TH_1(X^\varphi, Y_a^\varphi) & \xrightarrow{i_*} & TH_1(E_C^\varphi, Y_a^\varphi) \\ \cong \downarrow \vartheta_2 & & \cong \downarrow \lambda & & \downarrow \vartheta_1 \\ TH_1(E_C^\varphi, Y_a^\varphi)^\wedge & \xrightarrow{i^\wedge} & TH_1(X^\varphi, Y_a^\varphi)^\wedge & \xrightarrow{\partial^\wedge} & TH_2(E_C^\varphi, X^\varphi)^\wedge \end{array}$$

A standard argument shows how to deduce Proposition 2.3 from Lemma 2.4. We now provide the argument for the reader’s convenience.

Proof of Proposition 2.3. First note that the long exact sequence in homology of the pair (E_C^φ, X^φ) restricts to an exact sequence

$$TH_2(E_C^\varphi, X^\varphi) \longrightarrow TH_1(X^\varphi, Y_a^\varphi) \longrightarrow TH_1(E_C^\varphi, Y_a^\varphi)$$

since $TH_2(E_C^\varphi, X^\varphi) = H_2(E_C^\varphi, X^\varphi)$ by Lemma 2.2. This implies that the rows in the above commutative diagram are exact. Defining

$$P := \ker(i_*: TH_1(X^\varphi, Y_a^\varphi) \longrightarrow TH_1(E_C^\varphi, Y_a^\varphi)),$$

as in the statement of the proposition, we will now show that $P = P^\perp$.

First, let $x \in P$. By exactness there exists a $w \in TH_2(E_C^\varphi, X^\varphi)$ such that $\partial(w) = x$. Therefore

$$\lambda(x) = \lambda \circ \partial(w) = i^\wedge \circ \vartheta_2(w) = \vartheta_2(w) \circ i_*$$

with the second equality by commutativity of the left square. If also $y \in P$,

$$\lambda(x)(y) = \vartheta_2(w)(i_*(y)) = \vartheta_2(w)(0) = 0,$$

so that $x \in P^\perp$ and $P \subseteq P^\perp$.

Now let $y \in P^\perp$. This implies that $\partial^\wedge \circ \lambda(y) = 0$. Since the lower sequence is also exact it follows that $\lambda(y) = i^\wedge(x)$ for some $x \in TH_1(E_C^\varphi, Y_a^\varphi)^\wedge$. Since the left square of the diagram commutes, we see that $y = \partial(\vartheta_2^{-1}(x))$ i.e. $y \in P$. This shows that $P^\perp \subseteq P$.

The proof of Lemma 2.4 will require the remainder of this section, in which for brevity we omit the \mathbb{Z} coefficients for the homology groups.

To define ϑ_1 and ϑ_2 , we first use the analogue of the isomorphisms used to define λ . For ϑ_1 we have:

$$\begin{aligned} TH_1(E_C^\varphi, Y_a^\varphi) &\xrightarrow{\cong} TH^3(E_C^\varphi, \partial E_C^\varphi \setminus Y_a^\varphi) \\ &\xrightarrow{\cong} \text{Ext}_{\mathbb{Z}}^1(TH_2(E_C^\varphi, \partial E_C^\varphi \setminus Y_a^\varphi), \mathbb{Z}) \\ &\xrightarrow{\cong} \text{Ext}_{\mathbb{Z}}^0(TH_2(E_C^\varphi, \partial E_C^\varphi \setminus Y_a^\varphi), \mathbb{Q}/\mathbb{Z}) \\ &\xrightarrow{\cong} \text{Hom}(TH_2(E_C^\varphi, \partial E_C^\varphi \setminus Y_a^\varphi), \mathbb{Q}/\mathbb{Z}). \end{aligned}$$

Similarly, for ϑ_2 , we have

$$\begin{aligned} \vartheta_2: TH_2(E_C^\varphi, X^\varphi) &\xrightarrow{\cong} TH^2(E_C^\varphi, X_H^\varphi) \\ &\xrightarrow{\cong} \text{Ext}_{\mathbb{Z}}^1(TH_1(E_C^\varphi, X_H^\varphi), \mathbb{Z}) \\ &\xrightarrow{\cong} \text{Ext}_{\mathbb{Z}}^0(TH_1(E_C^\varphi, X_H^\varphi), \mathbb{Q}/\mathbb{Z}) \\ &\xrightarrow{\cong} \text{Hom}(TH_1(E_C^\varphi, X_H^\varphi), \mathbb{Q}/\mathbb{Z}). \end{aligned}$$

To finish the construction of the non-singular relative linking pairings, we will prove the following two lemmas:

LEMMA 2.5. *The inclusion $X^\varphi \rightarrow \partial E_C^\varphi \setminus Y_a^\varphi$ induces an isomorphism:*

$$TH_2(E_C^\varphi, X^\varphi) \xrightarrow{\cong} TH_2(E_C^\varphi, \partial E_C^\varphi \setminus Y_a^\varphi).$$

LEMMA 2.6. *The inclusion $Y_a^\varphi \rightarrow X_H^\varphi$ induces an isomorphism:*


$$TH_1(E_C^\varphi, Y_a^\varphi) \xrightarrow{\cong} TH_1(E_C^\varphi, X_H^\varphi).$$

Assuming Lemmas 2.5 and 2.6 we use these maps to finish the construction of ϑ_1 and ϑ_2 . It is straightforward to see that the forms ϑ_1 , ϑ_2 and λ fit into a commutative diagram as postulated in Lemma 2.4. It thus remains to prove Lemmas 2.5 and 2.6.

We continue with the notation as above. We denote the two components of Y_b by Y_{b1} and Y_{b2} . For $j = 1, 2$ we pick one component \tilde{Y}_{bj} of the preimage of Y_{bj} under the covering map $Y_b^\varphi \rightarrow Y_b$. We write $\tilde{Y}_b = \tilde{Y}_{b1} \cup \tilde{Y}_{b2}$. Note that each component of Y_b^φ is isotopic in ∂X to one of the two components of \tilde{Y}_b . We thus obtain a canonical map $Y_b^\varphi \rightarrow \tilde{Y}_b$.

We now have the following lemma:

LEMMA 2.7. *Let φ be admissible. Consider the following diagram*



Then the following hold:

- (i) the diagram commutes;
- (ii) the bottom vertical map is surjective;
- (iii) the bottom right diagonal map is an isomorphism;
- (iv) all top diagonal maps are isomorphisms with rational coefficients;
- (v) the top vertical map is rationally surjective;
- (vi) the bottom left diagonal map is injective and an isomorphism over \mathbb{Q} ;
- (vii) the bottom left diagonal map splits.

Proof of Lemma 2.7. It is evident that the diagram commutes and that the bottom vertical map is surjective (this is (i) and (ii)). Now recall that X_H is homeomorphic to a torus times an interval $S^1 \times S^1 \times I$. We denote the two meridians of H by x and y . Recall that we identified $H_1(X_H)$ with \mathbb{Z}^2 such that x corresponds to $(1, 0)$ and y corresponds to $(0, 1)$. Let $\varphi: H_1(X_H) \rightarrow A$ be any epimorphism onto a finite abelian group. We denote by m and n the orders of $\varphi(x) \in A$ and $\varphi(y) \in A$. Lifting to the covering space induced by φ , and then considering $H_1(X_H^\varphi)$ as a subgroup of $H_1(X_H)$, we see that

$$\text{im}(H_1(\tilde{Y}_b) \rightarrow H_1(X_H^\varphi) \rightarrow H_1(X_H)) = \mathbb{Z}x^m \oplus \mathbb{Z}y^n.$$

On the other hand we have $\text{im}(H_1(X_H^\varphi) \rightarrow H_1(X_H)) = \ker(\varphi: H_1(X_H) = \mathbb{Z}^2 \rightarrow A)$. It is straightforward to verify that $\text{im}(H_1(\tilde{Y}_b) \rightarrow H_1(X_H^\varphi)) = H_1(X_H^\varphi)$ if and only if

$$\ker(\varphi: \mathbb{Z}^2 \rightarrow A) = \ker(\varphi: \langle x \rangle \rightarrow A) \oplus \ker(\varphi: \langle y \rangle \rightarrow A).$$

The latter condition in turn is satisfied if and only if φ is admissible. This shows that the bottom right diagonal map is an isomorphism, which is (3).

Note that by Lemma 2.2 the top diagonal maps are isomorphisms with rational coefficients, which is (iv). Since the bottom right diagonal map is an isomorphism over the integers, it follows by commutativity, going both ways around the diamond, that the bottom left diagonal map is also an isomorphism over the rationals, which is part of (vi). It also follows by commutativity, (ii), (iii) and (iv), that the top vertical map is surjective over the rationals, which is (v). Note note that $H_1(\tilde{Y}_b)$ is torsion-free. It follows that the bottom left diagonal map is injective, which shows (vi).

Finally, consider the following commutative diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_1(\tilde{Y}_b^\varphi) & \longrightarrow & H_1(Y_b) & \longrightarrow & A \longrightarrow 0 \\ & & \downarrow & & \downarrow = & & \downarrow = \\ & & H_1(X^\varphi) & \longrightarrow & H_1(X) & \longrightarrow & A \longrightarrow 0. \end{array}$$

Note that the right two vertical maps are isomorphisms. It is a straightforward diagram chase to see that the left-hand map splits, which is (vii).

For future reference we record the following corollary to Lemma 2.7.

COROLLARY 2.8. *Let φ be admissible. Then we have an isomorphism*

$$TH_1(X^\varphi, Y_b^\varphi) \cong TH_1(X^\varphi).$$

Proof. By Lemma 2.7 (vii) the inclusion induced map $H_1(\tilde{Y}_b^\varphi) \rightarrow H_1(X^\varphi)$ admits a left inverse. It follows that

$$TH_1(X^\varphi, \tilde{Y}_b^\varphi) \cong TH_1(X^\varphi).$$

On the other hand it follows easily from comparing the long exact sequences of the pairs (X^φ, Y_b^φ) and $(X^\varphi, \tilde{Y}_b^\varphi)$ that

$$TH_1(X^\varphi, \tilde{Y}_b^\varphi) \cong TH_1(X^\varphi, Y_b^\varphi).$$

Proof of Lemma 2.5. The map which we claim is an isomorphism arises from restricting to the torsion part in the long exact sequence of the triple $(E_C^\varphi, \partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi \setminus Y_a^\varphi \simeq X^\varphi)$:

$$\begin{aligned} H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) &\rightarrow H_2(E_C^\varphi, X^\varphi) \\ &\rightarrow H_2(E_C^\varphi, \partial E_C^\varphi \setminus Y_a^\varphi) \\ &\rightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi). \end{aligned}$$

By Lemma 2.2, we have that

$$H_2(E_C^\varphi, X^\varphi) \cong TH_2(E_C^\varphi, X^\varphi).$$

It therefore suffices to show that

$$TH_1(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \cong 0$$

and that

$$H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \longrightarrow H_2(E_C^\varphi, X^\varphi)$$

is the zero map.

We first consider the following Mayer–Vietoris sequence to investigate the homology of $\partial E_C^\varphi \setminus Y_a^\varphi$, decomposing it as $X^\varphi \cup_{Y_b^\varphi} X_H^\varphi$.

$$H_1(Y_b^\varphi) \longrightarrow H_1(X^\varphi) \oplus H_1(X_H^\varphi) \longrightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi) \longrightarrow H_0(Y_b^\varphi) \longrightarrow \dots$$

We write $I := \text{im}(H_1(X^\varphi) \rightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi))$. By Lemma 2.7 (i), (ii) and (iii), the map $H_1(Y_b^\varphi) \rightarrow H_1(X_H^\varphi)$ is surjective. It then follows from the above exact sequence that

$$I = \text{im}(H_1(X^\varphi) \oplus H_1(X_H^\varphi) \rightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi)).$$

In particular, we get an inclusion

$$0 \longrightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi)/I \longrightarrow H_0(Y_b^\varphi).$$

Since $H_0(Y_b^\varphi)$ is torsion-free, it follows that $H_1(\partial E_C^\varphi \setminus Y_a^\varphi)/I$ is also torsion-free. On the other hand the pair $(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi)$ gives rise to the following exact sequence

$$0 \longrightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi)/I \longrightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \longrightarrow H_0(X^\varphi).$$

Since any subgroup of $H_0(X^\varphi)$ is torsion-free it now follows that $H_1(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi)$ is torsion-free. Put differently,

$$TH_1(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \cong 0,$$

as required.

In order to complete the proof of the lemma, we still have to show that

$$H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \longrightarrow H_2(E_C^\varphi, X^\varphi)$$

is the zero map. The following commutative diagram will guide us through the remainder of

the proof:

$$\begin{array}{ccccc}
 0 & \longrightarrow & H_2((\partial X^1)^\varphi) \oplus H_2((\partial X^2)^\varphi) & & \\
 \downarrow & & \downarrow & & \\
 H_2(X_H^\varphi) \oplus H_2(X^\varphi) & \xlongequal{\quad} & H_2(X_H^\varphi) \oplus H_2(X^\varphi) & & \\
 \downarrow & & \downarrow & & \\
 H_2(\partial E_C^\varphi \setminus Y_a^\varphi) & \longrightarrow & H_2(\partial E_C^\varphi) & \longrightarrow & H_2(\partial E_C^\varphi, X^\varphi) \\
 \downarrow \partial & & \downarrow & & \\
 H_1(Y_b^\varphi) & \xrightarrow{\quad g \quad} & H_1((\partial X^1)^\varphi) \oplus H_1((\partial X^2)^\varphi) & & \\
 \downarrow f & & \downarrow & & \\
 H_1(X_H^\varphi) \oplus H_1(X^\varphi) & \xlongequal{\quad} & H_1(X_H^\varphi) \oplus H_1(X^\varphi) & & \\
 \downarrow & & & & \\
 H_1((\partial E_C^\varphi \setminus Y_a^\varphi)). & & & &
 \end{array}$$

Note that the two vertical sequences are exact. We start out with the following claim:

Claim. The map

$$H_2(\partial E_C^\varphi \setminus Y_a^\varphi) \longrightarrow H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi)$$

is surjective.

Note that by Lemma 2.7 (i),(iii) and (vi) we know that

$$\begin{aligned}
 \ker(H_1(Y_b^\varphi) \longrightarrow H_1(X_H^\varphi)) &= \ker(H_1(Y_b^\varphi) \longrightarrow H_1(\tilde{Y}_b)) \\
 &= \ker(H_1(Y_b^\varphi) \longrightarrow H_1(X^\varphi)).
 \end{aligned}$$

Using the left vertical Mayer–Vietoris sequence in the above diagram and the fact that the map

$$H_1(Y_b^\varphi) \longrightarrow H_1(X_H^\varphi)$$

is surjective, by Lemma 2.7 (i), (ii) and (iii), this implies that the inclusion induced map

$$H_1(X^\varphi) \longrightarrow H_1(\partial E_C^\varphi \setminus Y_a^\varphi)$$

is injective. The claim follows from considering the long exact sequence in homology of the pair $(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi)$.

By the claim, we just have to show that the composition

$$H_2(\partial E_C^\varphi \setminus Y_a^\varphi) \longrightarrow H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \longrightarrow H_2(E_C^\varphi, X^\varphi)$$

is the zero map. Moreover, the map $H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \rightarrow H_2(E_C^\varphi, X^\varphi)$ factors as

$$H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \longrightarrow H_2(\partial E_C^\varphi, X^\varphi) \longrightarrow H_2(E_C^\varphi, X^\varphi),$$

so it suffices to show that the map

$$H_2(\partial E_C^\varphi \setminus Y_a^\varphi) \longrightarrow H_2(\partial E_C^\varphi, X^\varphi)$$

is the zero map.

First note that the maps $H_i(\partial(X^j)^\varphi) \rightarrow H_i(X_H^\varphi)$, $j = 1, 2$ in the above commutative diagram are induced by homotopy equivalences; in particular they are isomorphisms. It follows from the second vertical exact sequence that

$$\text{im}(H_2(X_H^\varphi) \longrightarrow H_2(\partial E_C^\varphi)) \subset \text{im}(H_2(X^\varphi) \longrightarrow H_2(\partial E_C^\varphi)).$$

The fact that $H_2(X^\varphi)$ maps to zero in $H_2(\partial E_C^\varphi, X^\varphi)$ implies that the map

$$H_2(X_H^\varphi) \oplus H_2(X^\varphi) \longrightarrow H_2(\partial E_C^\varphi) \longrightarrow H_2(\partial E_C^\varphi, X^\varphi)$$

is the zero map.

Recall that there are maps labelled as f and g in the preceding large commutative diagram.

Claim.

$$\ker(f) \subseteq \ker(g).$$

Suppose that $x \in \ker(f)$. Then in particular $x \in \ker(H_1(Y_b^\varphi) \rightarrow H_1(X_H^\varphi))$. But then, since X_H^φ is just a product $S^1 \times S^1 \times I$ and $\partial X_H^\varphi \cong \partial X^\varphi$, the inclusion induced maps $H_1((\partial X^1)^\varphi) \rightarrow H_1(X_H^\varphi)$ and $H_1((\partial X^2)^\varphi) \rightarrow H_1(X_H^\varphi)$ are isomorphisms. So indeed, using the inverses of these maps and commutativity of the bottom square, we have $x \in \ker(g)$. This concludes the proof of the claim.

A short diagram chase proves that the map

$$H_2(\partial E_C^\varphi \setminus Y_a^\varphi) \longrightarrow H_2(\partial E_C^\varphi \setminus Y_a^\varphi, X^\varphi) \longrightarrow H_2(\partial E_C^\varphi, X^\varphi)$$

is the zero map. Indeed, let $y \in H_2(\partial E_C^\varphi \setminus Y_a^\varphi)$. Then $\partial(y) \in H_1(Y_b^\varphi)$ lies in the kernel of f . Since $\ker(f) \subseteq \ker(g)$, and by the commutativity of the diagram it follows that the image of y in $H_2(\partial E_C^\varphi)$ lies in the image of $H_2(X_H^\varphi) \oplus H_2(X^\varphi)$. But we saw above that the map to $H_2(\partial E_C^\varphi, X^\varphi)$ vanishes on the image of $H_2(X_H^\varphi) \oplus H_2(X^\varphi)$.

For the proof of Lemma 2.6 we will need the following lemma, which we will also use in Section 2.4.

LEMMA 2.9. *Let $J = H$ or $J = L$. We write*

$$F := H_1(X_J^\varphi, Y_a^\varphi; \mathbb{Z})/\text{torsion}.$$

Then there exists a commutative diagram

$$\begin{array}{ccc} T H_1(X_J^\varphi, Y_a^\varphi) \oplus F & \xrightarrow{\cong} & H_1(X_J^\varphi, Y_a^\varphi) \\ \downarrow i_* \oplus \text{Id} & & \downarrow i_* \\ T H_1(E_C^\varphi, Y_a^\varphi) \oplus F & \xrightarrow{\cong} & H_1(E_C^\varphi, Y_a^\varphi). \end{array}$$

Proof. We have the following commutative diagram.

$$\begin{array}{ccccccccc} H_1(Y_a^\varphi) & \longrightarrow & H_1(X_J^\varphi) & \longrightarrow & H_1(X_J^\varphi, Y_a^\varphi) & \longrightarrow & H_0(Y_a^\varphi) & \longrightarrow & H_0(X_J^\varphi) \\ \parallel & & \downarrow & & \downarrow & & \parallel & & \cong \downarrow \\ H_1(Y_a^\varphi) & \longrightarrow & H_1(E_C^\varphi) & \longrightarrow & H_1(E_C^\varphi, Y_a^\varphi) & \longrightarrow & H_0(Y_a^\varphi) & \longrightarrow & H_0(E_C^\varphi) \end{array}$$

It follows from Lemma 2.7 that the two left horizontal maps are rationally surjective. We write $F' := \ker(H_0(Y_a^\varphi) \rightarrow H_0(X_J^\varphi))$, which equals $\ker(H_0(Y_a^\varphi) \rightarrow H_0(E_C^\varphi))$. Note that F' , as a subgroup of a torsion-free group is itself torsion-free. We now obtain the following commutative diagram of short exact sequences:

$$\begin{CD} 0 @>>> H_1(X_J^\varphi)/\text{im}(H_1(Y_a^\varphi)) @>>> H_1(X_J^\varphi, Y_a^\varphi) @>>> F' @>>> 0 \\ @. @VVV @VVV @| \\ 0 @>>> H_1(E_C^\varphi)/\text{im}(H_1(Y_a^\varphi)) @>>> H_1(E_C^\varphi, Y_a^\varphi) @>>> F' @>>> 0. \end{CD}$$

By the above the two groups on the left are torsion. Since F' is torsion-free it follows that the images of the groups on the left are precisely the torsion subgroups. We pick an isomorphism $F \rightarrow F'$ and a splitting $F' \rightarrow H_1(X_J^\varphi, Y_a^\varphi)$. Together with the induced splitting $F' \rightarrow H_1(X_J^\varphi, Y_a^\varphi) \rightarrow H_1(E_C^\varphi, Y_a^\varphi)$ we obtain the desired commutative diagram.

Proof of Lemma 2.6. Recall the statement of Lemma 2.6: the inclusion $Y_a^\varphi \rightarrow X_H^\varphi$ induces an isomorphism

$$TH_1(E_C^\varphi, Y_a^\varphi) \xrightarrow{\cong} TH_1(E_C^\varphi, X_H^\varphi).$$

We apply Lemma 2.9 to the case $J = H$. We obtain the following commutative diagram, where the right hand column contains a portion of the long exact sequence of the triple.

$$\begin{CD} TH_1(X_H^\varphi, Y_a^\varphi) \oplus F @>\cong>> H_1(X_H^\varphi, Y_a^\varphi) \\ @V i_* \oplus \text{Id} VV @VV i_* V \\ TH_1(E_C^\varphi, Y_a^\varphi) \oplus F @>\cong>> H_1(E_C^\varphi, Y_a^\varphi) \\ @. @VVV \\ @. H_1(E_C^\varphi, X_H^\varphi) = TH_1(E_C^\varphi, X_H^\varphi) \\ @. @VVV \\ @. H_0(X_H^\varphi, Y_a^\varphi) = 0 \end{CD}$$

We know that

$$H_1(E_C^\varphi, X_H^\varphi) = TH_1(E_C^\varphi, X_H^\varphi)$$

by Lemma 2.2. We immediately obtain the desired conclusion from the observation that $TH_1(X_H^\varphi, Y_a^\varphi) = 0$. This is a consequence of the proof of Lemma 2.9, which shows that $TH_1(X_H^\varphi, Y_a^\varphi)$ is precisely $H_1(X_H^\varphi)/\text{im}(H_1(Y_a^\varphi))$. But by Lemma 2.7 (i), (ii) and (iii), $H_1(Y_a^\varphi) \rightarrow H_1(X_H^\varphi)$ is surjective.

2.4. Extending characters

PROPOSITION 2.10. *Let L be an oriented 2-component link which is concordant to the Hopf link via a concordance C . We write $X = X_L$. Let $\varphi : H_1(X; \mathbb{Z}) = \mathbb{Z}^2 \rightarrow A$ be an admissible epimorphism onto a group of prime power order. Then there exists a metaboliser $P \subset TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z})$ for the linking form λ_L such that for any prime q and any homomorphism $\chi : H_1(X^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_{q^k}$ which factors through $H_1(X^\varphi; \mathbb{Z}) \rightarrow H_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_{q^k}$ and which vanishes on P , there exists an integer $l \geq k$ and a*

homomorphism $H_1(E_C^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_{q^l}$, such that following diagram commutes:

$$\begin{array}{ccc} H_1(X^\varphi; \mathbb{Z}) & \xrightarrow{\chi} & \mathbb{Z}_{q^k} \\ \downarrow & & \downarrow \cdot q^{l-k} \\ H_1(E_C^\varphi; \mathbb{Z}) & \longrightarrow & \mathbb{Z}_{q^l}. \end{array}$$

Proof. We start out with the following elementary observation: let $A \rightarrow B$ be a homomorphism between finite abelian groups and let $\chi : A \rightarrow \mathbb{Z}_{q^k}$ be a homomorphism to a cyclic group of prime power order which vanishes on $\ker(A \rightarrow B)$. Then there exists an integer $l \geq k$ and a homomorphism $\chi' : B \rightarrow \mathbb{Z}_{q^l}$, such that the following diagram commutes:

$$\begin{array}{ccc} A & \longrightarrow & \mathbb{Z}_{q^k} \\ \downarrow & & \downarrow \cdot q^{l-k} \\ B & \longrightarrow & \mathbb{Z}_{q^l}. \end{array}$$

For example, this observation can be proved using the classification of finite abelian groups.

Now let

$$P := \ker(TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \rightarrow TH_1(E_C^\varphi, Y_a^\varphi; \mathbb{Z})).$$

By Proposition 2.3 this is a metaboliser of the linking form λ_L . Now let χ be as in the statement of the lemma. We apply Lemma 2.9 to $J = L$. We write

$$F := H_1(X_L^\varphi, Y_a^\varphi; \mathbb{Z})/\text{torsion}$$

and we obtain the following commutative diagram

$$\begin{array}{ccccc} H_1(X_L^\varphi) & \longrightarrow & H_1(X_L^\varphi, Y_a^\varphi) & \xrightarrow{\cong} & TH_1(X_L^\varphi, Y_a^\varphi) \oplus F & \xrightarrow{\chi_T \oplus \chi_F} & \mathbb{Z}_{q^k} \\ \downarrow & & \downarrow i_* & & \downarrow i_* \oplus \text{Id} & & \\ H_1(E_C^\varphi) & \longrightarrow & H_1(E_C^\varphi, Y_a^\varphi) & \xrightarrow{\cong} & TH_1(E_C^\varphi, Y_a^\varphi) \oplus F & & \end{array}$$

Note that since χ vanishes on P , we can apply the above observation to $A := TH_1(X_L^\varphi, Y_a^\varphi) \rightarrow B := TH_1(E_C^\varphi, Y_a^\varphi)$ and to χ_T to obtain a homomorphism $\chi'_T : TH_1(E_C^\varphi, Y_a^\varphi) \rightarrow \mathbb{Z}_{q^l}$ with the appropriate properties. The homomorphisms

$$H_1(E_C^\varphi) \longrightarrow H_1(E_C^\varphi, Y_a^\varphi) \xrightarrow{\cong} TH_1(E_C^\varphi, Y_a^\varphi) \oplus F \xrightarrow{\chi'_T \oplus q^{l-k} \cdot \chi_F} \mathbb{Z}_{q^l}$$

complete the diagram as required.

For a two-component link L of linking number 1, we define the closed 3-manifold M_L to be given by the union

$$M_L := X_L \cup_{\partial X_L} -X_H$$

of the exterior of L and the exterior of the Hopf link glued together along their common boundary $\partial X_L = \partial X_H$, respecting the ordering of the link components and identifying each of the subsets $Y_a \subset \partial X_J$ for $J = L, H$. Note that $H_1(M_L; \mathbb{Z}) \cong \mathbb{Z}^3$ and that, if $L = H$, we have $M_H \cong T = S^1 \times S^1 \times S^1$.

LEMMA 2.11. Let $\varphi: H_1(M_L; \mathbb{Z}) \rightarrow A$ be a homomorphism to a finite abelian p -group A . Then $H_1(X_L^\varphi; \mathbb{Z})$ fits into a short exact sequence

$$0 \longrightarrow H_1(X_L^\varphi; \mathbb{Z}) \longrightarrow H_1(M_L^\varphi; \mathbb{Z}) \longrightarrow \mathbb{Z} \longrightarrow 0,$$

so that $H_1(X_L^\varphi; \mathbb{Z})$ is a subgroup of $H_1(M_L^\varphi; \mathbb{Z})$.

Proof. Consider the following exact sequence which arises from the Mayer–Vietoris sequence:

$$H_1((\partial X_L^1)^\varphi) \oplus H_1((\partial X_L^2)^\varphi) \longrightarrow H_1(X_L^\varphi) \oplus H_1(X_H^\varphi) \longrightarrow H_1(M_L^\varphi) \longrightarrow \mathbb{Z} \longrightarrow 0.$$

The \mathbb{Z} -term is given by

$$\ker(H_0((\partial X_L^1)^\varphi) \oplus H_0((\partial X_L^2)^\varphi) \longrightarrow H_0(X_L^\varphi) \oplus H_0(X_H^\varphi)).$$

We make the following claim.

Claim.

$$\begin{aligned} & \ker(H_1((\partial X_L^1)^\varphi) \oplus H_1((\partial X_L^2)^\varphi) \longrightarrow H_1(X_H^\varphi)) \\ & \subseteq \ker(H_1((\partial X_L^1)^\varphi) \oplus H_1((\partial X_L^2)^\varphi) \longrightarrow H_1(X_L^\varphi)). \end{aligned}$$

Since the maps $(\partial X_H^i)^\varphi \rightarrow X_H^\varphi$ are homotopy equivalences for $i = 1, 2$, the map $H_1((\partial X_L^1)^\varphi) \oplus H_1((\partial X_L^2)^\varphi) \rightarrow H_1(X_H^\varphi)$ is surjective. The claim then implies that the Mayer–Vietoris sequence for M_L^φ becomes the short exact sequence

$$0 \longrightarrow H_1(X_L^\varphi) \longrightarrow H_1(M_L^\varphi) \longrightarrow \mathbb{Z} \longrightarrow 0.$$

as required.

We now turn to the proof of the claim. Let $\pi: Y^\varphi \rightarrow Y$ be a finite covering: there are induced maps $\pi_*: H_*(Y^\varphi) \rightarrow H_*(Y)$, and transfer maps $t_*: H_*(Y) \rightarrow H_*(Y^\varphi)$. To prove the claim, we consider the following commutative diagram:

$$\begin{array}{ccccc} H_1(X_L^\varphi) & \xleftarrow{h} & H_1((\partial X_L^1)^\varphi) \oplus H_1((\partial X_L^2)^\varphi) & \xrightarrow{\quad} & H_1(X_H^\varphi) \\ \uparrow \pi_* & & \uparrow t_* \oplus t_* & & \uparrow \pi_* \\ H_1(X_L) & \xleftarrow{f} & H_1(\partial X_L^1) \oplus H_1(\partial X_L^2) & \xrightarrow{g} & H_1(X_H). \end{array}$$

Suppose that we have

$$x \in \ker(H_1((\partial X_L^1)^\varphi) \oplus H_1((\partial X_L^2)^\varphi) \longrightarrow H_1(X_H^\varphi)).$$

Then $x \mapsto 0 \in H_1(X_H)$. In the base spaces, we have that $\ker f = \ker g$, with both kernels generated by $[\mu_1 - \lambda_2]$ and $[\mu_2 - \lambda_1]$, where μ_i, λ_i is the meridian and longitude of the link component i . Since the right-hand square with down arrows is commutative, we see that

$$f((\pi_* \oplus \pi_*)(x)) = 0 \in H_1(X_L).$$

We write $y := (\pi_* \oplus \pi_*)(x)$. Since the left-hand square is commutative with up arrows, we have that

$$h((t_* \oplus t_*)(y)) = t_*(f(y)) = t_*(0) = 0 \in H_1(X_L^\varphi).$$

We then observe that for tori such as $\partial X_L^i \cong S^1 \times S^1$, we have that

$$t_* \circ \pi_*: H_1((\partial X_L^i)^\varphi) \longrightarrow H_1((\partial X_L^i)^\varphi)$$

is given by multiplication by p^α for some $\alpha \geq 0$, which may depend on x . Therefore

$$0 = h((t_* \oplus t_*)(y)) = h((t_* \oplus t_*)(\pi_* \oplus \pi_*)(x)) = h(p^\alpha x) = p^\alpha h(x).$$

We have that the image $h(x)$ of x in $H_1(X_L^\varphi)$ is either p -primary torsion, or is zero. To prove the claim we therefore wish to show that the latter case always holds. We therefore proceed to show that there is no p -primary torsion in $H_1(X_L^\varphi)$.

We use the fact that the inclusion map $\partial X_L^1 \rightarrow X_L$ induces a \mathbb{Z} -homology equivalence $H_*(\partial X_L^1; \mathbb{Z}) \xrightarrow{\cong} H_*(X_L; \mathbb{Z})$ to deduce that there is also an isomorphism $H_*((\partial X_L^1)^\varphi; \mathbb{Z}_{(p)}) \xrightarrow{\cong} H_*(X_L^\varphi; \mathbb{Z}_{(p)})$. This deduction is an extension of Lemma 2.2 to $\mathbb{Z}_{(p)}$ coefficients. We refer to [Cha10, lemma 3.3], which follows from [Lev94, lemma 4.3]. Levine shows that inverting integers coprime to p in the coefficients of the group ring of a p -group A , to obtain $\mathbb{Z}_{(p)}[A]$, is sufficient to invert square matrices over $\mathbb{Z}[A]$ which are invertible over \mathbb{Z}_p .

This then implies, since $H_1((\partial X_L^1)^\varphi; \mathbb{Z}) \cong \mathbb{Z}_{(p)}^2$, that there is no p -primary torsion in $H_1(X_L^\varphi; \mathbb{Z})$. We make use of the universal coefficient theorem and the fact that $H_0((\partial X_L^1)^\varphi; \mathbb{Z})$ and $H_0(X_L^\varphi; \mathbb{Z})$ are torsion free. This completes the proof of the claim.

Suppose that L is concordant to H with concordance C . The exterior E_C of the concordance is a 4-manifold with boundary $X_L \cup (\partial X_H \times I) \cup X_H$. We can glue E_C and $X_H \times I$ together along the $\partial X_H \times I$ components of their boundaries to define the manifold

$$W_C := E_C \cup (X_H \times I),$$

glued in such a way that W_C is a 4-manifold with boundary $\partial W_C \cong M_L \cup -M_H$.

For $J = L, H$ we have the following diagram with exact rows given by Mayer–Vietoris sequences with \mathbb{Z} coefficients and vertical maps induced by the inclusion maps:

$$\begin{array}{ccccccc} H_i(\partial X_J) & \longrightarrow & H_i(X_J) \oplus H_i(X_H) & \longrightarrow & H_i(M_J) & \longrightarrow & H_{i-1}(\partial X_J) \\ \cong \downarrow & & \cong \downarrow & & \downarrow & & \cong \downarrow \\ H_i(\partial X_J \times I) & \longrightarrow & H_i(E_C) \oplus H_i(X_H \times I) & \longrightarrow & H_i(W_C) & \longrightarrow & H_{i-1}(\partial X_J \times I). \end{array}$$


We already know that the vertical maps, except possibly the maps $H_i(M_J) \rightarrow H_i(W_C)$, are isomorphisms. But the 5-lemma then implies that the maps $H_i(M_J) \rightarrow H_i(W_C)$ are also isomorphisms, i.e. the inclusions $M_L \rightarrow W_C$ and $M_H \rightarrow W_C$ induce \mathbb{Z} -homology equivalences.

Now let $\varphi: H_1(M_L^\varphi; \mathbb{Z}) \rightarrow A$ be such that φ restricted to $H_1(X_L; \mathbb{Z}) \subset H_1(M_L; \mathbb{Z})$ is admissible. We also call such homomorphisms admissible and we denote the corresponding cover of M_L by M_L^φ .

Using the isomorphism on first homology $H_1(M_L; \mathbb{Z}) \xrightarrow{\cong} H_1(W_C; \mathbb{Z})$, the map φ extends to maps $\varphi: H_1(W_C; \mathbb{Z}) \rightarrow A$ and $H_1(M_H; \mathbb{Z}) \rightarrow A$. We can therefore define the corresponding covering spaces, which we denote by W_C^φ and M_H^φ .

PROPOSITION 2.12. *Let L be an oriented 2-component link which is concordant to the Hopf link via a concordance C . Let $\varphi: H_1(M_L; \mathbb{Z}) \rightarrow A$ be an admissible epimorphism onto a group of prime power order. Suppose that we have a character $\chi: H_1(M_L^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_{q^k}$ such that $\chi|_{H_1(X_L^\varphi)}: H_1(X_L^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_{q^k}$ extends to $\chi': H_1(E_C^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_{q^l}$. Then χ also extends to $\chi': H_1(W_C^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_{q^l}$. In other words, the map which is represented by the dotted arrow in*

the following commutative diagram exists:



Proof. We consider the exact Mayer-Vietoris sequence:

$$\begin{aligned}
 & H_1((\partial X_L^1 \times I)^\varphi) \oplus H_1((\partial X_L^2 \times I)^\varphi) \\
 & \longrightarrow H_1(E_C^\varphi) \oplus H_1(X_H^\varphi \times I) \longrightarrow H_1(W_C^\varphi) \longrightarrow \mathbb{Z} \longrightarrow 0,
 \end{aligned}$$

where the \mathbb{Z} term is given by

$$\ker(H_0((\partial X_L^1 \times I)^\varphi) \oplus H_0((\partial X_L^2 \times I)^\varphi) \longrightarrow H_0(E_C^\varphi) \oplus H_0(X_H^\varphi \times I)).$$

We apply the same argument as in the proof of Lemma 2.11, noting also that the inclusion induces isomorphisms

$$H_*(X_L; \mathbb{Z}) \xrightarrow{\cong} H_*(E_C; \mathbb{Z})$$

and therefore isomorphisms

$$H_*(X_L^\varphi; \mathbb{Z}_{(p)}) \xrightarrow{\cong} H_*(E_C^\varphi; \mathbb{Z}_{(p)}).$$

This means that we also have no p -primary torsion in $H_1(E_C^\varphi; \mathbb{Z})$, so the same argument with the transfer maps does indeed apply to show that the Mayer-Vietoris sequence becomes

$$0 \longrightarrow H_1(E_C^\varphi) \longrightarrow H_1(W_C^\varphi) \longrightarrow \mathbb{Z} \longrightarrow 0.$$

We have the commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_1(X_L^\varphi) & \longrightarrow & H_1(M_L^\varphi) & \longrightarrow & \mathbb{Z} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & H_1(E_C^\varphi) & \longrightarrow & H_1(W_C^\varphi) & \longrightarrow & \mathbb{Z} \longrightarrow 0.
 \end{array}$$

Choosing a splitting map $\mathbb{Z} \rightarrow H_1(M_L^\varphi)$ for the top row induces a splitting of the bottom row by commutativity, so that we have:

$$\begin{array}{ccc}
 H_1(X_L^\varphi) \oplus \mathbb{Z} & \xrightarrow{\cong} & H_1(M_L^\varphi) \\
 \downarrow i_* \oplus \text{Id} & & \downarrow \\
 H_1(E_C^\varphi) \oplus \mathbb{Z} & \xrightarrow{\cong} & H_1(W_C^\varphi).
 \end{array}$$

We then see that a character which extends from $H_1(X_L^\varphi)$ to $H_1(E_C^\varphi)$ also extends from $H_1(M_L^\varphi)$ over $H_1(W_C^\varphi)$, which completes the proof.

3. Obstructions

3.1. Preparation

We quote the following proposition, again from [FP12]. This is a consequence of Theorem 2.1, which comes from the same paper, and it will be used in key places in the construction of our obstruction theory. For our applications we will take \mathcal{H}' to have rank 3, and $A := \mathbb{Z}_{p^\alpha} \oplus \mathbb{Z}_{p^\beta}$.

PROPOSITION 3.1. *Let q be a prime. Suppose that S, Y are finite CW-complexes such that there is a map $i : S \rightarrow Y$ which induces an isomorphism*

$$i_* : H_*(S; \mathbb{Z}_q) \xrightarrow{\cong} H_*(Y; \mathbb{Z}_q).$$

Let $\phi' : \pi_1(Y) \rightarrow \mathcal{H}'$ be a homomorphism to a torsion-free abelian group \mathcal{H}' and let $\varphi : \pi_1(Y) \rightarrow \mathcal{H}' \rightarrow A$ be an epimorphism onto a finite abelian group which factors through ϕ' . Let Y^φ be the induced cover of Y , let $\phi : \pi_1(Y^\varphi) \rightarrow \mathcal{H}'$ be the restriction of ϕ' , let $\mathcal{H} \subseteq \mathcal{H}'$ be the image of ϕ , and let $\alpha' : \pi_1(Y^\varphi) \rightarrow \text{GL}(d, \mathbb{Q})$ be a d -dimensional representation to a field \mathbb{Q} of characteristic zero, such that α' restricted to the kernel of ϕ factors through a q -group. Define S^φ to be the pull-back cover $S^\varphi := i^*(Y^\varphi)$. Then

$$i_*^\varphi : H_*(S^\varphi; \mathbb{Q}(\mathcal{H})^d) \xrightarrow{\cong} H_*(Y^\varphi; \mathbb{Q}(\mathcal{H})^d)$$

is an isomorphism.

We now set the stage for the definition of our invariant. Let $L = L_1 \cup L_2$ be a two-component oriented link with $\text{lk}(L_1, L_2) = 1$. Choose a homomorphism

$$\phi' : \pi_1(M_L) \longrightarrow H_1(M_L) \xrightarrow{\cong} \mathcal{H}'$$

onto a free abelian group of rank 3. Choose a prime p , a p -group $A = \mathbb{Z}_{p^\alpha} \oplus \mathbb{Z}_{p^\beta}$ for integers α, β , and an admissible homomorphism $\varphi : \pi_1(M_L) \rightarrow H_1(M_L; \mathbb{Z}) \rightarrow A$. Define

$$\phi : \pi_1(M_L^\varphi) \longrightarrow \mathcal{H}'$$

to be the restriction of ϕ' to

$$\pi_1(M_L^\varphi) \cong \ker(\pi_1(M_L) \longrightarrow H_1(M_L) \xrightarrow{\varphi} A)$$

and let $\mathcal{H} \subseteq \mathcal{H}'$ be the image of ϕ . As a finite index subgroup \mathcal{H} is also free abelian of rank 3. Finally, we choose a homomorphism

$$\chi : \pi_1(M_L^\varphi) \longrightarrow H_1(M_L^\varphi) \longrightarrow \mathbb{Z}_{q^k},$$

where q is a prime and $k \geq 1$ is an integer. In this section, we proceed as follows. We will define an invariant of the isotopy class of L , $\tau(L, \chi) := \tau(L, \phi', \varphi, \chi)$, as a Witt class of a certain combination of intersection forms on a 4-manifold W whose boundary is a collection of copies of $M_L \sqcup -M_H$. We will then show that our invariant is well-defined, in that it does not depend on the choices we make to initially define it. Our main theorem is then that this invariant gives concordance obstructions for certain judicious choices of the representation χ .

3.2. Definition of $\tau(L, \chi)$

Note that X_J , for $J = L, H$, is an irreducible 3-manifold. It is well known that the result of gluing two irreducible 3-manifolds along incompressible tori is again irreducible. It follows that M_L is also irreducible.

The homomorphism $\phi: \pi_1(M_L^\varphi) \rightarrow \mathcal{H}$ induces a map

$$\phi_*: H_i(\pi_1(M_L^\varphi); \mathbb{Z}) \rightarrow H_i(\mathcal{H}; \mathbb{Z})$$

for all i . Since covering maps induce isomorphisms on homotopy groups, by the Papakyriakopoulos sphere theorem and the fact that M_L is irreducible, we have that M_L^φ is an Eilenberg-MacLane space $K(\pi_1(M_L^\varphi), 1)$. Therefore

$$H_i(\pi_1(M_L^\varphi); \mathbb{Z}) \cong H_i(M_L^\varphi; \mathbb{Z}).$$

Claim. The map $\phi: \pi_1(M_L^\varphi) \rightarrow \mathcal{H}$ gives rise to an isomorphism $\phi_*: H_3(M_L^\varphi; \mathbb{Z}) \rightarrow H_3(\mathcal{H}; \mathbb{Z})$.

First note that the map

$$\phi': \pi_1(M_L) \rightarrow \mathcal{H}'$$

allows us to define a map $M_L \rightarrow T$, by elementary obstruction theory, since T is a $K(\mathcal{H}', 1)$. Both M_L and T have the same cup product structure with \mathbb{Z} coefficients, which easily implies that

$$\phi'^*: H^3(\mathcal{H}'; \mathbb{Z}) = H^3(T; \mathbb{Z}) \xrightarrow{\cong} H^3(M_L; \mathbb{Z}) = H^3(\pi_1(M_L); \mathbb{Z})$$

is an isomorphism. By the universal coefficient theorem and Poincaré duality,

$$TH_2(M_L; \mathbb{Z}) \cong TH^1(M_L; \mathbb{Z}) \cong TH_0(M_L; \mathbb{Z}) \cong 0.$$

Similarly,

$$TH_2(T; \mathbb{Z}) \cong 0.$$

This implies, again by the universal coefficient theorem, making choices of fundamental classes $[M_L]$ and $[T]$, that

$$H^3(M_L; \mathbb{Z}) \cong H_3(M_L; \mathbb{Z})$$

and

$$H^3(T; \mathbb{Z}) \cong H_3(T; \mathbb{Z}).$$

The representation ϕ' therefore induces an isomorphism

$$\phi'_*: H_3(M_L; \mathbb{Z}) \xrightarrow{\cong} H_3(T; \mathbb{Z}).$$

We then restrict to the finite index subgroups $\pi_1(M_L^\varphi) \subseteq \pi_1(M_L)$ and $\mathcal{H} \subseteq \mathcal{H}'$. The transfer maps induce isomorphisms

$$t_*: H_3(M_L; \mathbb{Z}) \xrightarrow{\cong} H_3(M_L^\varphi; \mathbb{Z})$$

and

$$t_*: H_3(T; \mathbb{Z}) \xrightarrow{\cong} H_3(T^\varphi; \mathbb{Z}),$$

where we denote by T^φ the corresponding finite cover of T , which is nonetheless abstractly homeomorphic to T . We thus induce an isomorphism

$$H_3(\pi_1(M_L^\varphi); \mathbb{Z}) = H_3(M_L^\varphi; \mathbb{Z}) \xrightarrow{\cong} H_3(\mathcal{H}; \mathbb{Z}).$$

This concludes the proof of the claim.

Now let $T := S^1 \times S^1 \times S^1$ be the 3-torus. It follows from the above claim that there exists an isomorphism $\psi : \pi_1(T) \cong H_1(T; \mathbb{Z}) \xrightarrow{\sim} \mathcal{H}$ such that

$$\phi_*([M_L^\varphi]) = \psi_*([T]) \in H_3(\mathcal{H}) \cong \mathbb{Z}. \tag{3.1}$$

We now follow the Casson–Gordon program [CG86] for defining obstructions. We first have the following claim:

Claim. There exists an integer $s \geq 0$ and a 4-manifold W with a representation $\kappa \times \Phi : \pi_1(W) \rightarrow \mathbb{Z}_{q^k} \times \mathcal{H}$ and an orientation preserving homeomorphism

$$I : \bigsqcup_{q^s} (M_L^\varphi \sqcup -T) \longrightarrow \partial W$$

such that, restricted to each component of the boundary,

$$(\kappa \times \Phi)| \circ I_* = \chi \times \phi : \pi_1(M_L^\varphi) \longrightarrow \mathbb{Z}_{q^k} \times \mathcal{H}$$

and

$$(\kappa \times \Phi)| \circ I_* = \text{tr} \times \psi : \pi_1(T) \longrightarrow \mathbb{Z}_{q^k} \times \mathcal{H},$$

where tr is the trivial homomorphism.

First note that

$$(M_L^\varphi, \chi \times \phi) \cup -(T, \text{tr} \times \psi)$$

defines an element in $\Omega_3(\mathbb{Z}_{q^k} \times \mathcal{H})$. By the Atiyah–Hirzebruch spectral sequence,

$$\Omega_3(\mathbb{Z}_{q^k} \times \mathcal{H}) \cong H_3(\mathbb{Z}_{q^k} \times \mathcal{H}; \Omega_0) \cong H_3(\mathbb{Z}_{q^k} \times \mathcal{H}; \mathbb{Z}).$$

By the Künneth theorem,

$$H_3(\mathbb{Z}_{q^k} \times \mathcal{H}) \cong \bigoplus_{j=0}^3 H_j(\mathbb{Z}_{q^k}) \otimes H_{3-j}(\mathcal{H})$$

$$\begin{aligned} &\cong (H_0(\mathbb{Z}_{q^k}) \otimes H_3(\mathcal{H})) \oplus (H_1(\mathbb{Z}_{q^k}) \otimes H_2(\mathcal{H})) \oplus (H_3(\mathbb{Z}_{q^k}) \otimes H_0(\mathcal{H})) \\ &\cong (\mathbb{Z} \otimes H_3(\mathcal{H})) \oplus (\mathbb{Z}_{q^k} \otimes \mathbb{Z}^3) \oplus (\mathbb{Z}_{q^k} \otimes \mathbb{Z}) \end{aligned}$$

since $H_2(\mathbb{Z}_{q^k}; \mathbb{Z}) \cong 0$. Therefore

$$\Omega_3(\mathbb{Z}_{q^k} \times \mathcal{H}) \cong H_3(\mathcal{H}; \mathbb{Z}) \oplus \bigoplus_4 \mathbb{Z}_{q^k}.$$

By (3.1), the orientations are such that the image of

$$(M_L^\varphi, \chi \times \phi) \cup -(T, \text{tr} \times \psi)$$

in $H_3(\mathcal{H}; \mathbb{Z})$ is zero. Therefore $M_L \sqcup -T$, with the representations, is q -primary torsion in $\Omega_3(\mathbb{Z}_{q^k} \times \mathcal{H})$. This concludes the proof of the claim.

We choose an inclusion $\mathbb{Z}_{q^k} \hookrightarrow S^1 \subseteq \mathbb{C}$. Then, using the composition of ring homomorphisms

$$\mathbb{Z}[\pi_1(W)] \longrightarrow \mathbb{Z}[\mathbb{Z}_{q^k} \times \mathcal{H}] = \mathbb{Z}[\mathbb{Z}_{q^k}][\mathcal{H}] \longrightarrow \mathbb{Z}[\xi_{q^k}][\mathcal{H}] \longrightarrow \mathbb{Q}(\xi_{q^k})(\mathcal{H}),$$

where $\xi_{q^k} \in S^1 \subseteq \mathbb{C}$ is a primitive q^k th root of unity, we can define the twisted intersection form $\lambda_{\mathbb{Q}(\xi_{q^k})(\mathcal{H})}$ to be the non-singular part of the middle dimensional intersection form on

$H_2(W; \mathbb{Q}(\xi_{q^k})(\mathcal{H}))$. If we also consider the non-singular part of the ordinary intersection form $\lambda_{\mathbb{Q}}$ on rational homology, we obtain another intersection form, which we can also consider as a form of $\mathbb{Q}(\xi_{q^k})(\mathcal{H})$ -modules by tensoring up via the inclusion $\mathbb{Q} \rightarrow \mathbb{Q}(\xi_{q^k})(\mathcal{H})$. We define $\tau(L, \chi)$ to be

$$(\lambda_{\mathbb{Q}(\xi_{q^k})(\mathcal{H})}(W) - \mathbb{Q}(\xi_{q^k})(\mathcal{H}) \otimes_{\mathbb{Q}} \lambda_{\mathbb{Q}}) \otimes \frac{1}{q^s} \in L^0(\mathbb{Q}(\xi_{q^k})(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q].$$

We first show that $\tau(L, \chi)$ is well-defined. As usual when defining such invariants, the following lemma simplifies several key arguments. In particular, it implies that the intersection form $\lambda_{\mathbb{Q}(\xi_{q^k})(\mathcal{H})}(W)$ is non-singular on the whole of $H_2(W; \mathbb{Q}(\xi_{q^k})(\mathcal{H}))$. When $J = L$ we use $\alpha' = \chi$ and when $J = H$ we use the trivial representation for α' .

LEMMA 3.2. *Let p, q be primes. For any two-component link J with $\text{lk}(J_1, J_2) = 1$, with an isomorphism $\phi': H_1(M_J; \mathbb{Z}) \rightarrow \mathcal{H}'$, a homomorphism $\varphi: H_1(M_J; \mathbb{Z}) \rightarrow \mathcal{H}' \rightarrow A$ to a finite abelian p -group and a representation*

$$\alpha': H_1(M_J^{\varphi}; \mathbb{Z}) \longrightarrow \mathbb{Z}_{q^k} \hookrightarrow S^1 \subset \mathbb{C},$$

we have that

$$H_*(M_J^{\varphi}; \mathbb{Q}(\xi_{q^k})(\mathcal{H})) \cong 0,$$

where ξ_{q^k} is a primitive q^k th root of unity.

Proof. Since the linking number is 1, the inclusion maps $\partial X_J^1 \rightarrow X_J$ are \mathbb{Z} -homology equivalences. We wish to apply Proposition 3.1 with $S = S^1 \times S^1$ to see that

$$H_*(X_J^{\varphi}; \mathbb{Q}(\xi_{q^k})(\mathcal{H})) \cong H_*((\partial X_J^1)^{\varphi}; \mathbb{Q}(\xi_{q^k})(\mathcal{H})) \cong 0.$$

In order to apply Proposition 3.1, we need to note that $\mathbb{Z}[\xi_{q^k}]$ is a domain, so the cyclotomic field $\mathcal{Q} := \mathbb{Q}(\xi_{q^k})$ is a field of characteristic zero, and that α' factors through a q -group.

The Mayer–Vietoris sequence of $M_J^{\varphi} = X_J^{\varphi} \cup_{\partial X_J^{\varphi}} X_H^{\varphi}$ with twisted coefficients then yields the desired result.

LEMMA 3.3. *The Witt group class $\tau(L, \chi) \in L^0(\mathbb{Q}(\xi_{q^k})(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q]$ depends neither on the choice of 4-manifold W nor on the choice of isomorphism $\psi: \pi_1(T) \rightarrow \mathcal{H}$.*

Proof. To see that the object $\tau(L, \chi)$ as defined is independent of the choice of 4-manifold W , and so is an invariant, we need, by Novikov additivity for the rational intersection form and by Lemma 3.2 for twisted coefficients, to see that for a closed manifold

$$V := W \cup_{q^s M_L^{\varphi} \sqcup q^s T} -W',$$

the L -group classes

$$\lambda_{\mathbb{Q}(\xi_{q^k})(\mathcal{H})}(V) \text{ and } \lambda_{\mathbb{Q}}(V) \in L^0(\mathbb{Q}(\xi_{q^k})(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q]$$

coincide. This follows again from the Atiyah–Hirzebruch spectral sequence. More precisely, this follows from the argument of [Cha10, lemma 2.1], the only difference in our case is that $H_4(\mathbb{Z}_{q^k} \times \mathcal{H})$ is not zero, but is a finite q -group, and Cha’s argument thus implies that

$$\Omega_4(\mathbb{Z}_{q^k} \times \mathcal{H}) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q] \cong \Omega_4 \otimes_{\mathbb{Z}} \mathbb{Z}[1/q].$$

Therefore, modulo q -primary torsion, the twisted intersection form of V is Witt equivalent to the ordinary intersection form of V .

The obstruction does not depend on the choice of isomorphism $\psi : \pi_1(T) \rightarrow \mathcal{H}$: simply choose a different orientation preserving self-homeomorphism of T corresponding to the difference in two choices ψ and ψ' , and glue a product cobordism onto W for each copy of T in ∂W , using the new homeomorphism at the new boundary. We are using here the fact that any element of $SL(3, \mathbb{Z})$ can be realised by a homeomorphism of T , since $T \cong \mathbb{R}^3 / \mathbb{Z}^3$.

3.3. Main obstruction theorem

In order to prove Theorem 3.5 below, we will need the following lemma to justify having taken the trivial character on $\pi_1(T)$ to define $\tau(L, \chi)$.

LEMMA 3.4. *There is a cobordism of q^l copies of the 3-torus T with any character*

$$\chi : \pi_1(T) \xrightarrow{\cong} H_1(T; \mathbb{Z}) \longrightarrow \mathbb{Z}_{q^l},$$

via a cobordism W_χ which is depicted in Figure 1, to q^l copies of T with the trivial character, with an isomorphism $\psi : \pi_1(T) \rightarrow \mathcal{H}$ extending over W_χ . Moreover W_χ has trivial intersection form both with $\mathbb{Q}(\xi_{q^l})(\mathcal{H})$ -coefficients twisted using the extensions of ψ and χ , $\chi \times \psi : \pi_1(W_\chi) \rightarrow \mathbb{Z}_{q^l} \times \mathcal{H}$, and with rational coefficients.

Proof. First note that S^1 with a trivial character (S^1, tr) is cobordant on the one hand to q^l copies of S^1 , again with a trivial character, via a cobordism V_{tr} . On the other hand, (S^1, tr) is also cobordant to q^l copies of S^1 with the same non-trivial character $\chi : \pi_1(S^1) \rightarrow \mathbb{Z}_{q^l}$ on each copy, via a cobordism V_χ . If we glue together V_{tr} and V_χ , and cross the resulting cobordism with $S^1 \times S^1$, we obtain a cobordism from q^l copies of T to itself, with any character changed to the trivial character on the first S^1 of $S^1 \times S^1 \times S^1 = T$. That is, we define

$$W_\chi := (V_\chi \cup_{S^1} V_{\text{tr}}) \times S^1 \times S^1.$$

W_χ is just a union of several pairs of trousers (pants), crossed with a torus.

Part of the definition of a cobordism W_χ is a collection of homeomorphisms $T \xrightarrow{\cong} (\partial W_\chi)_i$ for each component $(\partial W_\chi)_i, i = 1, \dots, q^l$, of ∂W_χ with the non-trivial character associated to it. We pick a homeomorphism, which we use for each $i = 1, \dots, q^l, T \xrightarrow{\cong} (\partial W_\chi)_i$ such that the induced character $\chi : \pi_1((\partial W_\chi)_i) \rightarrow \mathbb{Z}_{q^l}$ factors through

$$\pi_1((\partial W_\chi)_i) \xrightarrow{\cong} \pi_1(S^1) \times \pi_1(S^1 \times S^1) \xrightarrow{\text{pr}} \pi_1(S^1) \cong \mathbb{Z} \longrightarrow \mathbb{Z}_{q^l}.$$

We therefore only need to change the character on one S^1 component of T in order to arrive at the trivial character associated to $(\partial W_\chi)_i$ for $i = q^l + 1, \dots, 2q^l$. This is precisely what W_χ achieves.

Since W_χ is a union of pairs of trousers, crossed with a torus, and since we have the same isomorphism $\psi : \pi_1(T) \xrightarrow{\cong} \mathcal{H}$ for each copy of T at the end of the cobordism with non-trivial character, then ψ extends over W_χ . If we require a different isomorphism ψ at the end of W_χ with trivial character, then, as in Lemma 3.3, we can glue on a product cobordism to each of the copies $(T, \text{tr} \times \psi) \subset \partial W_\chi$, of T with the trivial character, using a different self-homeomorphism of T .

The map $H_2(\partial W_\chi) \rightarrow H_2(W_\chi)$ is surjective. The non-singular part of the intersection form on $H_2(W_\chi)$ is therefore trivial. On the other hand, we have by the Eckmann–Shapiro lemma, that

$$\begin{aligned} H_2(W_\chi; \mathbb{Q}(\xi_{q^l})[\mathcal{H}]) &= H_2((V_\chi \cup_{S^1} V_{\text{tr}}) \times S^1 \times S^1; \mathbb{Q}(\xi_{q^l})[\mathcal{H}]) \\ &= H_2((V_\chi \cup_{S^1} V_{\text{tr}}); \mathbb{Q}(\xi_{q^l})[t^{\pm 1}]), \end{aligned}$$

but the last group is zero since $V_\chi \cup_{S^1} V_{tr}$ is homotopy equivalent to a 1-complex. It thus follows that the twisted intersection form on W_χ is also trivial.

The following is our main obstruction theorem. Note that we use the inclusion $\mathbb{Q}(\xi_{q^k}) \hookrightarrow \mathbb{C}$ in order to obviate dealing with the complications which arise from requiring higher prime powers to extend a character over a 4-manifold.

THEOREM 3.5. *Suppose that L is concordant to the Hopf link. Then for any admissible homomorphism $\varphi: H_1(M_L) \rightarrow A$ to a finite abelian p -group A , for a prime p , there exists a metaboliser $P = P^\perp$ for the linking form*

$$TH_1(X_L^\varphi, Y_a^\varphi) \times TH_1(X_L^\varphi, Y_a^\varphi) \longrightarrow \mathbb{Q}/\mathbb{Z}$$

with the following property: for any character of prime power order $\chi: H_1(M_L^\varphi) \rightarrow \mathbb{Z}_{q^k}$ which satisfies that $\chi|_{H_1(X_L^\varphi)}$ factors through

$$\chi|_{H_1(X_L^\varphi)}: H_1(X_L^\varphi) \longrightarrow H_1(X_L^\varphi, Y_a^\varphi) \xrightarrow{\delta} \mathbb{Z}_{q^k}$$

and that δ vanishes on P , we have that

$$\tau(L, \chi) = 0 \in L^0(\mathbb{C}(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q],$$

using the inclusion $\mathbb{Q}(\xi_{q^k}) \hookrightarrow \mathbb{C}$ to define the map $L^0(\mathbb{Q}(\xi_{q^k})(\mathcal{H})) \rightarrow \Lambda^0(\mathbb{C}(\mathcal{H}))$.

Proof. As above let E_C be the exterior of the concordance. The conclusion regarding the \mathbb{Q}/\mathbb{Z} -valued linking form was Proposition 2.3: a metaboliser is given by

$$P := \ker(TH_1(X^\varphi, Y_a^\varphi; \mathbb{Z}) \longrightarrow TH_1(E_C^\varphi, Y_a^\varphi; \mathbb{Z})).$$

Let χ satisfy the conditions set out. As in Section 2.4, let $W_C := E_C \cup X_H \times I$. By Propositions 2.10 and 2.12, we see that our character extends from $\pi_1(M_L^\varphi)$ to $\pi_1(W_C^\varphi)$, if we allow it to take values in \mathbb{Z}_{q^l} for a potentially higher prime power: $l \geq k$. We include $\mathbb{Q}(\xi_{q^k}) \subset \mathbb{C}$ so that our characters always extend.

Using the inverse of the inclusion induced isomorphism $H_1(M_L; \mathbb{Z}) \xrightarrow{\cong} H_1(W_C; \mathbb{Z})$, the map $\phi': \pi_1(M_L) \rightarrow H_1(M_L) \rightarrow \mathcal{H}'$ extends to

$$\phi': \pi_1(W_C) \longrightarrow H_1(W_C) \xrightarrow{\cong} H_1(M_L) \xrightarrow{\phi'} \mathcal{H}'.$$

As $\varphi: \pi_1(M_L) \rightarrow H_1(M_L) \rightarrow A$ extends to $\varphi: \pi_1(W_C) \rightarrow A$ using the same isomorphism $H_1(W_C) \xrightarrow{\cong} H_1(M_L)$, the restriction of ϕ' to

$$\ker(\varphi: \pi_1(W_C) \longrightarrow H_1(W_C) \xrightarrow{\cong} H_1(M_L) \longrightarrow A) \cong \pi_1(W_C^\varphi)$$

yields a homomorphism $\phi: \pi_1(W_C^\varphi) \rightarrow \mathcal{H}$, with $\mathcal{H} \subset \mathcal{H}'$ the image of $\phi: \pi_1(M_L^\varphi) \rightarrow \mathcal{H}'$ as above: since both $\phi: \pi_1(M_L^\varphi) \rightarrow \mathcal{H}$ and $\phi: \pi_1(W_C^\varphi) \rightarrow \mathcal{H}$ factor through $\ker(\varphi: H_1(M_L) \rightarrow A)$, we indeed have that

$$\text{im}(\phi: \pi_1(W_C^\varphi) \longrightarrow \mathcal{H}') \subseteq \text{im}(\phi: \pi_1(M_L^\varphi) \longrightarrow \mathcal{H}') = \mathcal{H}.$$

Since both χ and ϕ extend to W_C^φ , we can now use $(W_C^\varphi, \chi \times \phi)$ to calculate $\tau(L, \chi)$.

As χ extends to $\pi_1(W_C^\varphi)$, it also restricts to a map $\chi: \pi_1(T) \rightarrow \mathbb{Z}_{q^l}$, since $T \cong M_H^\varphi$. We need to know that for any character on $\pi_1(T) \cong \mathbb{Z}^3$, there is a cobordism from q^l copies of (T, χ) to q^l copies of (T, tr) , since we used the trivial character to define $\tau(L, \chi)$. This cobordism W_χ was described in Lemma 3.4. We take q^l copies of W_C , and glue them to W_χ :




Fig. 1. The cobordism with which we calculate $\tau(L, \chi)$ ($q^l = 3$).

this gives us a cobordism from $q^l \cdot M_L^\phi$ to $q^l \cdot T$, which we can then use to calculate $\tau(L, \chi)$: see Figure 1. We saw in Lemma 3.4 that the intersection forms of W_χ are trivial.

We therefore need to see that the twisted and ordinary Witt classes in $L^0(\mathbb{C}(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q]$ which arise from W_C^ϕ vanish.

We again apply Proposition 3.1, with $S = M_L$ and $Y = W_C$, to see, since $M_L \rightarrow W_C$ is a \mathbb{Z} -homology equivalence, that

$$H_*(W_C^\phi; \mathbb{Q}(\xi_{q^l})(\mathcal{H})) \cong H_*(M_L^\phi; \mathbb{Q}(\xi_{q^l})(\mathcal{H})) \cong 0.$$

Here we also use the conclusion of Lemma 3.2, so we are really applying Proposition 3.1 twice. Trivial homology groups necessarily carry Witt trivial forms; the rational intersection form is also trivial since

$$H_2(W_C^\phi; \mathbb{Q}) \subseteq \text{im}(H_2(M_L^\phi; \mathbb{Q}) \rightarrow H_2(W_C^\phi; \mathbb{Q}))$$

by Lemma 2.2 applied with $X = M_L$ and $E = W_C$. Therefore, including $\mathbb{Q}(\xi_{q^l}) \subset \mathbb{C}$, we have

$$\tau(L, \chi) = 0 \in L^0(\mathbb{C}(\mathcal{H})) \otimes_{\mathbb{Z}} \mathbb{Z}[1/q]$$

as claimed, since \mathbb{C} is flat over $\mathbb{Q}(\xi_{q^l})$.

4. Examples

4.1. Signature invariants

Let \mathcal{H} be a free-abelian group. In this section we will show how to recognise non-trivial elements in $L^0(\mathbb{C}(\mathcal{H}))$. Let P be a Hermitian matrix over $\mathbb{C}[\mathcal{H}]$. Let $\eta: \mathcal{H} \rightarrow S^1$ be any character. Note that η gives rise to a ring homomorphism $\mathbb{C}[\mathcal{H}] \rightarrow \mathbb{C}$. We denote by $\eta(P)$ the Hermitian matrix over \mathbb{C} obtained by applying the above ring homomorphism to all entries of P . We can then consider the signature $\text{sign}(\eta(P))$ of the matrix $\eta(P)$.

Let $\tau \in L^0(\mathbb{C}(\mathcal{H}))$. We represent τ by a Hermitian matrix P over $\mathbb{C}[\mathcal{H}]$ and we define

$$\sigma(P) := \int_{\eta \in \text{Hom}(\mathcal{H}, S^1)} \text{sign}(\eta(P)),$$

where we equip the torus $\text{Hom}(\mathcal{H}, S^1)$ with the Haar measure.

LEMMA 4.1. *The real number $\sigma(P)$ is independent of the choice of P representing a given $\tau \in L^0(\mathbb{C}(\mathcal{H}))$.*

Before proving Lemma 4.1, note that it allows us to define an invariant $\sigma(\tau) := \sigma(P)$ for some choice of representative P of τ . Finally given $\tau \otimes q \in L^0(\mathbb{C}(\mathcal{H})) \otimes \mathbb{Q}$ we define

$$\sigma(\tau \otimes q) = q \cdot \sigma(\tau).$$

Proof of Lemma 4.1. Since σ is additive it suffices to show that $\sigma : L^0(\mathbb{C}(\mathcal{H})) \rightarrow \mathbb{Z}$ is well-defined. Let $\tau \in L^0(\mathbb{C}(\mathcal{H}))$. Let P and P' be two Hermitian matrices over $\mathbb{C}[\mathcal{H}]$ which represent τ . We let

$$B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

and given $n \in \mathbb{N}$ we denote by nB the diagonal sum of n copies of B . It is well known that there exist n and n' such that $P \oplus nB$ and $P' \oplus n'B$ are congruent over $\mathbb{C}[\mathcal{H}]$, i.e. there exist matrices Q, Q' over $\mathbb{C}[\mathcal{H}]$ with non-zero determinant, such that

$$Q(P \oplus nB)Q^t = Q'(P' \oplus n'B)(Q')^t.$$

A priori the conjugation matrices Q are over $\mathbb{C}(\mathcal{H})$, but only one is required. We need two matrices Q and Q' here in order to ensure that both are over $\mathbb{C}[\mathcal{H}]$ rather than $\mathbb{C}(\mathcal{H})$. A straightforward calculation shows that

$$\begin{aligned} \text{sign}(\eta(P)) &= \text{sign}(\eta(Q(P \oplus nB)Q^t)) \\ &= \text{sign}(\eta(Q'(P' \oplus n'B)(Q')^t)) \\ &= \text{sign}(\eta(P')), \end{aligned}$$

for all η such that $\det(\eta(Q)) \neq 0$ and such that $\det(\eta(Q')) \neq 0$. Put differently, we have $\text{sign}(\eta(P)) = \text{sign}(\eta(P'))$ for all $\eta \in \text{Hom}(\mathcal{H}, S^1)$ outside a set of measure zero.

4.2. The satellite construction

Let L be a link, let K be a knot and let $\gamma \subset X_L$ be a curve which is unknotted in S^3 . Define $X_K := S^3 \setminus \nu K$. We form the space

$$S^3 \setminus \nu\gamma \cup_{\theta} X_K \cong S^3,$$

where $\theta: \partial(S^3 \setminus \nu\gamma) \rightarrow \partial X_K$ is a homeomorphism which sends the meridian of γ to the zero-framed longitude of K and the zero-framed longitude of γ to a meridian of K . The image of L is the *satellite link* $S(L, K, \gamma)$. When L, K and γ are understood, then we just write $S = S(L, K, \gamma)$.

Note that the canonical degree one map $S^3 \setminus \nu K \rightarrow S^1 \times D^2$ gives rise to a canonical degree one map $S^3 \setminus \nu S \rightarrow S^3 \setminus \nu L$. Let $\varphi: H_1(X_L; \mathbb{Z}) \rightarrow A$ be an epimorphism onto a finite group. We denote the induced homomorphism $H_1(X_S; \mathbb{Z}) \xrightarrow{\cong} H_1(X_L; \mathbb{Z}) \rightarrow A$ by φ as well. We now have the following lemma, which says that in many cases the satellite construction has no effect on the homology of finite covers or on the linking form.

LEMMA 4.2. *Let L, K and γ be as above such that in addition γ is null homologous. Let $\varphi: H_1(X_L; \mathbb{Z}) \rightarrow A$ be an epimorphism onto a finite group. Then the canonical degree one map $X_S \rightarrow X_L$ induces an isomorphism*

$$TH_1(X_S^\varphi, Y_a^\varphi) \xrightarrow{\cong} TH_1(X_L^\varphi, Y_a^\varphi)$$

and the linking form of Section 2.2 is preserved.

Proof. We write $k = |A|$. Since γ is null homologous, it lifts to curves $\tilde{\gamma}_1, \dots, \tilde{\gamma}_k \subset X_L^\varphi$. Then

$$X_S^\varphi \cong \text{cl}(X_L^\varphi \setminus \bigcup_{i=1}^k \nu \tilde{\gamma}_i) \cup_{\bigcup_{i=1}^k \partial X_K} \bigcup_{i=1}^k X_K.$$

We now refer to the argument of [Fri03, lemma 10.3], the ideas of which originate from [Lit84]. This argument carries over to our situation with translation of notation and, noting that $H_1(X_L^\varphi) \xrightarrow{\cong} H_S(X_L^\varphi)$, implies that

$$H_1(X_L^\varphi, Y_a^\varphi) \xrightarrow{\cong} H_1(X_S^\varphi, Y_a^\varphi)$$

by the five lemma, and the fact that the boundary of X_S is the image of the boundary of X_L in the satellite construction.

The key technical advantage of the satellite construction is that it allows us to calculate $\sigma(\tau(S, \chi))$ in terms of $\tau(L, \chi)$ and the Levine–Tristram signatures of K . Recall that given any knot K and $\omega \in S^1$ the Levine–Tristram signature $\sigma(K, \omega) \in \mathbb{Z}$ is defined and that it can be calculated using any Seifert matrix of K . It is well known that $\sigma(K, \omega) = 0$ for any slice knot K and any prime power root of unity. We refer to [Lev69] and [Tri69] for details.

The following theorem can easily be proved using the ideas and methods of Litherland [Lit84]; see also [Fri03, theorem 10.5]. Similar results have been obtained by many authors over the years, and we will therefore not provide a proof.

THEOREM 4.3. *Let L be a 2-component link with linking number one and let γ be a null-homologous curve in $S^3 \setminus \nu L$ which is unknotted in S^3 . Let K be a knot. We write $S := S(L, K, \gamma)$. Let $\varphi: H_1(X_L; \mathbb{Z}) \rightarrow A$ be an epimorphism onto a finite group of order k and let*

$$\chi: TH_1(X_S^\varphi, Y_a^\varphi) \xrightarrow{\cong} TH_1(X_L^\varphi, Y_a^\varphi) \longrightarrow S^1$$

be a character and let $\phi: H_1(M_S^\varphi) \xrightarrow{\cong} H_1(M_L^\varphi) \rightarrow \mathcal{H}$ be an epimorphism onto a free abelian group of rank 3. Suppose that γ lifts to $\tilde{\gamma}_1, \dots, \tilde{\gamma}_k$. Then

$$\sigma(\tau(S, \chi)) = \sigma(\tau(L, \chi)) + \sum_{i=1}^k \sigma(K, \omega_i),$$

where $\omega_i = \chi([\tilde{\gamma}_i])$ and $[\tilde{\gamma}_i]$ is the image of $\tilde{\gamma}_i$ in $TH_1(X_L^\varphi, Y_a^\varphi)$.

4.3. Example

We consider the link \mathcal{L} depicted in Figure 2. Note that \mathcal{L} is concordant to the Hopf link via a ribbon move. It has non-trivial multi-variable Alexander polynomial

$$\Delta_{\mathcal{L}}(s, t) = (ts + 1 - s)(ts + 1 - t),$$

which we computed using a Wirtinger presentation of the link group and the Fox calculus.




Fig. 2. A link \mathcal{L} which is concordant to the Hopf link, with non-trivial Alexander polynomial.

We take φ to be the admissible homomorphism given by two projections $\varphi: H_1(X_{\mathcal{L}}; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2$. Using the theorem in the appendix we can calculate that $TH_1(X_{\mathcal{L}}^{\varphi}, Y_a^{\varphi})$ of order 9. In fact we have a Maple program which calculates the homology explicitly. It starts with the Wirtinger presentation of $\pi_1(X_{\mathcal{L}})$, and uses it to construct the boundary maps in the chain complex $C_*(X_{\mathcal{L}}^{\varphi}; \mathbb{Z})$, by using the regular representation at the end of the composition:

$$\pi_1(X_{\mathcal{L}}) \longrightarrow H_1(X_{\mathcal{L}}) \cong \mathbb{Z} \oplus \mathbb{Z} \longrightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2 \longrightarrow \text{Aut}(\mathbb{Z}[\mathbb{Z}_2 \oplus \mathbb{Z}_2]) \longrightarrow \text{GL}(4, \mathbb{Z}),$$

which sends

$$(1, 0) \mapsto \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \text{ and } (0, 1) \mapsto \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

By doing row and column operations simultaneously on matrices representing the boundary maps $\partial_2: C_2 \rightarrow C_1$ and $\partial_1: C_1 \rightarrow C_0$, we directly compute that

$$H_1(X_{\mathcal{L}}^{\varphi}; \mathbb{Z}) \cong \mathbb{Z}_9 \oplus \mathbb{Z} \oplus \mathbb{Z},$$

which by Corollary 2.8 implies that:

$$TH_1(X_{\mathcal{L}}^{\varphi}, Y_a^{\varphi}; \mathbb{Z}) \cong \mathbb{Z}_9.$$

Using our Maple program we augment the matrix representing ∂_1 with an identity matrix, whose task is to keep track of the basis changes. With this extra information we can relate a generator of the \mathbb{Z}_9 summand of $TH_1(X_{\mathcal{L}}^{\varphi}, Y_a^{\varphi})$ to our original Wirtinger generators, which we understand geometrically and can exhibit in the diagram of \mathcal{L} . Using this approach we obtain the curve γ in Figure 3 which has the following three properties:

- (i) γ represents a generator of $TH_1(X_{\mathcal{L}}^{\varphi}, Y_a^{\varphi})$;
- (ii) there exists a Seifert surface F for \mathcal{L} , such that γ and F do not intersect (given by Seifert’s algorithm and tubing along γ); and
- (iii) the curve γ forms the unlink with either of the two components of L .

Claim. Let $K = 3_1$ be the trefoil knot. Then $S = S(\mathcal{L}, K, \gamma)$ is not concordant to the Hopf link.

Proof. Recall that the degree one map $X_S \rightarrow X_L$ induces an isomorphism $H_1(X_S; \mathbb{Z}) \rightarrow H_1(X_L; \mathbb{Z})$. We denote by φ the induced homomorphism $H_1(X_S; \mathbb{Z}) \rightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2$. Since




Fig. 3. (Colour online) The link \mathcal{L} with the curve γ shown.

$\gamma \subset X_L$ is null-homologous it follows from the previous section that the induced map

$$H_1(X_S^\varphi, Y_a^\varphi; \mathbb{Z}) \longrightarrow H_1(X_L^\varphi, Y_a^\varphi; \mathbb{Z})$$

is an isomorphism and in fact this map induces an isometry of linking forms. Note that any form $\mathbb{Z}_9 \times \mathbb{Z}_9 \rightarrow \mathbb{Q}/\mathbb{Z}$ has a unique metaboliser.

Now let $\chi : TH_1(X_S^\varphi, Y_a^\varphi; \mathbb{Z}) \rightarrow \mathbb{Z}_3 \rightarrow S^1$ be a non-trivial character which vanishes on the unique metaboliser of the linking pairing on $TH_1(X_S^\varphi, Y_a^\varphi; \mathbb{Z}) \cong \mathbb{Z}_9$ and let $\phi : H_1(M_S^\varphi) \rightarrow \mathcal{H}$ an epimorphism onto a free abelian group of rank 3. By Theorem 3.5 and Lemma 4.1 we are done once we show that $\sigma(\tau(S, \chi)) \neq 0$.

First note that χ and ϕ are induced from a character on $TH_1(X_L^\varphi, Y_a^\varphi; \mathbb{Z})$ and an epimorphism on $H_1(M_L^\varphi)$, which we denote with the same symbols. We denote by $\tilde{\gamma}_1, \dots, \tilde{\gamma}_4 \subset X_L^\varphi$ the preimages of γ . It follows from Theorem 4.3 that

$$\sigma(\tau(S, \chi)) = \sigma(\tau(L, \chi)) + \sum_{i=1}^4 \sigma(K, \omega_i),$$

where $\omega_i = \chi([\tilde{\gamma}_i])$ and $[\tilde{\gamma}_i]$ is the image of $\tilde{\gamma}_i$ in $TH_1(X_L^\varphi, Y_a^\varphi)$. Note that $\sigma(\tau(L, \chi)) = 0$ since L is concordant to the Hopf link and since $\chi : TH_1(X_L^\varphi, Y_a^\varphi; \mathbb{Z}) \rightarrow S^1$ also vanishes on the unique metaboliser of $TH_1(X_L^\varphi, Y_a^\varphi; \mathbb{Z}) \cong \mathbb{Z}_9$. Furthermore note that any ω_i is a third root of unity, and moreover at least one ω_i is a non-trivial third root of unity since γ represents a generator of $TH_1(X_L^\varphi, Y_a^\varphi)$. It is well known that $\sigma(K, 1) = 0$ and $\sigma(K, e^{\pm 2\pi i/3}) = 2$. We thus see that $\sigma(\tau(S, \chi)) \neq 0$.

4.4. Comparison with previous invariants

We now slightly modify the example of the previous section. By the proof of [Fri04, proposition 7.4] there exists a knot K such that $\int_{z \in S^1} \sigma(K, z) = 0$ but such that $\sigma(K, e^{2\pi i/3}) = 12$ and $\sigma(K, e^{4\pi i/3}) = 12$. We denote by S the satellite link which is constructed as in the previous section, with the sole difference that we replace the trefoil knot by the knot K . The argument of the previous section shows that our obstruction detects that S is not concordant to the Hopf link.

We will now quickly discuss the earlier obstructions to link concordance which are known to the authors. The enumeration here does not correspond exactly with the enumeration of link concordance obstructions which appears in the introduction.

- (i) The components of S and H are trivial knots, in particular components of S do not give rise to non-trivial concordance obstructions.
- (ii) The linking number of the two components of S is 1, which is the same as the linking number of the Hopf link. Note that for non-trivial linking numbers the higher Milnor $\overline{\mu}$ -invariants are not defined; in particular they do not give further concordance obstructions.
- (iii) A variation the proof of Lemma 4.2 shows that $\Delta_S = \Delta_{\mathcal{L}} \in \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$. It follows that

$$\begin{aligned} \Delta_S &= (ts + 1 - s)(ts + 1 - t) \\ &= s^{-1}t^{-1}(ts + 1 - s)(t^{-1}s^{-1} + 1 - s^{-1}) \in \mathbb{Z}[s^{\pm 1}, t^{\pm 1}] \end{aligned}$$

factors as a norm over $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$. In particular the link concordance obstructions of Murasugi [Mur67], Kawauchi [Kaw77] [Kaw78, theorem B] and Nakagawa [Nak78] are satisfied.

- (iv) Since the curve γ does not intersect a Seifert surface F of \mathcal{L} we see that the image of F in

$$S^3 \setminus v\gamma \cup_{\theta} X_K \cong S^3$$

is a Seifert surface for S . One can easily see that the corresponding Seifert matrices for \mathcal{L} and S agree. Since \mathcal{L} is concordant to the Hopf link it is now straightforward to see that the link concordance obstructions of Tristram [Tri69] are satisfied.

- (v) Let \mathbb{T} be one of the two boundary components of X_S . It follows from Stallings’s theorem (see [Sta65]) applied to the homology equivalence $\mathbb{T} \rightarrow \pi_1(X_S)$, that any epimorphism from $\pi_1(X_S)$ onto a nilpotent group (in particular a p -group) factors through the abelianisation of $\pi_1(X_S)$. It follows from standard arguments that all the twisted Alexander polynomials of S corresponding to representations factoring through groups of prime power order are in fact determined by Δ_S . Since Δ_S factors as a norm over $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$ it is now straightforward to verify that the link concordance obstruction of [CF10] vanishes.
- (vi) The fact that $\pi_1(X_S)$ has no non-abelian nilpotent quotients also implies that the signature invariants by Cha and Ko [CK99b, CK06], Friedl [Fri05], Levine [Lev94] and Smolinsky [Smo89] are determined by signature invariants corresponding to abelian representations. The results of Litherland [Lit84] can easily be modified to show that such signature invariants of S agree with the corresponding signature invariants of \mathcal{L} , which do not give any obstructions to being concordant to H , since \mathcal{L} is concordant to H .
- (vii) Similarly to the previous invariants one can also show that the obstructions of [Cha10, theorem 1.4] vanish in our case.
- (viii) Let $\mathcal{L}_1 \cup \mathcal{L}_2 \subset S^3$ be a 2-component link with linking number one. We can consider the k -fold branched cover $Y := Y(\mathcal{L}_1, k)$ of S^3 branched along \mathcal{L}_1 . The preimage of \mathcal{L}_2 under the projection map $Y \rightarrow S^3$ consists of one component which we denote by J (here we use that the linking number is one). If k is a prime power, then Y is a rational homology sphere. If \mathcal{L} is concordant to the Hopf link and if k is a prime

power, then it is straightforward to see that J is rationally concordant to the unknot. One can therefore apply the obstructions of a knot being rationally concordant to the unknot. Such obstructions are given by Levine–Tristram signatures (see Cha and Ko [CK02]) and by Cochran–Orr–Teichner type Von Neumann ρ -invariants (see [Cha07], [COT03]). In our case it is straightforward to see that the preimage of one component of the satellite link is the satellite of a slice knot such that the infection curve is null-homologous. It follows from standard arguments (see above) that the Levine–Tristram signatures are zero. Finally, since also $\int_{z \in S^1} \sigma(K, z) = 0$, it follows that the obstructions of [Cha07] cannot be used to show that the knot is not rationally slice.

- (ix) As mentioned in the introduction, blowing down along one of the link components yields a knot in S^3 to which we can apply knot concordance obstructions, not just rational concordance obstructions (see [CKRS10, corollary 2.4]). It seems likely that Casson–Gordon obstructions of this knot will also obstruct S from being concordant to the Hopf link. In our example, blowing down \mathcal{L} along the left hand component in the diagram (Figure 3) yields the slice knot 6_1 . The 2-fold branched cover of S^3 along 6_1 , which we denote by $L_2^{6_1}$, has homology $H_1(L_2^{6_1}; \mathbb{Z}) \cong \mathbb{Z}_9$, so we also take a non-trivial character $\chi: \mathbb{Z}_9 \rightarrow \mathbb{Z}_3$. Keeping track of the curve γ , one then needs our character on $H_1(L_2^{6_1}; \mathbb{Z})$ to map the image of γ to a non-zero element of \mathbb{Z}_3 . One way to check that this is possible would be to use our methods of the previous section; we have not done the calculation. Litherland’s formula [Lit84] for Casson–Gordon obstructions would then show that they do not vanish, thus giving another proof that S is not concordant to H .

In our obstruction theory, we make use of the fact that \mathcal{H} has rank greater than one when we take finite covers using maps $\varphi: H_1(X_L) \rightarrow A$ which are non-trivial on the meridians of both link components. In order to make use of the full power of our obstruction theory, we need an example which also makes use of the fact that \mathcal{H} has rank greater than one in the non-triviality of the Witt class $\tau(S, \chi)$ i.e. the fact that we define an obstruction in $L^0(\mathbb{C}(\mathcal{H}))$ and not just in $L^0(\mathbb{C}(t))$. Such examples do not arise from satellite constructions.

Added in proof: on the other hand, as mentioned in the introduction, Min Hoon Kim informs us that he has verified the conjecture that the obstructions of this paper vanishes for links which are height (3.5) Whitney tower/Grope concordant to the Hopf link. Therefore our satellite links are not height (3.5) concordant to the Hopf link, but the blow down technique cannot be used to prove this.

Appendix: homology of finite abelian covers

It is well known that the order of the homology of a cyclic cover of a knot complement can be computed from the Alexander polynomial. More precisely, let K be a knot, let $\Delta_K(t) \in \mathbb{Z}[t^{\pm 1}]$ be its Alexander polynomial, let $n \in \mathbb{N}$ and denote by X_K^n the n -fold cyclic cover of X_K . Then Fox [Fox56] (see also [Web79] and [Tur86, section 1.9]) showed that $H_1(X_K^n) \cong \mathbb{Z} \oplus G$, where G is a group which satisfies

$$|G| = \prod_{k=0}^{n-1} \Delta_K(e^{2\pi ik/n}). \tag{4.1}$$

Here we denote by $|G|$ the order of G . We use the convention that $|G| = 0$ if G is infinite.

In general a naïve extension of the above formula to the case of finite abelian covers of links does not hold. We refer to [MM82], [Sak95], [HS97] and [Por04] for results relating the size of the homology of finite abelian (branched) covers of link complements to evaluations of the multivariable Alexander polynomial and its sublinks.

In this paper we are only interested in 2–component links with linking number one. One of the guiding themes of this paper is that such links share many formal properties with knots, in particular many statements about abelian invariants of knots can be generalised fairly easily to such links. In this appendix we will give an elementary proof that the analogue of (4.1) holds in our case.

Before we state our theorem we have to fix some notation. Let L be an oriented 2–component link. We denote by s and t the canonical elements in $H_1(X_L; \mathbb{Z})$ corresponding to the meridians. If $\varphi: H_1(X_L) \rightarrow A$ is a homomorphism and $\eta: A \rightarrow S^1$ a character, then we can consider $\eta(\varphi(s)) \in S^1$ and $\eta(\varphi(t)) \in S^1$. We now have the following theorem.

Theorem. *Let L be an oriented 2–component link with linking number one. Let $\varphi: H_1(X_L) \rightarrow A$ be an epimorphism onto a finite abelian group. Then $H_1(X_L^\varphi) \cong \mathbb{Z}^2 \oplus G$, where G is a group which satisfies*

$$|G| = \prod_{\eta \in \text{Hom}(A, S^1)} \Delta_L(\eta(\varphi(s)), \eta(\varphi(t))).$$

Here $\Delta_L(s, t)$ denotes the Alexander polynomial of L .

Proof. We can view X_L as a finite CW complex. We denote by Y one of the two boundary components of X_L . We can collapse X_L onto a 2–complex X which contains Y and such that all 0–cells lie in Y . We denote by r the number of 1–cells in $X \setminus Y$. Note that the Euler characteristic of X and Y is zero; it follows that the number of 2–cells in $X \setminus Y$ also equals r . Furthermore note that the inclusions also induce isomorphisms $H_1(Y; \mathbb{Z}) \rightarrow H_1(X; \mathbb{Z}) \rightarrow H_1(X_L; \mathbb{Z})$. Also note that $H_1(Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) = 0$. It follows from the long exact sequence of $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$ –homology groups of the pair (X, Y) , that

$$H_1(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) \cong H_1(X; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]),$$

and the latter is of course isomorphic to $H_1(X_L; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}])$. Since X is a 2–complex and since Y contains all 0–cells of X we obtain the following exact sequence of $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$ –modules:

$$\begin{array}{ccccccc} 0 & & \longrightarrow & H_2(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) & & & \\ \longrightarrow & C_2(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) & \longrightarrow & C_1(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) & & & \\ \longrightarrow & H_1(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) & \longrightarrow & 0. & & & \end{array}$$

Note that the boundary map

$$C_2(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) \longrightarrow C_1(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}])$$

is represented by an $r \times r$ –matrix P . Furthermore note that by definition of the Alexander polynomial, and since $H_1(X, Y; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}]) \cong H_1(X_L; \mathbb{Z}[s^{\pm 1}, t^{\pm 1}])$ we have $\det(P) = \Delta_L$.

Now let $\varphi: H_1(X_L; \mathbb{Z}) \rightarrow A$ be an epimorphism onto a finite abelian group. We write $k = |A|$. We denote the resulting epimorphism $H_1(X; \mathbb{Z}) \cong H_1(X_L; \mathbb{Z}) \xrightarrow{\varphi} A$ by φ as well. We also denote by φ the induced map $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}] = \mathbb{Z}[H_1(X_L; \mathbb{Z})] \rightarrow \mathbb{Z}[A]$ and finally we denote by $\varphi(P)$ the $r \times r$ –matrix given by applying φ to each entry of P . We then obtain

the following exact sequence of $\mathbb{Z}[A]$ -modules:

$$\begin{array}{ccccccc} 0 & & \longrightarrow & H_2(X, Y; \mathbb{Z}[A]) & & & \\ & \longrightarrow & C_2(X, Y; \mathbb{Z}[A]) & \xrightarrow{\partial} & C_1(X, Y; \mathbb{Z}[A]) & & \\ & \longrightarrow & H_1(X, Y; \mathbb{Z}[A]) & \longrightarrow & 0, & & \end{array}$$

where the boundary map is presented by $\varphi(P)$.

We now pick an isomorphism $\mathbb{Z}[A] \rightarrow \mathbb{Z}^k$ of free \mathbb{Z} -modules. Note that the action of $H_1(X_L; \mathbb{Z})$ on $\mathbb{Z}[A]$ given by φ and by left multiplication of A on $\mathbb{Z}[A]$ gives rise to a representation $H_1(X_L; \mathbb{Z}) \rightarrow \text{GL}(k, \mathbb{Z})$, which we also denote by φ . Note that this map extends to a ring homomorphism $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}] \rightarrow \mathbb{Z}[H_1(X_L; \mathbb{Z})] \rightarrow M(k, \mathbb{Z})$ which we also denote by φ . Finally we denote by $\varphi(P)$ the $kr \times kr$ -matrix over \mathbb{Z} given by applying the above ring homomorphism to each entry of $\varphi(P)$. It follows, from the above, that

$$|H_1(X, Y; \mathbb{Z}[A])| = \det(\varphi(P)).$$

It is well known that for any finite abelian group B the regular representation $B \rightarrow \text{Aut}(\mathbb{C}[B]) = \text{GL}(|B|, \mathbb{C})$ is conjugate to the representation which is given by the direct sum of all characters $\eta: B \rightarrow S^1$. It now follows that

$$\begin{aligned} |H_1(X, Y; \mathbb{Z}[A])| &= \det(\varphi(P)) \\ &= \prod_{\eta \in \text{Hom}(A, S^1)} \det(\eta(P)) \\ &= \prod_{\eta \in \text{Hom}(A, S^1)} \eta(\det(P)) \\ &= \prod_{\eta \in \text{Hom}(A, S^1)} \eta(\Delta_L(s, t)) \\ &= \prod_{\eta \in \text{Hom}(A, S^1)} \Delta_L(\eta(s), \eta(t)). \end{aligned}$$

Now note that we have an exact sequence

$$\begin{array}{ccccccc} & & & & & & H_2(X, Y; \mathbb{Z}[A]) \\ \longrightarrow & H_1(Y; \mathbb{Z}[A]) & \longrightarrow & H_1(X; \mathbb{Z}[A]) & \longrightarrow & H_1(X, Y; \mathbb{Z}[A]) & \\ \longrightarrow & H_0(Y; \mathbb{Z}[A]) & \longrightarrow & H_0(X; \mathbb{Z}[A]) & & & \end{array}$$

Note that the last two groups are isomorphic to \mathbb{Z} and the last map is an isomorphism. We now consider the following commutative diagram

$$\begin{array}{ccc} H_1(Y; \mathbb{Z}[A]) & \longrightarrow & H_1(X; \mathbb{Z}[A]) \\ \downarrow & & \downarrow \\ H_1(Y) & \longrightarrow & H_1(X; \mathbb{Z}), \end{array}$$

where the vertical maps are induced by the surjection from the φ -cover to the base space. Note that the left hand vertical map is injective since Y is a torus, and the bottom horizontal map is injective since L is a 2-component link with linking number one. It follows that the top horizontal map is also injective. Finally note that $H_1(Y; \mathbb{Z}[A]) \cong \mathbb{Z}^2$. Combining all of the above we obtain a short exact sequence

$$0 \longrightarrow \mathbb{Z}^2 \longrightarrow H_1(X; \mathbb{Z}[A]) \longrightarrow H_1(X, Y; \mathbb{Z}[A]) \longrightarrow 0.$$

The above calculation of $|H_1(X, Y; \mathbb{Z}[A])|$ implies the theorem.

Acknowledgments. This research was mostly conducted while MP was a visitor at the Max Planck Institute for Mathematics in Bonn, which he would like to thank for its generous hospitality.

We furthermore would like to thank Jae Choon Cha, Jim Davis, Jonathan Hillman, Kent Orr, Danny Ruberman, Dan Silver and Lorenzo Traldi for helpful conversations, comments and feedback.

REFERENCES

- [CF10] J. C. CHA and S. FRIEDL. Twisted torsion invariants and link concordance. To appear in Forum Math., Preprint: <http://arxiv.org/abs/1001.0926>, (2010).
- [CG86] A. CASSON and C. MCA. GORDON. Cobordism of classical knots. In *A la Recherche de la Topologie Perdue Progr. Math.* vol. 62 (Birkhauser Boston, 1986), pages 181–199.
- [Cha07] J. C. CHA. The structure of the rational concordance group of knots. *Mem. Amer. Math. Soc.* **189** (885) (2007). x+95.
- [Cha10] J. C. CHA. Link concordance, homology cobordism, and Hirzebruch-type defects from iterated p -covers. *J. Eur. Math. Soc.* **12** (2010), 555–610.
- [Cha12] J. C. CHA. Symmetric Whitney tower cobordism for bordered 3–manifolds and links. arXiv:1204.4968 [math.GT] (2012).
- [CK99a] J. C. CHA and K. H. KO. On equivariant slice knots. *Proc. Amer. Math. Soc.* **127** (7) (1999), 2175–2182.
- [CK99b] J. C. CHA and K. H. KO. Signature invariants of links from irregular covers and non-abelian covers. *Math. Proc. Camb. Philos. Soc.* **127** (1) (1999), 67–81.
- [CK02] J. C. CHA and K. H. KO. Signatures of links in rational homology spheres. *Topology* **41** (2002), 1161–1182.
- [CK06] J. C. CHA and K. H. KO. Signature invariants of covering links. *Trans. Amer. Math. Soc.* **358** (2006), 3399–3412.
- [CK08] J. C. CHA and T. KIM. Covering link calculus and iterated Bing doubles. *Geom. Topol.* **12** (2008), 2172–2201.
- [CKRS10] J. C. CHA, T. KIM, D. RUBERMAN and S. STRLE. Smooth concordance of links topologically concordant to the Hopf link. arXiv:1012.2045v1[math.GT] (2010).
- [CLR08] J. C. CHA, C. LIVINGSTON and D. RUBERMAN. Algebraic and Heegaard–Floer invariants of knots with slice Bing doubles. *Math. Proc. Camb. Phil. Soc.* **144** (2008), 403–410.
- [CO90] T. D. COCHRAN and K. E. ORR. Not all links are concordant to boundary links. *Bulletin Amer. Math. Soc.* **23** (1) (1990), 99–106.
- [CO93] T. D. COCHRAN and K. E. ORR. Not all links are concordant to boundary links. *Ann. Math.* **138** (1993), 519–554.
- [CO09] J. C. CHA and K. E. ORR. $L^{(2)}$ -signatures, homology localization and amenable groups. To appear in *Comm. Pure Appl. Math.* Preprint: arXiv:09103.3700 (2009).
- [COT03] T. D. COCHRAN, K. E. ORR and P. TEICHNER. Knot concordance, Whitney towers and $L^{(2)}$ signatures. *Ann. of Math.* (2) **157** (2003), no. 2, 433–519.
- [CST12] J. CONANT, R. SCHNEIDERMAN and P. TEICHNER. Whitney tower concordance of classical links. *Geom. Topol.* **16** (2012), 1419–1479.
- [Dav06] J. DAVIS. A two component link with Alexander polynomial one is concordant to the Hopf link. *Math. Proc. Camb. Phil. Soc.* **140** (2006), no. 2, 265–268.
- [Fox56] R. H. FOX. Free differential calculus. III. Subgroups. *Ann. of Math.* (2) **64** (1956), 407–419.
- [FP12] S. FRIEDL and M. POWELL. An injectivity theorem for Casson–Gordon type representations relating to the concordance of knots and links. *Bull. Korean Math. Soc.* **49** (2012), 395–409.
- [FQ90] M. FREEDMAN and F. QUINN. *Topology of Four Manifolds* (Princeton University Press, 1990).
- [Fre84] M. H. FREEDMAN. The disk theorem for four-dimensional manifolds. *Proc. Internat. Congr. Math.* vol. 1, 2 (Warsaw, 1983), PWN (1984), pages 647–663.
- [Fri03] S. FRIEDL. *Eta invariants as sliceness obstructions and their relation to Casson–Gordon invariants*. PhD thesis, Brandeis University (2003).
- [Fri04] S. FRIEDL. Eta invariants as sliceness obstructions and their relation to Casson–Gordon invariants. *Alg. Geom. Topol.* **4** (2004), 893–934.
- [Fri05] S. FRIEDL. Link concordance, boundary link concordance and eta-invariants. *Math. Proc. Camb. Phil. Soc.* **138** (2005), no. 3, 437–460.
- [Har08] S. HARVEY. Homology cobordism invariants and the Cochran–Orr–Teichner filtration of the link concordance group. *Geom. Topol.* **12** (2008), no. 1, 387–430.
- [Hil02] J. A. HILLMAN. *Algebraic Invariants of Links*. Series on Knots and Everything **32** (World Scientific Publishing Co. 2002).
- [HS97] J. A. HILLMAN and M. SAKUMA. On the homology of finite abelian coverings of links. *Canad. Math. Bull.* **40** (1997), 309–315.

- [Kaw77] A. KAWAUCHI. On quadratic forms of 3-manifolds. *Invent. Math.* **43** (1977), 177–198.
- [Kaw78] A. KAWAUCHI. On the Alexander polynomials of cobordant links. *Osaka J. Math.* **15** (1978), no. 1, 151–159.
- [Let00] C. F. LETSCHE. An obstruction to slicing knots using the eta invariant. *Math. Proc. Camb. Phil. Soc.* **128** (2) (2000), 301–319.
- [Lev69] J. LEVINE. Knot cobordism groups in codimension two. *Comment. Math. Helv.* **44** (1969), 229–244.
- [Lev94] J. LEVINE. Link invariants via the eta invariant. *Comment. Math. Helv.* **69** (1994), 82–119.
- [Lev07] J. LEVINE. Concordance of boundary links. *J. Knot Theory Ramifications* **16** (2007), no. 9, 1111–1120.
- [Lit84] R. A. LITHERLAND. Cobordism of satellite knots. In *Four-manifold theory (Durham, N.H., 1982)*, volume 35 of *Contemp. Math.*, (Amer. Math. Soc. 1984) pages 327–362.
- [Mil57] J. W. MILNOR. Isotopy of links. *Alg. Geom. Top.* A symposium in honor of S. Lefschetz, (1957), 280–306.
- [MM82] J. P. MAYBERRY and K. MURASUGI. Torsion-groups of abelian coverings of links. *Trans. Amer. Math. Soc.* **271** (1982), 143–173.
- [Mur67] K. MURASUGI. On a certain numerical invariant of link types. *Trans. Amer. Math. Soc.* **117** (1967), 387–422.
- [Nak78] Y. NAKAGAWA. On the Alexander polynomials of slice links. *Osaka J. Math.* **15** (1978), no. 1, 161–182.
- [Por04] J. PORTI. Mayberry–Murasugi’s formula for links in homology 3-spheres. *Proc. Amer. Math. Soc.* **132** (2004), 3423–3431.
- [Sak95] M. SAKUMA. Homology of abelian coverings of links and spatial graphs. *Canad. J. Math.* **47** (1995), 201–224.
- [Smo89] L. SMOLINSKY. Invariants of link cobordism. *Proceedings of the 1987 Georgia Topology Conference (Athens, GA, 1987)*. *Topology Appl.* **32** (1989), 161–168.
- [Sta65] J. STALLINGS. Homology and central series of groups. *J. Algebra* **2** (1965), 170–181.
- [Tri69] A. G. TRISTRAM. Some cobordism invariants for links. *Proc. Camb. Phil. Soc.* **66** (1969), 251–264.
- [Tur86] V. TURAEV. Reidemeister torsion in knot theory. *Russian Math. Surveys* **41** (1986), 119–182.
- [Web79] C. WEBER. Sur une formule de R. H. Fox concernant l’homologie des revêtements cycliques. *Enseign. Math.* (2) **25** (1979), 261–272.