

**Regensburger
DISKUSSIONSBEITRÄGE
zur Wirtschaftswissenschaft**

University of Regensburg Working Papers in Business,
Economics and Management Information Systems

**Resource Allocation Heuristics
for Unknown Sales Response Functions
with Additive Disturbances**

Daniel Gahler* and Harald Hruschka*

November 17, 2016

Nr. 488

JEL Classification: M30 ; C61 ; C63

Key Words: Marketing Resource Allocation; Exploration-Exploitation Algorithm;
Monte Carlo Simulation; Optimization

* Marketing Department, Faculty of Business, Economics and Management Information Systems , University of Regensburg, 93040 Regensburg, Germany, Phone: +49-941-943-2279, E-mail: daniel.gahler@wiwi.uni-r.de

Resource Allocation Procedures for Unknown Sales Response Functions with Additive Disturbances

Daniel Gahler, Harald Hruschka

Abstract

We develop an exploration-exploitation algorithm which solves the allocation of a fixed resource (e.g., a budget, a sales force size, etc.) to several units (e.g., sales districts, customer groups, etc.) with the objective to attain maximum sales. This algorithm does not require knowledge of the form of the sales response function and is also able cope with additive random disturbances. The latter as a rule are a component of sales response functions estimated by econometric methods. We compare the algorithm to three rules of thumb which in practice are often used for this allocation problem. The comparison is based on a Monte Carlo simulation for five replications of 192 experimental constellations, which are obtained from four function types, four procedures (i.e., the three rules of thumb and the algorithm), similar/varied elasticities, similar/varied saturations, and three error levels. A statistical analysis of the simulation results shows that the algorithm performs better than the three rules of thumb if the objective consists in maximizing sales across several periods. We also mention several more general marketing decision problems which could be solved by appropriate modifications of the algorithm presented.

1 Introduction

Allocation decisions in marketing refer to decision variables like advertising budgets, sales budgets, sales force sizes, and sales calls which are allocated to sales units like sales districts, customer groups, individual customers, and prospects. Studies using optimization methods and empirical sales response functions provide evidence to the

importance of such allocation decisions. These studies demonstrate that sales or profits can be increased by changing allocation of budgets, sales force or sales calls (see, e.g., Beswick and Cravens 1977, LaForge and Cravens 1985, Sinha and Zoltners 2001). The average increase of profit contribution across studies analyzed in the review of Sinha and Zoltners (2001) compared to the current policies was 4.5 % of which 71 % are due to different allocations and 29 % are due to size changes. The smaller second percentage can be explained by the well known flat maximum principle (Mantrala et al. 1992).

All these studies require knowledge of the mathematical form of sales response functions which reproduce the dependence of sales on decision variables. In addition they require that parameters of sales response functions are estimated by econometric methods using historical data or by means of a decision calculus approach which draws upon managers' experiences (Gupta and Steenburgh 2008). Of course, there are situations in which both econometric methods and decision calculus cannot be applied. Lack of historical data (e.g., for new sales units), lack of variation of past allocations, lack of experiences with the investigated or similar markets constitute possible causes.

In such difficult situations the question arises how management may arrive at rational allocation decisions nonetheless. We are aware of only one appropriate approach developed by Albers (1998). He demonstrates that in spite of the lack of knowledge on functional form and parameters the allocation problem may be solved by an iteration along elasticities. Albers investigates several sales response functions with different properties. Note that these functions and their parameters are not used in the iterations, they only serve to generate values of the dependent variable sales by deterministic simulation. Of course, by applying deterministic simulation random disturbances are excluded though the latter usually are an additive component of sales response models estimated by econometric methods.

We develop an exploration-exploitation algorithm which extends the approach of Albers in order to cope with additive random disturbances. During a certain number of iterations, the algorithm collects data about the function and its distortion caused by random disturbances (exploration). In the second phase, the data gathered are used to solve the problem more efficiently (exploitation). In the exploitation phase we approximate the unknown functions by quadratic polynomials and obtain solutions by quadratic programming.

Chapter 2 describes the problem from a mathematical point of view. In chapter 3 we present the necessary preparations for the simulation study, the algorithm is describen

in chapter 4. Results of the simulation study are presented and discussed in chapter 5. In chapter 6 we investigate the performance of the algorithm under conditions different from those in the simulation study. We also mention several extensions of the allocation problem treated here which maybe could be solved by modifications of our algorithm. Appendix A contains a mathematical discussion of the general theory. The algorithm is presented as pseudocode in Appendix B. Further results not given in the main text can be found in Appendix C.

2 Decision Problem

A (scarce) resource B needs to be allocated to several ($n \in \mathbb{N}_{>1}$) units. We have one sales response function f_i for each unit $i = 1, \dots, n$ which depends on its allocated input x_i only. Allocations must be non-negative and lower than the resource, i.e., $0 \leq x_i \leq B$. In addition the sum of all inputs must not exceed the resource:

$$\sum_{i=1}^n x_i \leq B \quad (1)$$

The n -tuple (x_1, \dots, x_n) satisfying (1) is called *allocation*. Total sales, i.e, the sum of sales across all units $\sum_{i=1}^n f_i(x_i)$, represent the *objective* of this allocation. The goal is to find an allocation maximizing total sales:

$$\max \sum_{i=1}^n f_i(x_i) \quad (2)$$

As we assume that all sales response functions are monotonically increasing, we can conclude that condition (1) is binding and can therefore be rewritten as:

$$\sum_{i=1}^n x_i = B \quad (3)$$

The decision problem presented so far makes it possible that no (in other words: zero) resources are allocated to a unit (e.g., to make no sales calls in certain districts). Of course, decision makers may find it inappropriate to deprive a unit of all resources. To cope with such a situation one defines a new problem in which a modified total resource $B' := B - \sum_{i=1}^8 lb_i$ is allocated to the units with functions $g_i(x_i) := f_i(lb_i + x_i)$. Now the inputs of some (or all) units have a lower bound $lb_i > 0$ and B' must be zero or positive.

3 Preparing the Simulation Study

3.1 Sales Response Functions

We consider the same four functional forms investigated in Albers (1998), i.e., the multiplicative, the modified exponential, the concave and the S-shaped ADBUDG function:

$$f_{mul}(x) = ax^b \quad (4)$$

$$f_{exp}(x) = M_{exp}(1 - \exp(-xh)) \quad (5)$$

$$f_{adc}(x) = M_{adc} \frac{x^{f_c}}{g_c + x^{f_c}} \quad (6)$$

$$f_{adS}(x) = M_{adS} \frac{x^{f_s}}{g_s + x^{f_s}} \quad (7)$$

$$(8)$$

where $a, M_{exp}, M_{adc}, M_{adS}, h, g_c, g_s$ all are positive constants.

We now discuss properties of these four functional forms (for more details see, e.g., Hanssens et al. 2001). $M_{exp}, M_{adc}, M_{adS}$ symbolize maximum values of sales, in other words sales potentials. Given certain parameter restrictions the first three functions allow for a concave shape, i.e., they reproduce positive marginal effects which are increasing with higher values of x . These restrictions are $0 < b < 1, h > 0$ and $0 < f_c < 1$, respectively. The second version of the ADBUDG function f_{adS} leads to an S-shape for $f_s > 1$. S-shaped functions consist of two sections. The first section is characterized by increasing positive marginal effects, the second one by decreasing positive marginal effects.

One by one, a function-type is selected, eight different sets of parameters for these functions are chosen (each for one of eight sales units), and a fixed resource $B = 8,000,000$ is allocated among these units by the algorithm presented in section 4. In section 6, these conditions are varied. In Albers' original paper these parameters are generated from a table. To the best of our knowledge, this table contains inconsistencies for the last three function-types. For example, for the modified exponential function which has two parameters, M_{exp} and h Albers applies three restrictions. This procedure leads to an overdetermined equation system. The ADBUDG functions have an additional parameter, but also an additional restriction ($f_c < 1$ and $f_s > 1$). That is why again overdetermined equation systems result.

We therefore construct a new table and explain how parameters are obtained from this table in the next section.

	similar	varied	similar	varied
I	elasticities	elasticities	saturation levels	saturation levels
1	0.26	0.11	6,100,000	4,500,000
2	0.27	0.12	6,200,000	10,000,000
3	0.28	0.13	6,300,000	4,500,000
4	0.29	0.14	6,400,000	10,000,000
5	0.31	0.47	6,600,000	4,500,000
6	0.32	0.48	6,700,000	10,000,000
7	0.33	0.49	6,800,000	4,500,000
8	0.34	0.50	6,900,000	10,000,000

Table 1: Function Properties for Parameter Generation

As the multiplicative function does not have a saturation level, we define its 'saturation' as its value for the maximum input, i.e., $f_{mul}(B)$.

To cope with the inconsistencies in Albers' table we do not include contribution margins and use different saturation levels.

3.2 Generating Parameters

Parameters of the functions are generated starting from table 1. For the multiplicative function, matters are simple, as b is the elasticity, and $a := M \cdot B^{-b}$, where M is the saturation level.

For the other functions, we perform an iterative search for parameter values. As the saturation level is given, only the remaining parameters need to be determined.

One can derive the elasticity of the modified exponential function as

$$\varepsilon = \frac{\exp(-hx)hx}{1 - \exp(-hx)} \quad (9)$$

which can be transformed into a fix-point-formula

$$h = \frac{-\log\left(\frac{\varepsilon(1-\exp(-hx))}{hx}\right)}{x}. \quad (10)$$

Using a starting value $h_0 > 0$ and $x = R/d$, fix-point iterations quickly converge to a positive value for h which satisfies the conditions in the table.

For the ADBUDG-functions which have one parameter more we choose special values which yield reasonable shapes and ensure stability of the iteration process.

3.3 Disturbances

To each function we add normally distributed disturbances $u \sim \mathcal{N}(0, \sigma^2)$. Variances σ^2 are set to attain a desired share of explained variance for the dependent variable sales. These desired R^2 values amount to 0.9, 0.7, and 0.5. For each function the error variance σ^2 is set to the value which leads the R^2 -value closest to its desired value in a regression of that function using 2,000 uniformly distributed integer values of the inputs.

3.4 Rules of Thumb

We start from the same rules of thumb as Albers (1998):

1. Allocation proportional to sales of a unit in the previous period.
2. Allocation proportional to sales of a unit divided by its allocation in the previous period.
3. Allocation proportional to the saturation level of a unit.

The third rule cannot be implemented as saturation levels like all parameters of functions are unknown. That is why we replace it by the following rule.

- 3.' Allocation proportional to maximum sales of a unit observed so far.

For the remainder of the paper we will refer to these as first, second and third rule of thumb.

3.5 Estimators for Elasticities

On the basis of several definitions of point and arc elasticity to be found in the literature (see, e.g., Vázquez (1995) P.223 and Seldon (1986) P.122) we obtain four different estimators:

$$\frac{\log(\frac{y_2}{y_1})}{\log(\frac{x_2}{x_1})}, \frac{\frac{y_2-y_1}{y_1}}{\frac{x_2-x_1}{x_1}}, \frac{\frac{y_2-y_1}{y_2}}{\frac{x_2-x_1}{x_2}}, \frac{\frac{y_2-y_1}{\bar{y}}}{\frac{x_2-x_1}{\bar{x}}} \quad (11)$$

where $\bar{x} := \frac{x_1+x_2}{2}$ and $\bar{y} := \frac{y_1+y_2}{2}$.

x_1 and x_2 denote inputs of two consecutive periods, y_1 and y_2 their corresponding outputs (sales).

The results we present later are obtained using the third estimator which leads to highest values for the objective “Sales” (see section 5). But note that the difference to the other three estimators is not statistically significant according to a regression analysis with main effects only.

3.6 Elimination of correlation conditions

Albers also compares different starting conditions with respect to the correlation of starting allocations with their optimal values. These starting conditions are constructed by changing an equal distribution by roughly 5% towards a positive or negative correlation to the optimum. However, this only makes sense in a situation without random disturbances. In our case, the magnitude of the error terms, even in the case of $R^2 = 0.9$, greatly exceed a 5% boundary and hence these starting condition variations need not be considered. Therefore the starting allocations are equal with $B/8 = 1,000,000$ for each unit.

4 Developed Algorithm

Albers (1998) intends to show that in allocation problems iterations along elasticities always outperform rules of thumb, independent from functional form, other properties of the functions and correlations of starting values with optimal allocations. He considers for each of the four functions discussed in the previous section every constellation of different values of properties and correlations. Sales are computed on the basis of these functions without adding disturbances though the latter are a component of most econometric models. Results obtained by his iterative algorithm are compared to those computed by the three rules of thumb explained in the previous section. Overall, his iterative algorithm outperforms all rules of thumb by far.

In our study the algorithm of Albers turns out not to work well for sales response functions with additive disturbances. Disturbances which directly affect elasticities cause elasticities and new allocation values to fluctuate. Very small allocations are computed quite often which in their turn lead to a low function value before a disturbance is added. Under such circumstances adding a disturbance frequently results in a negative value and the logarithmic estimator of elasticities which is used in the algorithm of Albers does not work. But note that any of the other three estimators shown in section 3.5 also leads to heavily biased, sometimes even negative, elasticities.

4.1 Exploration

In order to bring stability to the process, we apply first order exponential smoothing in the following way:

$$\tilde{\epsilon}_t := (1 - \beta)\tilde{\epsilon}_{t-1} + \beta\hat{\epsilon}_t \quad (12)$$

with $\beta \in [0, 1]$ (we use $\beta := 0.85$ based on a comparison of different values).

$\tilde{\epsilon}_t$ is the smoothed elasticity and $\hat{\epsilon}_t$ the estimated elasticity in period t . Moreover, each elasticity value is projected into the interval $[0.01, 0.5]$, i.e., if the calculated value is above 0.5 it was set to 0.5, and it was set to 0.01 if it was below 0.01.

So far we have described the exploration phase of the algorithm. As allocations generated in this fashion are close to the optimum we consider it to be useful for exploration. But even the exponential smoothing version still jumps around too much. Therefore there is dire need for a method which dampens disturbances.

4.2 Exploitation

The general idea can be easily understood when looking at modified exponential functions with varying parameters. The optimal allocation for the function in the example drawn in figure 1 is roughly 640,000. Nevertheless, even when the error term is small, the algorithm and each of the three rules of thumb still fluctuate a lot, showing no sign of stability, although they do not leave a certain interval of the domain in each variable (and hence of the codomain).

This area shown in the figure looks like it can be easily approximated by a parabola, i.e. a polynomial of degree two. Assuming the functions were actually polynomials of degree two, quadratic programming could then give an exact solution, as the boundary condition is linear.

The two steps necessary for optimization are hence: exploration and exploitation, a concept first introduced by March (1991). During a fixed number of periods, the algorithm generates data points close enough to the optimum. After that, a quadratic regression of the form

$$y \sim ax^2 + bx + c \quad (13)$$

is performed for each unit in order to approximate its unknown function based on all values of sales and allocations available so far (using a smaller number of the most




Figure 1: True Function and Fitted Parabola

recent values only does not improve results). Then quadratic programming yields an allocation which is optimal *for these approximations*, obeys the total resource restriction and provides a new data point for the regression. We use the method of Goldfarb and Idnani (1983) for quadratic programming in our implementation. This process is repeated until a total of 40 iterations is reached.

A problem arises when a is estimated as a positive number for any sales unit, as the matrix in the quadratic program is no longer positive definite. This, however, has a surprisingly easy fix: a can be set to a very small value (we choose -10^{-15}) and b is set to the slope of a linear regression line, thereby “fooling” the quadratic program into accepting something that is basically a straight line rather than a parabola.

In the worst-case-scenario, additionally, the slope of the regression line may be negative. This case is very rare and the allocation to this unit will almost certainly be zero. One should remember however, what this actually means: The shape of the data points resembles a monotonically decreasing, convex (!) function and would hence arise either from a few very unlucky error terms in a row or an outer influence that cannot be ex-

plained by an additive error term. Surely in this case the function should be thoroughly analyzed instead of continuing the application of *any* algorithm. The situation will be briefly mentioned in section 6.

A final question that arises is when to switch from exploration to exploitation. As the exact functions and variances are unknown to the algorithm, there can be no mathematical derivation of the optimal switching point. The main idea is that for higher error variances, more exploration is necessary to make sure the parabolas are sufficiently accurate. This was confirmed in a separate simulation wherein the R^2 -Levels of 0.9, 0.7 and 0.5 optimally had 10, 14 and 18 iterations respectively. Hence, an estimator for the switching period was defined based on these results, dependent on the estimated variance of the data. After the first nine iterations, this number is estimated, and lies in the interval $[10, 25]$.

4.3 Related Allocation Problem

The form of the investigated allocation problem presented in section 2 also gives rise to the solution of a different, but related problem for free. In the related decision problem one part of a given budget B may be spent (allocated), while the other part may be saved. Instead of (2) we obtain the following objective function

$$\sum_{i=1}^n f_i(x_i) + (B - \sum_{i=1}^n x_i). \quad (14)$$

in which the remaining, unspent budget $B - \sum_{i=1}^n x_i$ is added to the sum of sales across all units.

This problem looks different from the one we have investigated so far, but rewritten it turns out to be just a special case. Define the function $f_{n+1}(x) := x$, and consider the allocation of B onto the $n + 1$ functions f_1, \dots, f_{n+1} . A complete allocation now means finding (x'_1, \dots, x'_{n+1}) that sum up to B . But that is equivalent to finding (x_1, \dots, x_n) that sum up to a value $R \leq B$, since we can then put $x'_1 := x_1, \dots, x'_n := x_n, x'_{n+1} := B - R$, and they evidently add up to B .

So now we can see that this problem can be solved by the algorithm discussed above, but we know even more: as $f_{n+1}(x) = x$, its elasticity is exactly 1 and hence need not be estimated, no problems arise in the exploration phase. As f_{n+1} is a straight line, we use the fix discussed in 4.2 by adding a tiny parabola to make it compatible with the quadratic program. As this function has no additive disturbance, the problem is actually easier to solve than a regular allocation to $n + 1$ units. The optimum is found when the

slopes of all functions are equal to one (as the Lagrange-multiplier becomes 1, see A.3) and the remaining budget is saved. This is to be expected, as the slope of $f(x) = x$ is constant 1 and every further unit of the budget will be added to the function with the steepest slope.

5 Evaluation of Procedures

We want to compare four different procedures, i.e., three rules of thumb and the algorithm introduced in section 4. To this end we conduct a simulation study and consider two different dependent variables which both are computed as arithmetic means across 40 periods. The first dependent variable “Sales” is based on total sales attained by rules of thumb or the algorithm in each period. The second dependent variable “Optimality” is a relative measure, the ratio of total sales and optimal total sales. Optimal total sales are determined by optimizing on the basis of the true response functions without disturbances, i.e., assuming knowledge of sales response functions and their parameters. Optimality therefore shows to what extent a procedure which lacks knowledge of the response functions attains optimal total sales on average. A value of 1.0 for optimality indicates as a rule that average total sales equal their optimal value.

As mentioned above, the S-shaped ADBUDG-function is not concave, and hence a problem arises with local and global optima. In particular, the nine control algorithms that search for the optimal solution may get stuck in local optima. Therefore optimality values greater than 1.0 maybe obtained.

We look at 192 constellations, which result from four function types, four procedures, similar/varied elasticities, similar/varied saturations, three error levels and generate five replications for each constellation. A seed is set to ensure reproducibility.

The dependent variables are normalized in the following manner: As “Optimality” usually is already a number between 0 and 1, no normalization is necessary. Since the four function types yield quite different values for total sales (which is especially pronounced for the ADBUDG-functions), “Sales” are divided by the maximum value *of the respective function type*. This allows to compare the effectiveness of procedures across different functional forms.

Finally, two linear regressions are performed in the following way:

$$Sales \sim \beta_{1,0} + \beta_{1,1}Proc + \beta_{1,2}Form + \beta_{1,3}Dist + \beta_{1,4}Elas + \beta_{1,5}Satu \quad (15)$$

$$Optimality \sim \beta_{2,0} + \beta_{2,1}Proc + \beta_{2,2}Form + \beta_{2,3}Dist + \beta_{2,4}Elas + \beta_{2,5}Satu \quad (16)$$

All independent variables are categorical, where “Proc” takes values corresponding to the three rules of thumb and the algorithm described above, “Form” takes the four values corresponding to the four types of functions, “Dist” takes three values corresponding to the R^2 values of 0.9, 0.7 and 0.5, and “Elas” and “Satu” take two values corresponding to “similar” and “varied”, as explained above.

Moreover, we also estimate regression models which in addition include certain sets of interaction terms. As models with main effects only are inferior to models with interactions, a discussion of the main effect regressions can be found in Appendix C.

5.1 Results for Regression Models with Interactions

We consider several model types with interaction terms. The most simple of these models includes only pairwise interactions of the “Procedure”-variable with all other variables. In this model we can easily analyze the interactions between procedures and the other variables, while also, with the use of F-tests, being able to compare the performance of procedures (results of these models and models with all pairwise interactions can found in Appendix C).

To consider additional interaction terms we also investigate models (one for each dependent variable) with all pairwise interactions and models with double, triple, quadruple and quintuple interactions. Quite interestingly, the full models with quintuple interactions not only lead to the best variance explanations (adjusted R^2 values amount to 0.8786 and 0.85 for “Sales” and “Optimality”, respectively), but F-tests also confirm that the full models outperform all the restricted models.

A complete explanation of the F-tests used to compare procedures is given in Appendix C section C.3, here we present the main results.

These comparisons consist of two steps (C.2 is a more comprehensive example, since it has fewer coefficients and a simpler null hypothesis). In the first step we calculate the effect of each procedure by summing all its interaction terms. Then for each pair of procedures we subtract their two respective effects (for a detailed explanation see the example in section C.2). The corresponding matrix (whose product with the coefficient vector is zero under the null hypothesis) was the basis of an F-test. Hence the sign of

the difference indicates if the algorithm performs better than a rule of thumb, and the F-statistic reveals the significance of a difference .

For the dependent variable “Sales” we obtain average differences between the algorithm and each of the three rules of thumb of 4.8325, 7.2464, and 3.7383, respectively. These differences are highly significant at levels lower than 0.001 (the corresponding F-statistics are 46.426, 104.389 and 27.782). Based on these results we conclude that the algorithm clearly beats the three rules of thumb in terms of Sales.

For the other dependent variable “Optimality” we obtain differences of -0.9251, 1.7389 and -2.2644 (F-statistics are 1.4663, 5.1813 and 8.7861), respectively. At a significance level of 0.05 the algorithm performs better than the second rule of thumb, but worse than the third rule of thumb. In addition no significant performance difference with respect to the first rule of thumb is found.

5.2 Discussion

We begin by discussing the results for “Sales”. As seen in the full model, the algorithm is vastly superior to all three rules of thumb. This is the most important result, which leads to clear implications for marketing decisions: Whenever a scarce resource needs to be allocated to different units, it is highly beneficial to use all the data like in the algorithm, instead of considering only the most recent observation like in the rules of thumb investigated here. In applying our algorithm two situations can be distinguished. If historical data with enough variation are lacking, begin with exploration and switch to exploitation later on. If on the other hand appropriate historical data are at hand, one may start with the exploitation stage right away, i.e., use quadratic regression and quadratic optimization. *This approach should highly improve “Sales” as compared to its value obtained by rule of thumbs.*

With respect to “Optimality” we obtain much lower F-statistics for procedure comparisons, probably due to the fact that this dependent variable is limited to the unit interval with respect to a global maximum. As long as procedures do not behave chaotically, not a lot of variance and difference between algorithms can be expected. This will be discussed shortly, but first let us consider the results.

As no significant difference can be seen between the algorithm and the first rule of thumb, no further discussion is necessary here. The second rule is outperformed because of certain chaotic patterns that can be visualized by manually starting optimiza-

tions with low levels of R^2 . We can see that our algorithm clearly outperforms the second rule of thumb both with respect to “Sales” and “Optimality”. This disadvantage of the second rule of thumb is also confirmed by the second and hence by both F-tests explained above. The third rule of thumb leaves us with mixed signals. While we seem to obtain significantly better sales with our algorithm, the third rule seems to perform significantly closer to the optimum. This is not as contradictory as it first seems, since the rule literally says: “take the highest value you can get” but might then refuse to move from the position it has settled in.

We see three reasons for not using the third rule of thumb:

1. The rule should not be used if decision makers have more interest in high achieved sales and do not care how close sales are to an optimal value which is based on unknown functions and ignores disturbances. This recommendation follows from the results of our simulation study for the objective “Sales” which clearly show that the algorithm performs better.
2. The rule cannot be modified to deal with extended marketing decision problems (e.g., if an allocation affects other units as well, if sales depend on marketing variables of different types, etc.) which may be investigated by future research. We shortly discuss several extensions in the next section and indicate that the algorithm can be modified to handle these more general problems.
3. This rule uses almost no information from previous periods to determine the allocation.

In Albers’ original version it was just defined as the allocation proportional to the saturation levels, which as a rule are not known in practice. Providing examples of functions for which this rule of thumb behaves very badly is rather simple. For example, given a resource of 700,000 and eight S-shaped ADBUDG-functions with different elasticities and varied saturation levels and medium disturbances the third rule leads to solutions which are clearly worse compared to those determined by our algorithm (see table 2). This weakness of the third rule can be consistently observed if resources are low.

6 Conclusions

In addition to the simulation study presented we examine whether performance of the developed algorithm remains stable if it has to deal with different conditions. First of all we analyze how procedures behave given a different number of units. The performance

Period	Algorithm	ROT1	ROT2	ROT3
1	0.6327	0.5438	0.2995	0.6981
2	0.7307	0.5108	0.6494	0.8169
3	0.7801	0.6119	0.2068	0.7991
...
17	0.6499	0.6421	0.3466	0.6683
18	0.9479	0.6447	0.3087	0.6683
19	0.9055	0.6612	0.6145	0.6683
20	0.9625	0.6500	0.4258	0.6683
21	0.9727	0.6603	0.1992	0.6683
22	0.9789	0.6725	0.4874	0.6683

Table 2: Period optimalities allocating a resource of 700,000 for S-shaped ADBUDG functions (the exploitation stage begins at iteration 18)

of the algorithm remains similar to that obtained in the simulation no matter whether we consider less (four) or more (twenty) units.

Remember that in the simulation study in each experimental constellation sales are generated by functions which have the same form for all units. Of course, in real life situations sales response of units may be very nonuniform due to, e.g., economic or cultural factors which may differ sharply between regions or customer groups. Now sales response can no longer be reproduced by different coefficient values and requires the use of different functional forms. Combinatorics tells us right away that an exhaustive proof of the superiority of our algorithm in this case is not possible (as we would need to consider $4^8 = 65536$ different function constellations), so we look at several random as well as explicitly designed conditions. We randomly choose the function type for each unit and repeated this exercise by varying the number of units. Still for these constellations our algorithm usually determines better solutions than the rules of thumb.

Summing up we recommend to use the developed algorithm to solve marketing allocation problems of the form shown in section 2 or the related problem introduced in section 4 if sales response functions are unknown. We justify this recommendation by the good performance of the algorithm demonstrated by the simulation study and its stability under changing conditions. Future work might consider modifications of the algorithm for the decision problem investigated here. What effect would it have to

choose a different algorithm in the exploration phase? A next goal would be to check if other algorithms are more suitable for the exploitation phase. This might mean small adjustments of parameters, or a completely new algorithm.

As mentioned in section 4, a problem arises if the parabola has a positive leading coefficient and the regression line has a negative slope. This effect usually occurs if a unit repeatedly receives allocations near zero. Due to the additive disturbances the data set for the regressions then consists of very similar x -values, while the y -values vary a lot. As low allocations are usually not optimal, an amendment of the algorithm may be beneficial for such situations.

Modifying the algorithm to solve more general marketing decision problems also seems to be an interesting task of future research. One extended decision problem results if one or several sales functions may change suddenly. We suspect that under such circumstances the exploration phase will have to start once again. For situations with gradual change on the other hand an easy fix would be to delete older data points before each iteration. One could also investigate multi-variable generalizations. In one generalization allocations affect sales of the same unit as well as sales of other units. Another more challenging generalization allows for marketing variables of different types. Examples of such variables are advertising and price or advertising and sales effort, where both variables have an effect on sales of different units.

A Appendix: Mathematical Discussion

This section is a repetition of the results from Albers (1998).

As mentioned above, a scarce resource B is to be allocated to n different units, where we want to find the maximum objective

$$\max \sum_{i=1}^n f_i(x_i) \tag{A.1}$$

of all allocations (x_1, \dots, x_n) .

The most suitable mathematical context for this problem is the Lagrangian formalism for optimizing a function (2) with a given condition (3).

A.1 Albers' Lagrange-Ansatz

The Lagrangian one needs to consider takes the form

$$L = \sum_{i=1}^n f_i(x_i) + \lambda \cdot \left(\sum_{i=1}^n x_i - B \right), \quad (\text{A.2})$$

and one shall differentiate (A.2) by the x_i and λ and set the derivations equal to zero to get the conditions an extremum needs to satisfy.

One receives

$$\frac{\partial L}{\partial x_i} = \frac{\partial f_i}{\partial x_i} - \lambda \stackrel{!}{=} 0 \quad (\text{A.3})$$

and of course

$$\sum_{i \in I} x_i = B. \quad (\text{A.4})$$

As the point elasticity is defined as $\frac{\partial f_i}{\partial x_i} \cdot \frac{x_i}{f_i} = \varepsilon_i$ we can rewrite $\frac{\partial f_i}{\partial x_i}$ as $\varepsilon_i \cdot \frac{f_i}{x_i}$ and insert into (A.3)

$$\varepsilon_i \cdot \frac{f_i}{x_i} = \lambda \Rightarrow x_i = \frac{f_i \varepsilon_i}{\lambda}. \quad (\text{A.5})$$

In particular, the resource becomes

$$B = \sum_{i \in I} x_i = \sum_{i \in I} \frac{f_i \varepsilon_i}{\lambda}, \quad (\text{A.6})$$

which can be solved for λ and inserted into (A.5) to obtain

$$x_i = \frac{f_i \varepsilon_i}{\sum_{j \in I} f_j \varepsilon_j} B. \quad (\text{A.7})$$

In general, the exact form of the functions f_i , in particular its point elasticity is unknown.

However, given two values x_i and x'_i , and their respective outputs $y_i = f_i(x_i)$, $y'_i = f_i(x'_i)$

, the arc elasticity may be estimated as

$$\varepsilon_i = \frac{\ln\left(\frac{y_i}{y'_i}\right)}{\ln\left(\frac{x_i}{x'_i}\right)}, \quad (\text{A.8})$$

where \ln is the natural logarithm. Of course, other estimators may be used here as well, as explained above. This gives rise to an iterative algorithm, where in each step the elasticities are estimated dependent on the last two iterations and the components x_i of the new allocation (x_1, \dots, x_n) are obtained from that elasticity.

A.2 Further results

As the functions are concave and the process closes in on the Lagrange solution, the method presented will find a solution to the allocation problem, *if it exists*. To guarantee the existence of a (unique) solution, a descent into mathematics is necessary. If the reader is unfamiliar with concepts of continuity, compactness and the extreme value theorem for topological spaces (or subsets of \mathbb{R}^n) he may consult Rudin (1976).

Neither negative inputs nor outputs are to be expected, so consider $\mathbb{R}_0^+ := \{x \in \mathbb{R} | x \geq 0\}$.

Definition 1

A function $f : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$ satisfies the *conditions of diminishing returns*, if

1. $f \in C^2(\mathbb{R}_0^+)$ (double differentiability)
2. $f'(x) \geq 0 \quad \forall x \in \mathbb{R}_0^+$ (monotony)
3. $f''(x) < 0 \quad \forall x \in \mathbb{R}_0^+$ (strict concavity)

(This condition may actually be weakened. We only require single differentiability; condition 3 then becomes “ f' is strictly monotonically decreasing”)

A.3 Existence

Proposition 1

Let $(f_i)_{i \in I}$ be a finite family ($\#I = n$) of continuous functions. Then for every $B > 0$ there is a solution to the allocation problem (2) & (3).

Proof:

The map

$$F : (\mathbb{R}_0^+)^n \rightarrow \mathbb{R}_0^+, (x_i)_{i \in I} \mapsto \sum_{i=1}^n f_i(x_i) \quad (\text{A.9})$$

is continuous, since the f_i are, which follows from Rudin P.87 Theorem 9 and Theorem 10. The bounded hyperplane

$$H := \{(x_i)_{i \in I} | \sum_{i=1}^n x_i = B\} \quad (\text{A.10})$$

is a closed (in the topological sense; this is equivalent to “the limit of every convergent sequence with terms in H also lies in H ”; the reader may verify that H is in fact closed, by using that the function $(x_1, \dots, x_n) \mapsto \sum_{i=1}^n x_i$ is sequentially continuous) subset of $(\mathbb{R}_0^+)^n$ and hence compact (by the Heine-Borel-theorem, see Rudin P.40 Theorem 2.41). The restriction of F to H is continuous as well, since continuity is a local property.

Since every continuous function from a compact set to \mathbb{R} attains its maximum (see Rudin P.89/90 Theorem 4.16), so does $F|_H$ which ends the proof. \square

A.4 Uniqueness

Proposition 2

Let $(f_i)_{i \in I}$ be a finite family of functions satisfying the conditions of diminishing returns. Then for every $B > 0$ the solution to the allocation problem (2) & (3) is unique.

Proof:

Each of the functions f_i is concave in the multidimensional sense as a map

$$\mathbb{R}^n \rightarrow \mathbb{R}, \quad (x_i)_{i=1,\dots,n} \mapsto f_i(x_i) \quad (\text{A.11})$$

and hence, so is their sum

$$\mathbb{R}^n \rightarrow \mathbb{R}, \quad (x_i)_{i=1,\dots,n} \mapsto \sum_{i=1}^n f_i(x_i) \quad (\text{A.12})$$

and the restriction to the convex subset of allocations in $(\mathbb{R}_0^+)^n$. The maximum must therefore be unique. \square

B Appendix: The Algorithm in Pseudocode

Notation: for two vectors a and b of the same dimension, we denote by $\langle a, b \rangle$ their dot product, and by $a * b$ the componentwise multiplication, i.e. the vector $(a_i b_i)_{i=1}^n$

Data: $x^1 = (x_1^1, \dots, x_n^1)$, x^2 start values, $0 \leq \beta \leq 1$ smoothing parameter, maxit: maximum number of iterations, B: resource, I_n : identity matrix of dimension n , exex=10: first estimate of number of iterations in exploration phase, \tilde{f} : multidimensional map consisting of the separate functions in each component, DF : array which will be filled with the data points of all functions

$y^1 := \tilde{f}(x^1); y^2 := \tilde{f}(x^2);$

while $i < exex$ **do**

Test(x^1, x^2, y^1, y^2);

$\varepsilon := \frac{y^2 - y^1}{\frac{x^2 - x^1}{x^1}}$; Epstest(ε);

if $i \geq 2$ **then**

$\varepsilon_{sm} := \varepsilon \cdot (1 - \beta) + \varepsilon_{old} \cdot \beta$

else

$\varepsilon_{sm} := \varepsilon;$

end

$up := \frac{y^1 \cdot \varepsilon_{sm}}{\langle y^1, \varepsilon_{sm} \rangle} \cdot B;$

$\varepsilon_{old} := \varepsilon_{sm};$

$x^2 := x^1; y_2 := y^1;$

Test2(up, x^2);

$x^1 := up; y^1 := \tilde{f}(x^1);$

if $i = 9$ **then**

 estimate variance and exex

end

add (x^1, y^1) to DF ;

end

for $i = exex$ to maxit **do**

for $j = 1$ to n **do**

 Perform Regression $y_j \sim a(x_j)^2 + bx_j + c$

end

D : Diagonal Matrix containing a -values $\cdot (-2)$

A : n by $n + 1$ -Matrix with -1 in the first column followed by I_n

d : Vector containing b -values

bv : Vector of length $n + 1$ containing $(-B, 0, 0, \dots, 0)$

$up := \text{solve.QP}(D, d, A, bv)$

$x^1 := up; y^1 := \tilde{f}(x^1);$

add (x^1, y^1) to DF ;

end

Remark: The functions Test and Test2 check if the entries are too small or too close to each other, epstest projects elasticities into the Interval $[0.01, 0.5]$.

Note that the estimator for “exex” was determined empirically and is directly dependent on *maxit*

The function solve.QP solves the quadratic program as described in Goldfarb/Idnani (1983) with the notation from the R-Package “quadprog”.

Variable	Dependent Variable: Sales		Dependent Variable: Optimality	
	Coefficient	t-Value	Coefficient	t-Value
Intercept	0.881	118.582***	1.023	128.317***
Proc2	-0.014	-2.271*	-0.017	-2.510*
Proc3	-0.065	-10.209***	-0.073	-10.676***
Proc4	0.008	1.328	0.011	1.595
Form2	0.034	5.306***	0.024	3.462***
Form3	0.102	16.171***	0.034	5.030***
Form4	-0.009	-1.401	-0.020	-2.896**
Dist2	-0.018	-3.324***	-0.021	-3.636***
Dist3	-0.047	-8.573***	-0.057	-9.722***
Elas2	-0.021	-4.762***	-0.051	-10.544***
Satu2	0.058	12.882***	-0.040	-8.332***
* p-value<0.05, ** p-value<0.01, *** p-value<0.001				

Table 3: Main Effect Regression Models

C Appendix: Detailed Regression Results

C.1 Main effect models

The regression results of the main effect models both for “Sales” and “Optimality” are shown in table 3. The independent variables are coded by binary dummy variables with the developed algorithm, the multiplicative function, similar elasticities and similar saturations as reference categories.

C.2 Models with procedure interactions

In addition to main effects we now consider pairwise interaction between procedures and each of the other independent variables. We compute the total effect of our algorithm in the following manner. We add the intercept to each main effect coefficient and sum these intermediate values. To this sum we add the intercept once more to also consider the constellation in which all independent variables are in their reference category. The first rule of thumb would be represented by the coefficients of the interactions of the rule of thumb with each condition plus the coefficient of the rule of thumb plus intercept, plus another sum of the rule of thumb plus intercept. For the corresponding

F-test we construct a matrix that, when multiplied with the coefficient vector, gives us the difference of these two real numbers, where the null hypothesis is that number being zero. The matrix is hence a 1×32 matrix with a -8 in second entry, a 1 in entries (5,6,7,8,9,10,11), a -1 in entries (12,15,18,21,24,27,30), and zeroes elsewhere.

The resulting coefficient for “Sales” is 0.2073, implying that the algorithm is better than the first rule of thumb. The F-statistic takes the value 9.4066 and follows an $F(1,928)$ -distribution given the null hypothesis. Therefore the algorithm is better than the first rule of thumb at a significance level of 0.0022. Furthermore it outperforms the third rule of thumb with respect to “Sales” with a p-value of 0.056 and the second rule of thumb with respect to optimalities with a p-value of 0.025.

C.3 F-Tests in the Full Models

In order to compare the strengths of the algorithms in the full model, F-Tests were performed in the following manner. The coefficient vector of the OLS-Estimator was multiplied with a matrix representing the difference of the algorithms. This matrix was constructed analogously to the ones in the previous section: For every constellation of conditions a matrix was defined as the sum of the coefficients containing the algorithm together with the corresponding interaction terms minus the coefficients containing the rule of thumb that was to be discussed, again, including their respective interactions. All these matrices were added yielding three matrices, one for each rule of thumb. This data is represented in table 4, where the variables are shortened to their first letter.

The coefficients in the table hence represent the number of effects it appears in (as the null hypothesis is the difference of sums of the effects). Hence if k_i is the number of categories of the variable i , the number of the effects the l -fold interaction term $i_1 \times \dots \times i_l$ appears in (for $l \in \{0, \dots, 5\}$), is

$$\prod_{i \notin \{i_1, \dots, i_l\}} k_i. \quad (\text{C.1})$$

For example, the interaction term $P1 : F2 : D2 : E1$ appears in its own effect (together with lower terms) where $S = 2$ (as $S2$ is the reference category) and its counterpart where $S = 1$, represented by itself, lower terms and the coefficient of $P1 : F2 : D2 : E2 : S2$. Via the formula we get

$$\prod_{i \notin \{F,D,E\}} k_i = \prod_{i \in \{S\}} k_i = k_S = 2. \quad (\text{C.2})$$

Name	β_1	β_2	M1	M2	M3	Name	β_1	β_2	M1	M2	M3
(Intercept)	0.82277	0.99693	0	0	0	P3:F4:S1	0.01025	0.00945	0	0	-6
P1	-0.00588	-0.00007	-48	0	0	P1:D2:S1	-0.02060	-0.00951	-8	0	0
P2	0.00250	-0.00070	0	-48	0	P2:D2:S1	-0.13402	-0.13451	0	-8	0
P3	-0.00285	0.00094	0	0	-48	P3:D2:S1	-0.00755	-0.00361	0	0	-8
F2	0.05685	-0.00393	12	12	12	P1:D3:S1	0.02295	-0.00087	-8	0	0
F3	0.13735	-0.00244	12	12	12	P2:D3:S1	-0.22894	-0.26788	0	-8	0
F4	0.04394	-0.00504	12	12	12	P3:D3:S1	0.00415	0.00721	0	0	-8
D2	-0.01137	-0.00690	16	16	16	F2:D2:S1	-0.02247	-0.01135	2	2	2
D3	0.00722	-0.01061	16	16	16	F3:D2:S1	-0.00932	-0.00337	2	2	2
E1	0.00765	-0.05026	24	24	24	F4:D2:S1	-0.02821	-0.03078	2	2	2
S1	0.11797	-0.00405	24	24	24	F2:D3:S1	0.00805	-0.00366	2	2	2
P1:F2	0.01079	0.00578	-12	0	0	F3:D3:S1	0.01135	0.01056	2	2	2
P2:F2	0.00172	0.00520	0	-12	0	F4:D3:S1	0.01334	0.00228	2	2	2
P3:F2	0.00915	0.00554	0	0	-12	P1:E1:S1	-0.00701	0.00069	-12	0	0
P1:F3	0.00670	0.00368	-12	0	0	P2:E1:S1	-0.01137	-0.01288	0	-12	0
P2:F3	-0.00379	0.00352	0	-12	0	P3:E1:S1	-0.00836	-0.00124	0	0	-12
P3:F3	0.00290	0.00280	0	0	-12	F2:E1:S1	-0.00561	0.00644	3	3	3
P1:F4	0.00852	0.00575	-12	0	0	F3:E1:S1	-0.02180	-0.00900	3	3	3
P2:F4	-0.00041	0.00452	0	-12	0	F4:E1:S1	0.03958	0.02374	3	3	3
P3:F4	0.00769	0.00642	0	0	-12	D2:E1:S1	0.00931	0.00566	4	4	4
P1:D2	0.01757	0.00264	-16	0	0	D3:E1:S1	0.05078	0.04154	4	4	4
P2:D2	0.00401	-0.00052	0	-16	0	P1:F2:D2:E1	0.03734	0.01509	-2	0	0
P3:D2	0.00485	0.00678	0	0	-16	P2:F2:D2:E1	0.01296	0.02702	0	-2	0
P1:D3	-0.02295	-0.01215	-16	0	0	P3:F2:D2:E1	-0.00739	-0.00683	0	0	-2
P2:D3	-0.06001	-0.07274	0	-16	0	P1:F3:D2:E1	0.02650	0.01755	-2	0	0
P3:D3	-0.00095	0.00907	0	0	-16	P2:F3:D2:E1	0.00549	0.03205	0	-2	0
F2:D2	0.00771	0.00420	4	4	4	P3:F3:D2:E1	-0.00878	-0.00335	0	0	-2
F3:D2	0.00903	0.00705	4	4	4	P1:F4:D2:E1	0.01622	0.00747	-2	0	0
F4:D2	-0.00071	0.00061	4	4	4	P2:F4:D2:E1	0.00529	0.02670	0	-2	0
F2:D3	-0.02408	-0.00006	4	4	4	P3:F4:D2:E1	-0.01882	-0.00812	0	0	-2
F3:D3	-0.01386	0.00868	4	4	4	P1:F2:D3:E1	-0.04176	-0.02704	-2	0	0
F4:D3	-0.04265	-0.02514	4	4	4	P2:F2:D3:E1	0.07127	0.11882	0	-2	0
P1:E1	-0.01688	-0.02401	-24	0	0	P3:F2:D3:E1	-0.04038	-0.02815	0	0	-2
P2:E1	0.00566	0.01336	0	-24	0	P1:F3:D3:E1	-0.03123	-0.01473	-2	0	0
P3:E1	0.00566	-0.00240	0	0	-24	P2:F3:D3:E1	0.06680	0.10899	0	-2	0
F2:E1	-0.03689	0.03150	6	6	6	P3:F3:D3:E1	-0.02832	-0.03552	0	0	-2
F3:E1	0.02150	0.01499	6	6	6	P1:F4:D3:E1	-0.04615	-0.03372	-2	0	0
F4:E1	-0.06042	0.01980	6	6	6	P2:F4:D3:E1	-0.03058	-0.02472	0	-2	0
D2:E1	-0.00439	-0.01031	8	8	8	P3:F4:D3:E1	-0.02570	-0.04034	0	0	-2
D3:E1	-0.04049	-0.03881	8	8	8	P1:F2:D2:S1	0.02769	0.00794	-2	0	0
P1:S1	0.01147	0.00334	-24	0	0	P2:F2:D2:S1	0.14667	0.14205	0	-2	0
P2:S1	-0.00663	-0.00812	0	-24	0	P3:F2:D2:S1	0.02404	0.02016	0	0	-2
P3:S1	0.00542	0.00170	0	0	-24	P1:F3:D2:S1	0.02292	0.01260	-2	0	0
F2:S1	-0.02037	-0.00832	6	6	6	P2:F3:D2:S1	0.12912	0.13675	0	-2	0
F3:S1	-0.11627	0.00533	6	6	6	P3:F3:D2:S1	0.00472	0.00639	0	0	-2
F4:S1	-0.02085	-0.02512	6	6	6	P1:F4:D2:S1	0.05832	0.05041	-2	0	0
D2:S1	0.00910	0.00030	8	8	8	P2:F4:D2:S1	-0.00869	-0.00287	0	-2	0
D3:S1	-0.01624	-0.01329	8	8	8	P3:F4:D2:S1	0.02224	0.03694	0	0	-2
E1:S1	0.00649	-0.00382	12	12	12	P1:F2:D3:S1	-0.01288	-0.00345	-2	0	0
P1:F2:D2	-0.01831	-0.00332	-4	0	0	P2:F2:D3:S1	-0.01396	0.01430	0	-2	0
P2:F2:D2	-0.00501	-0.00189	0	-4	0	P3:F2:D3:S1	0.01623	0.02089	0	0	-2
P3:F2:D2	-0.00391	-0.00503	0	0	-4	P1:F3:D3:S1	-0.02459	0.00289	-2	0	0
P1:F3:D2	-0.01561	-0.00405	-4	0	0	P2:F3:D3:S1	0.21926	0.26922	0	-2	0
P2:F3:D2	-0.00069	-0.00164	0	-4	0	P3:F3:D3:S1	-0.00413	-0.00461	0	0	-2
P3:F3:D2	-0.00034	-0.00705	0	0	-4	P1:F4:D3:S1	0.02709	0.05167	-2	0	0
P1:F4:D2	-0.01683	-0.00600	-4	0	0	P2:F4:D3:S1	-0.18091	-0.17894	0	-2	0
P2:F4:D2	-0.00673	-0.00852	0	-4	0	P3:F4:D3:S1	0.00778	0.01419	0	0	-2
P3:F4:D2	0.00627	-0.00263	0	0	-4	P1:F2:E1:S1	0.01962	0.00579	-3	0	0
P1:F2:D3	0.02174	0.01333	-4	0	0	P2:F2:E1:S1	0.01413	0.01229	0	-3	0
P2:F2:D3	0.06059	0.06655	0	-4	0	P3:F2:E1:S1	0.01888	0.01407	0	0	-3
P3:F2:D3	0.01702	-0.00029	0	0	-4	P1:F3:E1:S1	0.01940	0.01002	-3	0	0
P1:F3:D3	0.03274	0.01063	-4	0	0	P2:F3:E1:S1	0.02513	0.02382	0	-3	0
P2:F3:D3	0.07041	0.06891	0	-4	0	P3:F3:E1:S1	0.02301	0.01373	0	0	-3
P3:F3:D3	0.01022	-0.00825	0	0	-4	P1:F4:E1:S1	0.06140	0.06699	-3	0	0
P1:F4:D3	0.02682	0.00708	-4	0	0	P2:F4:E1:S1	-0.01491	-0.00450	0	-3	0
P2:F4:D3	0.04701	0.05246	0	-4	0	P3:F4:E1:S1	-0.01088	-0.00501	0	0	-3
P3:F4:D3	0.02732	0.02399	0	0	-4	P1:D2:E1:S1	0.02296	0.01593	-4	0	0
P1:F2:E1	-0.01126	-0.00378	-6	0	0	P2:D2:E1:S1	-0.01793	-0.01520	0	-4	0
P2:F2:E1	-0.01543	-0.02444	0	-6	0	P3:D2:E1:S1	-0.01578	-0.00198	0	0	-4
P3:F2:E1	-0.02510	-0.01880	0	0	-6	P1:D3:E1:S1	-0.07029	-0.02085	-4	0	0
P1:F3:E1	-0.00629	-0.00194	-6	0	0	P2:D3:E1:S1	0.09140	0.16368	0	-4	0
P2:F3:E1	-0.03463	-0.04123	0	-6	0	P3:D3:E1:S1	-0.01535	-0.03492	0	0	-4
P3:F3:E1	-0.03231	-0.02421	0	0	-6	F2:D2:E1:S1	0.01191	0.00461	1	1	1
P1:F4:E1	-0.00354	-0.00443	-6	0	0	F3:D2:E1:S1	0.00690	0.01695	1	1	1
P2:F4:E1	-0.00486	-0.01447	0	-6	0	F4:D2:E1:S1	-0.00313	0.01058	1	1	1
P3:F4:E1	-0.01679	-0.01378	0	0	-6	F2:D3:E1:S1	-0.05328	-0.03556	1	1	1
P1:D2:E1	-0.02094	-0.00716	-8	0	0	F3:D3:E1:S1	-0.02033	-0.02430	1	1	1
P2:D2:E1	-0.00947	-0.02233	0	-8	0	F4:D3:E1:S1	-0.03121	-0.02013	1	1	1
P3:D2:E1	0.01801	0.01635	0	0	-8	P1:F2:D2:E1:S1	-0.03417	-0.01113	-1	0	0
P1:D3:E1	0.04271	0.02513	-8	0	0	P2:F2:D2:E1:S1	-0.01006	0.00169	0	-1	0
P2:D3:E1	-0.06649	-0.10742	0	-8	0	P3:F2:D2:E1:S1	0.00708	-0.01190	0	0	-1
P3:D3:E1	0.03975	0.04894	0	0	-8	P1:F3:D2:E1:S1	-0.03752	-0.03731	-1	0	0
F2:D2:E1	-0.00502	0.00135	2	2	2	P2:F3:D2:E1:S1	0.01602	-0.00499	0	-1	0
F3:D2:E1	-0.00352	-0.00303	2	2	2	P3:F3:D2:E1:S1	0.00027	-0.01988	0	0	-1
F4:D2:E1	0.01407	0.00867	2	2	2	P1:F4:D2:E1:S1	-0.03991	-0.03174	-1	0	0
F2:D3:E1	0.03808	0.02488	2	2	2	P2:F4:D2:E1:S1	-0.26438	-0.30237	0	-1	0
F3:D3:E1	0.02224	0.02436	2	2	2	P3:F4:D2:E1:S1	0.01045	-0.01697	0	0	-1
F4:D3:E1	0.03351	0.03253	2	2	2	P1:F2:D3:E1:S1	0.08873	0.05392	-1	0	0
P1:F2:S1	-0.04063	-0.03238	-6	0	0	P2:F2:D3:E1:S1	-0.08777	-0.16810	0	-1	0
P2:F2:S1	0.01484	0.01894	0	-6	0	P3:F2:D3:E1:S1	0.00641	0.01366	0	0	-1
P3:F2:S1	-0.00773	-0.00613	0	0	-6	P1:F3:D3:E1:S1	0.05272	0.00698	-1	0	0
P1:F3:S1	-0.01374	-0.00476	-6	0	0	P2:F3:D3:E1:S1	-0.11177	-0.17661	0	-1	0
P2:F3:S1	0.00700	0.00634	0	-6	0	P3:F3:D3:E1:S1	-0.00881	0.01932	0	0	-1
P3:F3:S1	-0.00347	-0.00301	0	0	-6	P1:F4:D3:E1:S1	0.04961	0.00943	-1	0	0
P1:F4:S1	-0.09881	-0.09962	-6	0	0	P2:F4:D3:E1:S1	-0.06499	-0.11115	0	-1	0
P2:F4:S1	0.03659	0.03503	0	-6	0	P3:F4:D3:E1:S1	-0.00188	0.01291	0	0	-1

Table 4: Full table of F-Tests (the β_1 and β_2 columns contain coefficients for “Sales” and “Optimality”, respectively; the M -columns the matrices to compare the algorithm to the three rules of thumb)

Similarly, for $P1 : D3 : S1$ we have

$$\prod_{i \notin \{D, S\}} k_i = \prod_{i \in \{F, E\}} k_i = k_F \cdot k_E = 4 \cdot 2 = 8. \quad (\text{C.3})$$

As these contain $P1$ they belong to the part of the null hypothesis that is subtracted, hence they are equipped with a negative sign. As is easily seen, the intercept appears on both sides of the minus-sign exactly

$$\prod_{i \notin \emptyset} k_i = \prod_{i \in \{F, D, E, S\}} k_i = k_F \cdot k_D \cdot k_E \cdot k_S = 4 \cdot 3 \cdot 2 \cdot 2 = 48 \quad (\text{C.4})$$

times, and its coefficient is $48 - 48 = 0$.

References

- [1] Albers, S. (1998): Regeln für die Allokation eines Marketing-Budgets auf Produkte oder Marktsegmente, Schmalenbachs Zeitschrift für betriebswirtschaftliche Forschung, 50(3): 211-235
- [2] Beswick, C.A.; Cravens, D.W. (1977): A Multistage decision Model for sales force management. Journal of Marketing Research 14(2): 135-144
- [3] Doyle, P.; Saunders, J. (1990): Multiproduct Advertising Budgeting. Marketing Science 9(2): 97-113
- [4] Goldfarb, D.; Idnani, A. (1983): A numerically stable dual method for solving strictly convex quadratic programs, Columbia University, Department of Industrial Engineering and Operations Research. New York
- [5] Gupta, S.; Steenburgh, T. (2008): Allocating Marketing Resources. Working Paper, Harvard Business School, Boston, MA
- [6] Hanssens, D.M.; Parsons, L.J.; Schultz, R.L. (2001): Market Response Models. Econometric and Time Series Analysis, 2nd Edition, Kluwer Academic Publishers, Boston, MA.
- [7] Kotzan, J.A.; Evanson, R.V. (1969): Responsiveness of Drug Store Sales to Shelf Space Allocations. Journal of Marketing Research 6(4): 465-469
- [8] LaForge, R.; Cravens, D.W. (1985): Empirical and judgment based sales force decision models: A comparative analysis. Decision Sciences 16: 177-195
- [9] Mantrala, M.K. (2006): Allocating Marketing Resources. in: Handbook of Marketing, herausgegeben von Weitz, B.; Wensley, R., London: SAGE
- [10] Mantrala, M.K.; Sinha, P.; Zoltners, A.A. (1992): Impact of Resource Allocation Rules on Marketing Investment-Level Decisions and Profitability. Journal of Marketing Research 52(2): 147-165
- [11] March, J. G. (1991): Exploration and Exploitation in Organizational Learning. Organization Science 2.1: 71-87
- [12] Rudin, W. (1976): Principles of Mathematical Analysis, McGraw-Hill, New York

- [13] Seldon, J.A. (1986): A Note on the Teaching of Arc Elasticity. *Journal of economic education* 12(2): 120-124
- [14] Sinha, P.; Zoltners, A.A. (2001): Sales-force decision models: Insights from 25 years of implementation. *Interfaces* 31(3): S8-S44
- [15] Vázquez, A. (1995): A Note on the Arc Elasticity of Demand. *Estudios Económicos* 10(2): 221-228