

Integral foliated simplicial volume of hyperbolic 3-manifolds

Clara Löh and Cristina Pagliantini

Abstract. Integral foliated simplicial volume is a version of simplicial volume combining the rigidity of integral coefficients with the flexibility of measure spaces. In this article, using the language of measure equivalence of groups we prove a proportionality principle for integral foliated simplicial volume for aspherical manifolds and give refined upper bounds of integral foliated simplicial volume in terms of stable integral simplicial volume. This allows us to compute the integral foliated simplicial volume of hyperbolic 3-manifolds. This is complemented by the calculation of the integral foliated simplicial volume of Seifert 3-manifolds.

Mathematics Subject Classification (2010). 57M27, 57M50, 20F67, 55N99.

Keywords. Simplicial volume, integral foliated simplicial volume, hyperbolic 3-manifolds, measure equivalence.

Contents

1	Introduction	826
2	Simplicial volume and (stable) integral simplicial volume	829
3	Hyperbolic 3-manifolds with small stable integral simplicial volume	830
4	Integral foliated simplicial volume	834
5	A proportionality principle for integral foliated simplicial volume	847
6	Comparing integral foliated simplicial volume and stable integral simplicial volume	855
7	Integral foliated simplicial volume of hyperbolic 3-manifolds	860
8	Stable integral simplicial volume and integral foliated simplicial volume of Seifert 3-manifolds	861
	References	862

1. Introduction

Integral foliated simplicial volume is a version of simplicial volume combining the rigidity of integral coefficients with the flexibility of measure spaces. If M is an oriented closed connected manifold, then the integral foliated simplicial volume $|M|$ fits into the sandwich

$$\|M\| \leq |M| \leq \|M\|_{\mathbb{Z}}^{\infty}, \quad (1)$$

where $\|M\|$ is the classical simplicial volume [13] and where $\|M\|_{\mathbb{Z}}^{\infty}$ is the stable integral simplicial volume (stabilised over all finite coverings of M).

Gromov [15, p. 305f] suggested a definition of integral foliated simplicial volume and an upper estimate of L^2 -Betti numbers in terms of integral foliated simplicial volume, which was confirmed by Schmidt [29]. It is an open problem whether integral foliated simplicial volume coincides with simplicial volume in the case of aspherical manifolds; an affirmative answer would show that aspherical manifolds with vanishing simplicial volume have vanishing Euler characteristic, which is a long-standing open problem [14, p. 232]. The only cases in which the integral foliated simplicial volume has been computed are manifolds that split off an S^1 -factor and simply connected manifolds [29, Chapter 5.2]. Moreover, Sauer proved an upper bound of a related invariant in terms of minimal volume [28, Section 3].

In the present article, we will prove the following:

Theorem 1.1 (integral foliated simplicial volume of hyperbolic 3-manifolds). *For all oriented closed connected hyperbolic 3-manifolds M the integral foliated simplicial volume and the simplicial volume of M coincide:*

$$|M| = \|M\| = \frac{\text{vol}(M)}{v_3}.$$

Here, v_3 denotes the maximal volume of ideal geodesic 3-simplices in \mathbb{H}^3 .

The second equality in Theorem 1.1 is the classic proportionality principle for hyperbolic manifolds of Gromov [13] and Thurston [32], which also holds in more generality [7, 10, 11, 5, 30, 4, 9, 1, 20].

The proof of Theorem 1.1 consists of the following steps. Using the language of measure equivalence of groups and techniques of Bader, Furman and Sauer [1] we show that, similarly to classical simplicial volume [1, Theorem 1.9], also integral foliated simplicial volume of aspherical manifolds satisfies a proportionality principle with respect to certain parameter spaces (see Section 5 and Section 4 for the definitions):

Theorem 1.2 (proportionality principle for integral foliated simplicial volume). *Let M and N be oriented closed connected aspherical manifolds of the same dimension satisfying $\|M\| > 0$ and $\|N\| > 0$. Suppose that there exists an ergodic bounded measure equivalence coupling (Ω, μ) of the fundamental groups Γ and Λ of M and N , respectively; let c_Ω be the coupling index of this coupling.*

(1) Then

$$|M|^{\Lambda \setminus \Omega} = c_\Omega \cdot |N|^{\Gamma \setminus \Omega}.$$

(2) If the coupling (Ω, μ) is mixing, then

$$|M| = c_\Omega \cdot |N|.$$

Considering locally symmetric spaces of non-compact type, we obtain:

Corollary 1.3 (a proportionality principle for integral foliated simplicial volume of locally symmetric spaces of non-compact type). *Let X be an aspherical symmetric space of non-compact type and let $\Gamma, \Lambda < G = \text{Isom}^0(X)$ be uniform lattices. Then*

$$\frac{|X/\Gamma|^{G/\Lambda}}{\text{covol}(\Gamma)} = \frac{|X/\Lambda|^{G/\Gamma}}{\text{covol}(\Lambda)}$$

and

$$\frac{|X/\Gamma|}{\text{covol}(\Gamma)} = \frac{|X/\Lambda|}{\text{covol}(\Lambda)}.$$

In particular, the coupling of uniform hyperbolic lattices given by the isometry group satisfies Corollary 1.3. Therefore, when estimating integral foliated simplicial volume of a hyperbolic manifold, we can call other hyperbolic manifolds for help.

In particular, we obtain the following refinement of the upper bound of integral foliated simplicial volume in terms of stable integral simplicial volume:

Corollary 1.4 (comparing integral foliated simplicial volume and stable integral simplicial volume, hyperbolic case). *Let $n \in \mathbb{N}$, and let M and N be oriented closed connected hyperbolic n -manifolds. Then*

$$|M| \leq \frac{\text{vol}(M)}{\text{vol}(N)} \cdot \|N\|_{\mathbb{Z}}^\infty.$$

Moreover, we exhibit concrete examples of parameter spaces that realize stable integral simplicial volume as integral foliated simplicial volume.

Theorem 1.5 (comparing integral foliated simplicial volume and stable integral simplicial volume, generic case). *Let M be an oriented closed connected manifold with fundamental group Γ and let S be the set of finite index subgroups of Γ . Then*

$$\|M\|_{\mathbb{Z}}^{\infty} = |M|^{\prod_{\Lambda \in S} \Gamma/\Lambda}.$$

While Theorem 1.5 is not necessary to prove Theorem 1.1, it is of independent interest in the context of Question 1.7.

As last step in the proof of Theorem 1.1, in dimension 3, we will use the following sequence of hyperbolic manifolds, based on a variation of a result by Francaviglia, Frigerio and Martelli [8]:

Theorem 1.6 (hyperbolic 3-manifolds with small stable integral simplicial volume). *There exists a sequence $(M_n)_{n \in \mathbb{N}}$ of oriented closed connected hyperbolic 3-manifolds with*

$$\lim_{n \rightarrow \infty} \frac{\|M_n\|_{\mathbb{Z}}^{\infty}}{\|M_n\|} = 1.$$

Notice that it is still unknown whether stable integral simplicial volume and simplicial volume coincide for hyperbolic 3-manifolds. Francaviglia, Frigerio, and Martelli [8] proved that in all dimensions bigger than 3 a sequence of hyperbolic manifolds as in Theorem 1.6 does *not* exist. Therefore, our approach does not allow to compute the integral foliated simplicial volume of higher-dimensional hyperbolic manifolds. Moreover, we do not know whether integral foliated simplicial volume can be different from stable integral simplicial volume for aspherical manifolds with enough finite coverings:

Question 1.7. *What is the difference between integral foliated simplicial volume and stable integral simplicial volume of aspherical oriented closed connected manifolds with residually finite fundamental group?*

In dimension 2 the situation is completely understood. Indeed, as an application of inequality (1), we show that the integral foliated simplicial volume of an oriented closed connected surface M of genus bigger than 0 is equal to both the simplicial volume and the stable integral simplicial volume (see Example 6.2).

Organisation of this article. In Section 2, we recall the definition and basic properties of (stable integral) simplicial volume. In Section 3, we construct hyperbolic 3-manifolds with small stable integral simplicial volume, which proves Theorem 1.6. Section 4 is an introduction into integral foliated simplicial volume and basic operations on parameter spaces. We prove the proportionality principles Theorem 1.2 and Corollary 1.3 in Section 5. Section 6 is devoted to the refinements of the comparison between integral foliated simplicial volume and stable integral

simplicial volume and includes a proof of Theorem 1.5. Finally, in Section 7, we complete the proof of Corollary 1.4 and Theorem 1.1. Section 8 contains the computation of integral foliated simplicial volume of Seifert manifolds.

Acknowledgements. We would like to thank Roman Sauer, Roberto Frigerio, Bruno Martelli, and Marco Schmidt for numerous helpful discussions. In particular, we would like to thank Roman Sauer for pointing out a mistake in the first version. This work was partially supported by the Graduiertenkolleg *Curvature, Cycles, and Cohomology* (Universität Regensburg). The second author was also partially supported by the Swiss National Science Foundation, under the grant 2000200-144373.

2. Simplicial volume and (stable) integral simplicial volume

In this section we will recall the definition of simplicial volume introduced by Gromov [13, 19] and its integral version, which uses integral homology instead of real homology.

Let X be a topological space. Let R be a normed ring. In this section, we restrict only to the cases $R = \mathbb{R}$ or \mathbb{Z} . For $i \in \mathbb{N}$ we denote by $S_i(X)$ the set of singular i -simplices in X , by $C_i(X, R)$ the module of singular i -chains with R -coefficients. The homology of the complex $(C_*(X, R), \partial_*)$, where ∂_* is the usual differential, is the singular homology $H_*(X, R)$ of X with coefficients in R .

We endow the R -module $C_i(X, R)$ with the ℓ^1 -norm defined by

$$\left\| \sum_{\sigma \in S_i(X)} a_\sigma \cdot \sigma \right\|_1^R = \sum_{\sigma \in S_i(X)} |a_\sigma|,$$

where $|\cdot|$ is the norm on R . We denote the norm $\|\cdot\|_1^R$ simply by $\|\cdot\|_1$. The norm $\|\cdot\|_1^R$ descends to a semi-norm on $H_*(X, R)$, which is also denoted by $\|\cdot\|_1^R$ and is defined as follows: if $\alpha \in H_i(X, R)$, then

$$\|\alpha\|_1^R = \inf\{\|c\|_1^R \mid c \in C_i(X, R), \partial c = 0, [c] = \alpha\}.$$

Note that $\|\cdot\|_1^{\mathbb{Z}}$ on $H_*(\cdot, \mathbb{Z})$ is technically not a semi-norm as it is not multiplicative in general (see below).

If M is a closed connected oriented n -manifold, then we denote the fundamental class of M by $[M]_{\mathbb{Z}}$, i.e., the positive generator of $H_n(M, \mathbb{Z}) \cong \mathbb{Z}$. The change of coefficients homomorphism $H_n(M, \mathbb{Z}) \rightarrow H_n(M, \mathbb{R})$ sends the fundamental class to the real fundamental class $[M]_{\mathbb{R}} \in H_n(M, \mathbb{R})$ of M . The following definition is due to Gromov [13].

Definition 2.1 ((integral) simplicial volume). The *simplicial volume* of M is

$$\|M\| := \|[M]_{\mathbb{R}}\|_1 \in \mathbb{R}_{\geq 0}.$$

The *integral simplicial volume* of M is defined as $\|M\|_{\mathbb{Z}} := \|[M]_{\mathbb{Z}}\|_1^{\mathbb{Z}} \in \mathbb{N}$.

Of course we have the inequality $\|M\| \leq \|M\|_{\mathbb{Z}}$ but in general no equality (for instance, $\|S^1\| = 0$ but $\|S^1\|_{\mathbb{Z}} = 1$). The integral simplicial volume does not behave as nicely as the simplicial volume. For example, it follows from the definition that $\|M\|_{\mathbb{Z}} \geq 1$ for every manifold M . Therefore, the integral simplicial volume cannot be multiplicative with respect to finite coverings (otherwise it would vanish on manifolds that admit finite non-trivial self-coverings, such as S^1). Moreover, as we mentioned before, the ℓ^1 -semi-norm on integral homology is not really a semi-norm, since the equality $\|n \cdot \alpha\|_1^{\mathbb{Z}} = |n| \cdot \|\alpha\|_1^{\mathbb{Z}}$, may not hold for all $\alpha \in H_*(X; \mathbb{Z})$ and all $n \in \mathbb{Z}$. Indeed, it is easy to see that $\|n \cdot [S^1]_{\mathbb{Z}}\|_1^{\mathbb{Z}} = 1$ for every $n \in \mathbb{Z} \setminus \{0\}$.

We may consider a *stable* version of the integral simplicial volume:

Definition 2.2 (stable integral simplicial volume). The *stable integral simplicial volume* of an oriented closed connected manifold M is

$$\|M\|_{\mathbb{Z}}^{\infty} := \inf \left\{ \frac{1}{d} \cdot \|\bar{M}\|_{\mathbb{Z}} \mid d \in \mathbb{N}, \text{ there is a } d\text{-sheeted covering } \bar{M} \rightarrow M \right\}.$$

Since the simplicial volume is multiplicative under finite coverings [19, Proposition 4.1], it is clear that $\|M\| \leq \|M\|_{\mathbb{Z}}^{\infty}$, but in general they are not equal:

Theorem 2.3 ([8, Theorem 2.1]). *For every $n \in \mathbb{N}_{\geq 4}$ there exists a constant $C_n < 1$ such that the following holds. Let M be an oriented closed connected hyperbolic n -manifold. Then*

$$\|M\| \leq C_n \cdot \|M\|_{\mathbb{Z}}^{\infty}.$$

For hyperbolic 3-manifolds it is still an open question whether the simplicial volume and the stable integral simplicial volume are the same: our Theorem 1.6 gives a partial answer.

3. Hyperbolic 3-manifolds with small stable integral simplicial volume

In this section we will prove Theorem 1.6 following an argument of Francaviglia, Frigerio, and Martelli [8, Corollary 5.16]. For the sake of completeness, we recall some background on triangulations and special complexity.

Definition 3.1 (triangulation). A *triangulation* of a closed 3-manifold M is a realization of the manifold M as the gluing of finitely many tetrahedra via some simplicial pairing of their faces. Turning to the case of a compact manifold M with non-empty boundary ∂M , one can adapt the notion of triangulation: an *ideal triangulation* of M is a decomposition of its interior $\text{Int}(M)$ into tetrahedra with their vertices removed. An (ideal) triangulation is *semi-simplicial* if all the edges have distinct vertices.

Definition 3.2 (special complexity). Let M be a compact 3-manifold, possibly with boundary. The *special complexity* $c_S(M)$ of M is the minimal number of vertices in a special spine for M .

We refer to the works of Matveev [24, 25] for the definition of special spines and their properties. For our purpose we just need to recall that a special spine is dual to a triangulation. In particular, there is a bijection between the simplices in a triangulation and the vertices in the dual special spine [25, Theorem 1.1.26 and Corollary 1.1.27]. With an abuse of notation we call the *true* vertices of a special spine in Matveev's definition simply vertices of a special spine.

Theorem 3.3 (Matveev [25, Corollary 1.1.28]). *Let M be a compact 3-manifold whose interior $\text{Int}(M)$ admits a complete finite volume hyperbolic structure. Then there is a bijection between special spines and ideal triangulations of M such that the number of vertices in the special spine is equal to the number of tetrahedra in the corresponding triangulation.*

Remark 3.4. Matveev [24] introduced the more general notion of complexity, which involves spines that are not necessarily dual to triangulations and showed that complexity and special complexity are equal for any closed orientable irreducible 3-manifold distinct from S^3 , \mathbb{RP}^3 , and $L(3, 1)$.

Remark 3.5. Matveev [24] and Martelli [21] used two slightly different definitions of spines but the two notions are equivalent and lead to the same complexity [21, Section 7]. Coherently with [8] we use Martelli's definition.

Remark 3.6. Let M be an oriented closed connected manifold and let T be a semi-simplicial triangulation of M . We fix an order of the vertices of T . Parametrising the tetrahedra according to this order yields singular simplices $\sigma_1, \dots, \sigma_k$ of M . We then choose signs $\varepsilon_1, \dots, \varepsilon_k \in \{1, -1\}$ indicating whether these singular simplices $\sigma_1, \dots, \sigma_k$ are compatible with the orientation on M or not. Then $\sum_{j=1}^k \varepsilon_j \cdot \sigma_j$ is a fundamental cycle of M . Therefore, the number of tetrahedra in a semi-simplicial triangulation provides an upper bound for the integral simplicial volume of M .

In order to adapt the proof of Francaviglia, Frigerio and Martelli [8, Corollary 5.16] we need the following results.

Proposition 3.7. *Let $\bar{M}_{(5)}$ be the compactification of the 5-chain link complement $M_{(5)}$. Then we have*

$$c_S(\bar{M}_{(5)}) = 10 = \|\bar{M}_{(5)}\|.$$

Moreover, the value of $c_S(\bar{M}_{(5)})$ is realized by a special spine dual to a semi-simplicial triangulation.

Proof. The 5-chain link complement has a hyperbolic structure [27] and admits an ideal triangulation with 10 ideal and regular tetrahedra such that each edge has vertices in different cusps [26, Section 5.2]. By the proportionality between simplicial volume and Riemannian volume (which holds both in the compact case [13, 32] and in the cusped case [7, 10, 11, 5]) we have

$$\|\bar{M}_{(5)}\| = \frac{\text{vol}(M_{(5)})}{v_3} = 10.$$

Moreover, Theorem 3.3 implies

$$c_S(\bar{M}_{(5)}) \leq 10.$$

The equality follows from the fact that for every oriented connected finite volume hyperbolic 3-manifold M with compactification \bar{M} the inequality $\|\bar{M}\| \leq c_S(\bar{M})$ holds. Indeed, an argument by Francaviglia [7, Theorem 1.2 and Proposition 3.8] guarantees that the volume of M can be computed by straightening any ideal triangulation of M and then summing the volume of the straight version of the tetrahedra. \square

Proposition 3.8. *Let N be the compactification of a finite volume oriented connected hyperbolic 3-manifold and suppose that N admits a semi-simplicial triangulation that realizes the value of $c_S(N)$. Let M be a manifold obtained by Dehn filling on N . Then*

$$\|M\|_{\mathbb{Z}}^{\infty} \leq c_S(N).$$

Proof. As pointed out in Remark 3.6 we estimate the integral simplicial volume by the number of vertices of the special spine dual to a semi-simplicial triangulation.

From the special spine P dual to a semi-simplicial triangulation of N that realizes $c_S(N)$, we construct a special spine for M , and hence for its finite coverings, such that the associated triangulations are still semi-simplicial. We obtain $\|M\|_{\mathbb{Z}}^{\infty} \leq c_S(N)$ following step by step the argument of Francaviglia, Frigerio, and Martelli [8, Proposition 5.15].

More precisely, let T_1, \dots, T_k be the boundary tori of N . For every $i \in \{1, \dots, k\}$ let V_i be an open solid torus in $M \setminus P$ created by Dehn filling on the boundary component T_i . Let D_i^1 and D_i^2 be a pair of parallel meridian discs of V_i . If D_i^1 and D_i^2 are generic with respect to the cellularization induced by P on T_i [8, Lemma 5.9], the spine $P \cup D_i^1 \cup D_i^2$ is special, dual to a semi-simplicial triangulation, and with $v_\partial^{T_i}$ vertices added to the ones of P . Gluing a pair of parallel discs for each boundary torus T_i we obtain a special spine $Q = P \cup \bigcup_{i=1}^k (D_i^1 \cup D_i^2)$ for M with $c_S(N) + \sum_{i=1}^k v_\partial^{T_i} + v_I$ vertices, where v_I is the number of vertices created by intersections between discs added in different boundary components.

Since $\pi_1(M)$ is residually finite, for every $n > 0$ there exist $n_0 > n, h > 0$ and a regular covering $p: \bar{M} \rightarrow M$ of degree hn_0 such that, for every $i \in \{1, \dots, k\}$, the preimage $p^{-1}(V_i)$ consists of h open solid tori $\bar{V}_i^1, \dots, \bar{V}_i^h$ each winding n_0 times along V_i via p . The special spine Q of M lifts to a special spine $\bar{Q} := p^{-1}(Q)$ of \bar{M} . In particular, each pair of parallel discs added to P lifts to n_0 copies of pairs of parallel discs in each open solid torus \bar{V}_i^j . Removing $2n_0 - 2$ discs for each open solid torus in \bar{Q} , we obtain again a special spine \bar{Q}' dual to a semi-simplicial triangulation of \bar{M} .

By Remark 3.6 we now estimate $\|\bar{M}\|_{\mathbb{Z}}$ with the number of vertices of \bar{Q}' :

$$\|M\|_{\mathbb{Z}}^\infty \leq \frac{\|\bar{M}\|_{\mathbb{Z}}}{hn_0} \leq c_S(N) + \frac{\sum_{i=1}^k v_\partial^{T_i} + v_I}{n_0}.$$

Since this holds for every $n > 0$ and since $n_0 > n$, we get the conclusion. □

We can now complete the proof of Theorem 1.6.

Proof of Theorem 1.6. Let $(M_n)_{n \in \mathbb{N}}$ be a family of hyperbolic 3-manifolds obtained by Dehn filling on $\bar{M}_{(5)}$. By Thurston’s Dehn filling Theorem [32, Chapter 5, page 118] it follows that $\lim_{n \rightarrow \infty} \text{vol}(M_n) = \text{vol}(M_{(5)})$, which implies $\lim_{n \rightarrow \infty} \|M_n\| = \|\bar{M}_{(5)}\|$ using the proportionality principle for hyperbolic manifolds (which holds both in the compact case [13, 32] and in the cusped case [7, 10, 11, 5]). Then we have

$$1 \leq \frac{\|M_n\|_{\mathbb{Z}}^\infty}{\|M_n\|} \leq \frac{c_S(\bar{M}_{(5)})}{\|M_n\|} \xrightarrow{n \rightarrow \infty} \frac{c_S(\bar{M}_{(5)})}{\|\bar{M}_{(5)}\|} = 1.$$

where the second inequality follows by Proposition 3.8, and the last equality by Proposition 3.7. □

4. Integral foliated simplicial volume

In the following, we will recall the precise definition of integral foliated simplicial volume by Schmidt [29] and discuss basic facts about the effect of changing parameter spaces.

4.1. Definition of integral foliated simplicial volume. Integral foliated simplicial volume is defined via homology with twisted coefficients in function spaces of probability spaces that carry an action of the fundamental group. Background on the well-behaved category of standard Borel spaces can be found in the book by Kechris [16].

Definition 4.1 (parametrised fundamental cycles). Let M be an oriented closed connected n -manifold with fundamental group Γ and universal covering $\tilde{M} \rightarrow M$.

- A *standard Borel space* is a measurable space that is isomorphic to a Polish space with its Borel σ -algebra. A *standard Borel probability space* is a standard Borel space together with a probability measure.
- A *standard Γ -space* is a standard Borel probability space (X, μ) together with a measurable μ -preserving (left) Γ -action. If the probability measure is clear from the context, we will abbreviate (X, μ) by X .
- If (X, μ) is a standard Γ -space, then we equip $L^\infty(X, \mu, \mathbb{Z})$ with the right Γ -action

$$\begin{aligned} L^\infty(X, \mu, \mathbb{Z}) \times \Gamma &\longrightarrow L^\infty(X, \mu, \mathbb{Z}), \\ (f, g) &\longmapsto (x \mapsto (f \cdot g)(x) := f(g \cdot x)), \end{aligned}$$

and we write i_M^X for the change of coefficients homomorphism

$$\begin{aligned} i_M^X: C_*(M, \mathbb{Z}) \cong \mathbb{Z} \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}) &\longrightarrow L^\infty(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}), \\ 1 \otimes c &\longmapsto 1 \otimes c, \end{aligned}$$

induced by the inclusion $\mathbb{Z} \hookrightarrow L^\infty(X, \mathbb{Z})$ as constant functions.

- If (X, μ) is a standard Γ -space, then

$$\begin{aligned} [M]^X &:= H_n(i_M^X)([M]_{\mathbb{Z}}) \in H_n(M, L^\infty(X, \mathbb{Z})), \\ &= H_n(L^\infty(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z})), \end{aligned}$$

is the X -parametrised fundamental class of M . All cycles in the chain complex $C_*(M, L^\infty(X, \mathbb{Z})) = L^\infty(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z})$ representing $[M]^X$ are called X -parametrised fundamental cycles of M .

The integral foliated simplicial volume is now defined as the infimum of ℓ^1 -norms over all parametrised fundamental cycles.

Definition 4.2 (integral foliated simplicial volume). Let M be an oriented closed connected n -manifold with fundamental group Γ , and let (X, μ) be a standard Γ -space.

- Let $\sum_{j=1}^k f_j \otimes \sigma_j \in C_*(M, L^\infty(X, \mathbb{Z}))$ be a chain in *reduced form*, i.e., the singular simplices $\sigma_1, \dots, \sigma_k$ on \tilde{M} satisfy $\pi \circ \sigma_j \neq \pi \circ \sigma_\ell$ for all $j, \ell \in \{1, \dots, k\}$ with $j \neq \ell$ (where $\pi: \tilde{M} \rightarrow M$ is the universal covering map). Then we define

$$\left| \sum_{j=1}^k f_j \otimes \sigma_j \right|^X := \sum_{j=1}^k \int_X |f_j| d\mu \in \mathbb{R}_{\geq 0}.$$

- The X -parametrised simplicial volume of M , denoted by $|M|^X$, is the infimum of the ℓ^1 -norms of all X -parametrised fundamental cycles of M .
- The *integral foliated simplicial volume* of M , denoted by $|M|$, is the infimum of all $|M|^X$ over all isomorphism classes of standard Γ -spaces X .

Remark 4.3. Let Γ be a countable group. The class of isomorphism classes of standard Γ -spaces indeed forms a set [29, Remark 5.26].

Remark 4.4. Schmidt’s original definition [29, Definition 5.25] requires the actions of the fundamental group on the parameter spaces to be essentially free. However, allowing also parameter spaces with actions that are not essentially free does not change the infimum (Corollary 4.14).

Example 4.5 (trivial parameter space). Let M be an oriented closed connected manifold with fundamental group Γ . If (X, μ) is a standard Γ -space consisting of a single point, then $L^\infty(X, \mathbb{Z}) \cong \mathbb{Z}$ (as $\mathbb{Z}\Gamma$ -modules with trivial Γ -action) and so

$$|M|^X = \|M\|_{\mathbb{Z}}.$$

More generally, in combination with Proposition 4.13 (2), we obtain: if (X, μ) is a standard Γ -space with trivial Γ -action, then

$$|M|^X = \|M\|_{\mathbb{Z}}.$$

In particular, if M is simply connected, then $|M|^X = \|M\|_{\mathbb{Z}}$ for all standard Borel probability spaces (X, μ) [29, Proposition 5.29].

Proposition 4.6 (comparison with (integral) simplicial volume [29, Remark 5.23]). *Let M be an oriented closed connected n -manifold with fundamental group Γ , and let (X, μ) be a standard Γ -space. Then*

$$\|M\| \leq |M| \leq |M|^X \leq \|M\|_{\mathbb{Z}}.$$

Proof. The linear map $L^\infty(X, \mathbb{Z}) \rightarrow \mathbb{R}$ given by integration with respect to μ maps the constant function 1 to 1 and is norm non-increasing with respect to the ℓ^1 -norm on $L^\infty(X, \mathbb{Z})$. From the first property, we easily deduce that the induced map $C_n(M, L^\infty(X, \mathbb{Z})) \rightarrow C_n(M, \mathbb{R})$ maps X -parametrised fundamental cycles to \mathbb{R} -fundamental cycles, and so

$$\|M\| \leq |M|^X.$$

Since the inequality holds for every standard Γ -space X , we have

$$\|M\| \leq |M|.$$

The inclusion $\mathbb{Z} \hookrightarrow L^\infty(X, \mathbb{Z})$ as constant functions is isometric with respect to the ℓ^1 -norm and the induced map $C_n(M, \mathbb{Z}) \rightarrow C_n(M, L^\infty(X, \mathbb{Z}))$ maps fundamental cycles to X -parametrised fundamental cycles. Hence,

$$|M|^X \leq \|M\|_{\mathbb{Z}}. \quad \square$$

Remark 4.7 (real coefficients). Arguments analogous to the ones in the proof of the previous proposition show that

$$|M|^{L^\infty(X, \mu, \mathbb{R})} = \|M\|$$

holds for all oriented closed connected manifolds M and all standard $\pi_1(M)$ -spaces (X, μ) . Here, $|M|^{L^\infty(X, \mu, \mathbb{R})}$ denotes the number defined like $|M|^X$, but using $L^\infty(X, \mu, \mathbb{R})$ instead of $L^\infty(X, \mu, \mathbb{Z})$.

We recall two “indecomposability” notions for parameter spaces from ergodic theory:

Definition 4.8 (ergodic/mixing parameter spaces). Let Γ be a countable group.

- A standard Γ -space (X, μ) is *ergodic* if every Γ -invariant measurable subset $A \subset X$ satisfies $\mu(A) \in \{0, 1\}$ (equivalently, $L^\infty(X, \mu, \mathbb{Z})^\Gamma$ contains only the constant functions).
- A standard Γ -space (X, μ) is called *mixing* if for all measurable subsets $A, B \subset X$ and all sequences $(g_n)_{n \in \mathbb{N}}$ in Γ with $\lim_{n \rightarrow \infty} g_n = \infty$ we have

$$\lim_{n \rightarrow \infty} \mu(A \cap g_n \cdot B) = \mu(A) \cdot \mu(B).$$

Here, $\lim_{n \rightarrow \infty} g_n = \infty$ means that the sequence $(g_n)_{n \in \mathbb{N}}$ eventually leaves any finite subset of Γ , i.e., that for all finite subsets $F \subset \Gamma$ there is an $N \in \mathbb{N}$ such that for all $n \in \mathbb{N}_{\geq N}$ we have $g_n \in \Gamma \setminus F$.

Clearly, for infinite discrete groups, any mixing parameter space is also ergodic. Moreover, any countably infinite group admits an essentially free mixing parameter space:

Example 4.9 (Bernoulli shift). The *Bernoulli shift* of a countable group Γ is the standard Borel space $(\{0, 1\}^\Gamma, \otimes_\Gamma (1/2 \cdot \delta_0 + 1/2 \cdot \delta_1))$, endowed with the translation action. If Γ is infinite, this standard Γ -space is essentially free and mixing (and hence ergodic) [29, Lemma 3.37].

For ergodic parameter spaces, the parametrised fundamental class indeed is a generator of the corresponding top homology with twisted coefficients:

Remark 4.10. Let M be an oriented closed connected n -manifold with fundamental group Γ , and let (X, μ) be an ergodic standard Γ -space. Then the inclusion $\mathbb{Z} \hookrightarrow L^\infty(X, \mathbb{Z})^\Gamma$ as constant functions is an isomorphism, and so the change of coefficients homomorphism

$$H_n(i_M^X): \mathbb{Z} \cong H_n(M, \mathbb{Z}) \longrightarrow H_n(M, L^\infty(X, \mathbb{Z})) \cong L^\infty(X, \mathbb{Z})^\Gamma \cong \mathbb{Z}$$

is an isomorphism; the isomorphism $H_n(M, L^\infty(X, \mathbb{Z})) \cong L^\infty(X, \mathbb{Z})^\Gamma$ is a consequence of Poincaré duality with twisted coefficients [34, Theorem 2.1, p. 23].

Furthermore, we will see that ergodic parameters suffice to describe the integral foliated simplicial volume (Proposition 4.17).

Remark 4.11 (lack of functoriality). Ordinary simplicial volume has the following functoriality property: if $f: M \rightarrow N$ is a continuous map between oriented closed connected manifolds of the same dimension of degree d , then

$$|d| \cdot \|N\| = \|d \cdot [N]_{\mathbb{R}}\|_1 = \|H_*(f, \mathbb{R})([M]_{\mathbb{R}})\|_1 \leq \|M\|.$$

However, when dealing with integral coefficients, the first equality might fail in general (because we will not be able to divide representatives of $d \cdot [N]_{\mathbb{Z}}$ by d). Therefore, integral simplicial volume, stable integral simplicial volume and integral foliated simplicial volume suffer from a lack of good estimates in terms of mapping degrees.

In the following, we will investigate some of the effects of changing parameter spaces. To this end, we will use the following comparison mechanism:

Proposition 4.12 (comparing parameter spaces). *Let M be an oriented closed connected n -manifold with fundamental group Γ , let (X, μ) and (Y, ν) be standard Γ -spaces, and let $\varphi: X \rightarrow Y$ be a measurable Γ -map. Moreover, suppose that*

$$\mu(\varphi^{-1}(A)) \leq \nu(A) \tag{2}$$

holds for all measurable sets $A \subset Y$. Then

$$|M|^X \leq |M|^Y.$$

Proof. We consider the (well-defined) chain map $\Phi := L^\infty(\varphi, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} \text{id}_{C_*(\tilde{M}, \mathbb{Z})}$:

$$\begin{aligned} \Phi: L^\infty(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}) &\longrightarrow L^\infty(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}) \\ f \otimes \sigma &\longmapsto f \circ \varphi \otimes \sigma. \end{aligned}$$

In view of the compatibility of φ with the measures (equation (2)), we see that

$$|\Phi(c)|^X \leq |c|^Y$$

holds for all chains $c \in L^\infty(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z})$. Moreover, equation (2) shows that φ is ν -almost surjective. In particular, $L^\infty(\varphi, \mathbb{Z})$ maps ν -almost constant functions to μ -almost constant functions (with the same value). From this we can easily conclude that Φ maps Y -parametrised fundamental cycles to X -parametrised fundamental cycles. Taking the infimum over all Y -parametrised fundamental cycles of M thus leads to $|M|^X \leq |M|^Y$. \square

We will now consider products, convex combinations, ergodic decomposition and induction/restriction of parameter spaces.

4.2. Products of parameter spaces

Proposition 4.13 (products of parameter spaces). *Let M be an oriented closed connected n -manifold with fundamental group Γ .*

- (1) *Let I be a non-empty, countable (or finite) set. If $(X_i, \mu_i)_{i \in I}$ is a family of standard Γ -spaces, then also the product*

$$(Z, \zeta) := \left(\prod_{i \in I} X_i, \bigotimes_{i \in I} \mu_i \right),$$

equipped with the diagonal Γ -action, is a standard Γ -space, and

$$|M|^Z \leq \inf_{i \in I} |M|^{X_i}.$$

- (2) *Let (X, μ) be a standard Γ -space and let (Y, ν) be some standard Borel probability space. Then*

$$|M|^Z = |M|^X,$$

where $Z := X \times Y$ is given the Γ -action induced by the Γ -action on X and where $\zeta := \mu \otimes \nu$ is the product measure on Z .

Proof. The first part follows by applying Proposition 4.12 for all $i \in I$ to the projection $\prod_{j \in I} X_j \rightarrow X_i$.

We now show the second part (following a similar argument by Schmidt [29, Proposition 5.29]). We can view (Y, ν) as standard Γ -space with trivial Γ -action. Then we obtain

$$|M|^Z \leq |M|^X$$

from the first part. For the converse inequality, we consider a Z -parametrised fundamental cycle $c = \sum_{j=0}^k f_j \otimes \sigma_j \in L^\infty(Z, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})$ in reduced form. So, if $c_Z \in \mathbb{Z} \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})$ is a fundamental cycle of M , there is a chain $d \in L^\infty(Z, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_{n+1}(\tilde{M}, \mathbb{Z})$ such that

$$c - c_Z = \partial d \in L^\infty(Z, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}).$$

Therefore, for ν -almost all $y \in Y$, the chain

$$c_y := \sum_{j=0}^k (x \mapsto f_j(x, y)) \otimes \sigma_j \in L^\infty(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})$$

is well-defined (Γ acts trivially on Y) and an X -parametrised fundamental cycle in reduced form (witnessed by the corresponding evaluation of d at y). By Fubini's theorem,

$$\begin{aligned} |c|^Z &= \int_{X \times Y} \sum_{j=0}^k |f_j| d(\mu \otimes \nu) \\ &= \int_Y \int_X \sum_{j=0}^k |f_j(x, y)| d\mu(x) d\nu(y) \\ &= \int_Y |c_y|^X d\nu(y). \end{aligned}$$

Hence, there is a $y \in Y$ such that c_y is an X -parametrised fundamental cycle and $|c_y|^X \leq |c|^Z$. Taking the infimum over all Z -parametrised fundamental cycles c shows $|M|^X \leq |M|^Z$, as desired. \square

Taking products of parameter spaces hence shows that the infimum in the definition of integral foliated simplicial volume is a *minimum*:

Corollary 4.14. *Let M be an oriented closed connected manifold with fundamental group Γ . Then there exists a standard Γ -space (X, μ) with essentially free Γ -action satisfying*

$$|M| = |M|^X.$$

Proof. Let (X_0, μ_0) be a standard Γ -space with essentially free Γ -action, e.g., the Bernoulli shift of Γ (Example 4.9) (or, in the case of finite Γ just Γ with the normalised counting measure). For $n \in \mathbb{N}_{>0}$ let (X_n, μ_n) be a standard Γ -space with

$$|M|^{X_n} \leq |M| + \frac{1}{n}.$$

Then the diagonal Γ -action on $(X, \mu) := (\prod_{n \in \mathbb{N}} X_n, \otimes_{n \in \mathbb{N}} \mu_n)$ is essentially free and we obtain $|M|^X = |M|$ from Proposition 4.13. \square

4.3. Convex combinations of parameter spaces

Proposition 4.15 (convex combinations of parameter spaces). *Let M be an oriented closed connected n -manifold with fundamental group Γ , let (X, μ) and (Y, ν) be standard Γ -spaces, and let $t \in [0, 1]$. Then*

$$|M|^Z = t \cdot |M|^X + (1-t) \cdot |M|^Y,$$

where $Z := X \sqcup Y$ is the disjoint union of X and Y endowed with the obvious Γ -action and the probability measure $\zeta := t \cdot \mu \sqcup (1-t) \cdot \nu$.

Proof. Under the mutually inverse $\mathbb{Z}\Gamma$ -isomorphisms

$$\begin{aligned} L^\infty(Z, \zeta, \mathbb{Z}) &\longleftrightarrow L^\infty(X, \mu, \mathbb{Z}) \oplus L^\infty(Y, \nu, \mathbb{Z}), \\ f &\longmapsto (f|_X, f|_Y), \\ \chi_X \cdot f + \chi_Y \cdot g &\longleftarrow (f, g), \end{aligned}$$

the constant function 1 on Z corresponds to $(1, 1)$, and for all $f \in L^\infty(Z, \zeta, \mathbb{Z})$ we have

$$\int_Z f d\zeta = t \cdot \int_X f|_X d\mu + (1-t) \cdot \int_Y f|_Y d\nu.$$

Therefore, the same arguments as in Proposition 4.12 show that under the induced mutually inverse chain isomorphisms

$$\begin{aligned} L^\infty(Z, \zeta, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}) &\longleftrightarrow L^\infty(X, \mu, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}) \\ &\oplus L^\infty(Y, \nu, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}) \end{aligned}$$

Z -parametrised fundamental cycles correspond to pairs of X -parametrised and Y -parametrised fundamental cycles and that (by applying the arguments in both directions)

$$|M|^Z \geq t \cdot |M|^X + (1-t) \cdot |M|^Y,$$

and

$$|M|^Z \leq t \cdot |M|^X + (1-t) \cdot |M|^Y. \quad \square$$

In combination with Example 4.5 we obtain:

Corollary 4.16. *Let M be an oriented closed connected manifold with fundamental group Γ . Then*

$$\{ |M|^X \mid (X, \mu) \text{ is a standard } \Gamma\text{-space} \} = [|M|, \|M\|_{\mathbb{Z}}] \subset \mathbb{R}.$$

4.4. Ergodic decomposition of parameter spaces. We will now show that ergodic parameter spaces suffice to describe the integral foliated simplicial volume:

Proposition 4.17 (ergodic parameters suffice). *Let M be an oriented closed connected manifold with fundamental group Γ .*

- (1) *If (X, μ) is a standard Γ -space and $\varepsilon \in \mathbb{R}_{>0}$, then there is a Γ -invariant ergodic probability measure μ' on the measurable Γ -space X with*

$$|M|^{(X, \mu')} \leq |M|^{(X, \mu)} + \varepsilon.$$

- (2) *In particular, for every $\varepsilon \in \mathbb{R}_{>0}$ there is an ergodic standard Γ -space X with*

$$|M|^X \leq |M| + \varepsilon.$$

The proof of this proposition relies on the ergodic decomposition theorem:

Theorem 4.18 (ergodic decomposition [6, Theorem 5][33, Theorem 4.2]). *Let Γ be a countable group and let (X, μ) be a standard Γ -space. Then there is a probability space (P, ν) and a family $(\mu_p)_{p \in P}$ of Γ -ergodic probability measures on the measurable Γ -space X with the following property: for each Borel subset $A \subset X$, the function*

$$\begin{aligned} P &\longrightarrow [0, 1], \\ p &\longmapsto \mu_p(A), \end{aligned}$$

is measurable and

$$\mu(A) = \int_P \mu_p(A) d\nu(p).$$

In view of this theorem all standard Γ -spaces can be seen as an assembly of ergodic Γ -spaces. However, we have to be careful about the sets of measure 0 with respect to the involved measures. Therefore, we consider the following “strict” function spaces and chain complexes:

Definition 4.19. Let X be a measurable space. We write $B(X, \mathbb{Z})$ for the set of bounded measurable functions of type $X \rightarrow \mathbb{Z}$. If μ is a measure on X , we write

$$N(X, \mu, \mathbb{Z}) := \{ f \in B(X, \mathbb{Z}) \mid \mu(f^{-1}(\mathbb{Z} \setminus 0)) = 0 \}$$

for the set of all functions vanishing μ -almost everywhere.

Remark 4.20. Let M be an oriented closed connected manifold with fundamental group Γ and let (X, μ) be a standard Γ -space. By definition,

$$L^\infty((X, \mu), \mathbb{Z}) \cong B(X, \mathbb{Z})/N(X, \mu, \mathbb{Z}),$$

and this isomorphism of $\mathbb{Z}\Gamma$ -modules gives rise to an isomorphism

$$L^\infty((X, \mu), \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}) \cong \frac{B(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z})}{N(X, \mu, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z})}$$

of chain complexes (because the chain modules of $C_*(\tilde{M}, \mathbb{Z})$ are free, and hence flat, over $\mathbb{Z}\Gamma$).

Proof of Proposition 4.17. It suffices to prove the first part. Let (X, μ) be a standard Γ -space, let $\varepsilon \in \mathbb{R}_{>0}$, and let $n := \dim M$. Then there is an (X, μ) -parametrised fundamental cycle

$$c = \sum_{j=0}^k f_j \otimes \sigma_j \in L^\infty((X, \mu), \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})$$

with

$$\sum_{j=0}^k \int_X |f_j| d\mu \leq |M|^{(X, \mu)} + \varepsilon.$$

Let $c_{\mathbb{Z}} \in \mathbb{Z} \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})$ be an integral fundamental cycle of M . Because c is an (X, μ) -parametrised fundamental cycle of M , we can find a chain $d \in L^\infty((X, \mu), \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_{n+1}(\tilde{M}, \mathbb{Z})$ satisfying

$$c - c_{\mathbb{Z}} = \partial d \in L^\infty((X, \mu), \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}).$$

In view of Remark 4.20, we can assume that the coefficients f_0, \dots, f_k of c and those of d lie in the “strict” function space $B(X, \mathbb{Z})$ and that there is a (without loss of generality, Γ -invariant) μ -null set $A \subset X$ and a chain $c' \in B(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})$ satisfying the relation

$$c - c_{\mathbb{Z}} = \partial d + \chi_A \cdot c' \in B(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}).$$

Here, $\chi_A \cdot c'$ uses the canonical $B(X, \mathbb{Z})^\Gamma$ - $\mathbb{Z}\Gamma$ -bimodule structure on $B(X, \mathbb{Z})$.

By the ergodic decomposition theorem (Theorem 4.18), we obtain a probability space (P, ν) and Γ -invariant ergodic probability measures $(\mu_p)_{p \in P}$ on X with

$$\mu(B) = \int_X \mu_p(B) \, d\nu(p)$$

for all Borel sets $B \subset X$. Hence, for all $f \in B(X, \mathbb{Z})$ we have

$$\int_X f \, d\mu = \int_P \int_X f \, d\mu_p \, d\nu(p).$$

Taking $f := \sum_{j=0}^k |f_j|$ and keeping in mind that A is a μ -null set, we thus find a $p \in P$ with

$$\mu_p(A) = 0 \quad \text{and} \quad \int_X f \, d\mu_p \leq \int_X f \, d\mu.$$

We now show that

$$|M|^{(X, \mu_p)} \leq \int_X f \, d\mu \leq |M|^{(X, \mu)} + \varepsilon.$$

To this end, we consider the chain

$$c_p := [c] \in \frac{B(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})}{N(X, \mu_p, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z})} \cong L^\infty((X, \mu_p), \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}).$$

Then c_p is an (X, μ_p) -parametrised fundamental cycle of M , because

$$c - c_{\mathbb{Z}} = \partial d + \chi_A \cdot c'$$

holds in the “strict” twisted chain complex $B(X, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z})$ and $\mu_p(A) = 0$, and so

$$c_p - c_{\mathbb{Z}} = \partial[d] \in L^\infty((X, \mu_p), \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}).$$

Furthermore, we obtain the desired estimate for the norms, namely

$$|c_p|^{(X, \mu_p)} \leq \sum_{j=0}^k \int_X |f_j| \, d\mu_p = \int_X f \, d\mu_p \leq \int_X f \, d\mu \leq |M|^{(X, \mu)} + \varepsilon. \quad \square$$

As products of ergodic spaces are not necessarily ergodic, it is not clear that there is an analogue of Corollary 4.14 for ergodic parameter spaces:

Question 4.21. *Is the integral foliated simplicial volume always given by an ergodic parameter space? Is the integral foliated simplicial volume always given by the Bernoulli shift of the fundamental group?*

4.5. Integral foliated simplicial volume and finite coverings. We will now prove that integral foliated simplicial volume is multiplicative with respect to finite coverings:

Theorem 4.22 (multiplicativity of integral foliated simplicial volume). *Let M be an oriented closed connected n -manifold and let $p: N \rightarrow M$ be a d -sheeted covering with $d \in \mathbb{N}_{>0}$. Then*

$$|M| = \frac{1}{d} \cdot |N|.$$

The theorem will follow from compatibility with respect to restriction and induction of parameter spaces (Proposition 4.29 and 4.26).

Setup 4.23. Let M be an oriented closed connected n -manifold with fundamental group Γ , and let $p: N \rightarrow M$ be a d -sheeted covering with $d \in \mathbb{N}_{>0}$. Let Λ be the fundamental group of N , and let $\Lambda' = \pi_1(p)(\Lambda) \cong \Lambda$ be the subgroup of Γ associated with p (which has index d in Γ). For notational simplicity, in the following, we will identify the groups Λ and Λ' via the isomorphism given by p .

For the discussion of induction spaces and associated constructions, it will be necessary to choose representatives:

Setup 4.24. Let Γ be a countable group, let $\Lambda \subset \Gamma$ be a subgroup of finite index $d := [\Gamma : \Lambda]$, and let $g_1, \dots, g_d \in \Gamma$ be a set of representatives of Λ in Γ :

$$\{g_1 \cdot \Lambda, \dots, g_d \cdot \Lambda\} = \Gamma/\Lambda.$$

Definition 4.25 (induction). In the situation of Setup 4.24, let (Y, ν) be a standard Λ -space. Then the *induction* $(\Gamma \times_{\Lambda} Y, \mu)$ of (Y, ν) from Λ to Γ is the standard Γ -space defined as follows.

– The set

$$\Gamma \times_{\Lambda} Y := \Gamma \times Y / \{(g \cdot h, y) \sim (g, h \cdot y) \mid g \in \Gamma, h \in \Lambda, y \in Y\}$$

is endowed with the measurable structure induced from the bijection

$$\begin{aligned} \Gamma \times_{\Lambda} Y &\longrightarrow \Gamma/\Lambda \times Y, \\ [g_j \cdot y] &\longmapsto (g_j \cdot \Lambda, y) \end{aligned}$$

(where Γ/Λ is given the discrete Borel structure). Moreover, the probability measure μ is the pull-back of the measure $1/d \cdot \nu' \otimes \nu$ on $\Gamma/\Lambda \times Y$ under this bijection, where ν' denotes the counting measure on Γ/Λ .

– The Γ -action on $\Gamma \times_{\Lambda} Y$ is defined by

$$\begin{aligned} \Gamma \times (\Gamma \times_{\Lambda} Y) &\longrightarrow \Gamma \times_{\Lambda} Y, \\ (g, [g', y]) &\longmapsto [g \cdot g', y]. \end{aligned}$$

Notice that in the above definition μ is indeed Γ -invariant, and that the measurable structure and the probability measure μ on the induction space do *not* depend on the chosen set of representatives.

Proposition 4.26 (induction of parameter spaces). *In the situation of Setup 4.23, let (Y, ν) be a standard Λ -space. Then*

$$|M|^{\Gamma \times_{\Lambda} Y} = \frac{1}{d} \cdot |N|^Y.$$

Proof. We choose representatives g_1, \dots, g_d for the index d subgroup $\Lambda \subset \Gamma$ as in Setup 4.24. Induction of parameter spaces is compatible with algebraic induction of modules: we have (well-defined) mutually inverse $\mathbb{Z}\Gamma$ -isomorphisms

$$\begin{aligned} \varphi: L^{\infty}(\Gamma \times_{\Lambda} Y, \mathbb{Z}) &\longrightarrow L^{\infty}(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} \mathbb{Z}\Gamma, \\ f &\longmapsto \sum_{j=1}^d (y \mapsto f([g_j, y])) \otimes g_j, \\ \psi: L^{\infty}(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} \mathbb{Z}\Gamma &\longrightarrow L^{\infty}(\Gamma \times_{\Lambda} Y, \mathbb{Z}) \\ f \otimes g_j &\longmapsto \left([g_k, y] \mapsto \begin{cases} f(y) & \text{if } k = j \\ 0 & \text{if } k \neq j \end{cases} \right). \end{aligned}$$

Because $p: M \rightarrow N$ is a finite covering, M and N share the same universal covering space $\tilde{M} = \tilde{N}$ and the Λ -action on \tilde{N} is nothing but the restriction of the Γ -action on \tilde{M} . Therefore, the above maps induce mutually inverse chain complex isomorphisms

$$\begin{aligned} \Phi: L^{\infty}(\Gamma \times_{\Lambda} Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}) &\longrightarrow L^{\infty}(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} \mathbb{Z}\Gamma \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}) \\ &\cong L^{\infty}(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} C_*(\tilde{N}, \mathbb{Z}), \\ f \otimes c &\longmapsto \sum_{j=1}^d f([g_j, \cdot]) \otimes g_j \cdot c, \\ \Psi: L^{\infty}(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} C_*(\tilde{N}, \mathbb{Z}) &\longrightarrow L^{\infty}(\Gamma \times_{\Lambda} Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z}), \\ f \otimes c &\longmapsto \psi(f \otimes 1) \otimes c. \end{aligned}$$

It is not difficult to see that Φ and Ψ map $\Gamma \times_{\Lambda} Y$ -parametrised fundamental cycles of M to Y -parametrised fundamental cycles of N , and vice versa: it suffices to prove this claim for Φ . For this, we use the following transfer type argument. Let $c_{\mathbb{Z}} = \sum_{j=1}^k a_j \otimes \sigma_j \in \mathbb{Z} \otimes_{\mathbb{Z}\Gamma} C_n(\tilde{M}, \mathbb{Z}) \cong C_n(M, \mathbb{Z})$ be a fundamental cycle of M . By construction,

$$\Phi \circ i_M^{\Gamma \times_{\Lambda} Y}(c_{\mathbb{Z}}) = i_N^Y \left(\sum_{j=1}^k a_j \otimes \sum_{\ell=1}^d g_{\ell} \cdot \sigma_j \right),$$

and $g_1 \cdot \sigma_j, \dots, g_d \cdot \sigma_j$ are π_N -lifts of the d different p -lifts of $\pi_M \circ \sigma_j$, where $\pi_N: \tilde{N} \rightarrow N$ and $\pi_M: \tilde{M} \rightarrow M$ denote the universal covering maps. Therefore, $\sum_{j=1}^k a_j \otimes \sum_{\ell=1}^d g_{\ell} \cdot \sigma_j$ is a fundamental cycle of N , which proves the claim about parametrised fundamental cycles.

By definition of the induction space (Definition 4.25), the d copies of Y inside $\Gamma \times_{\Lambda} Y$ are each given the weight $1/d$. Therefore, it is not difficult to show that

$$|\Phi(c)|^Y \leq d \cdot |c|^{\Gamma \times_{\Lambda} Y}$$

holds for all chains $c \in L^{\infty}(\Gamma \times_{\Lambda} Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Gamma} C_*(\tilde{M}, \mathbb{Z})$, and that

$$|\Psi(c)|^{\Gamma \times_{\Lambda} Y} \leq \frac{1}{d} \cdot |c|^Y$$

holds for all chains $c \in L^{\infty}(Y, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} C_*(\tilde{N}, \mathbb{Z})$. Taking the infimum over all parametrised fundamental cycles therefore yields that

$$|M|^{\Gamma \times_{\Lambda} Y} \leq \frac{1}{d} \cdot |N|^Y \quad \text{and} \quad |N|^Y \leq d \cdot |M|^{\Gamma \times_{\Lambda} Y}. \quad \square$$

Corollary 4.27 (coset spaces as parameter space). *In the situation of Setup 4.23 we have*

$$|M|^{\Gamma/\Lambda} = \frac{1}{d} \cdot \|N\|_{\mathbb{Z}}.$$

Here, we equip the finite set Γ/Λ with the left Γ -action given by translation and the normalised counting measure.

Proof. Let X be a standard Λ -space consisting of a single point. Then

$$\Gamma/\Lambda \cong \Gamma \times_{\Lambda} X$$

(in the category of standard Γ -spaces). Hence, Proposition 4.26 and Example 4.5 show that

$$|M|^{\Gamma/\Lambda} = \frac{1}{d} \cdot |N|^X = \frac{1}{d} \cdot \|N\|_{\mathbb{Z}}. \quad \square$$

Conversely, we will now consider restriction of parameter spaces:

Definition 4.28 (restriction). Let Γ be a group, let (X, μ) be a standard Γ -space, and let $\Lambda \subset \Gamma$ be a subgroup. Restricting the Γ -action on X to Λ (and keeping the same probability measure) results in a standard Λ -space, the *restriction* $\text{res}_\Lambda^\Gamma(X, \mu)$ of (X, μ) from Γ to Λ .

Proposition 4.29 (restriction of parameter spaces). *In the situation of Setup 4.23 let (X, μ) be a standard Γ -space. Then*

$$\frac{1}{d} \cdot |N|^{\text{res}_\Lambda^\Gamma X} \leq |M|^X.$$

Proof. In view of Proposition 4.26, it suffices to show that

$$|M|^{\Gamma \times_\Lambda \text{res}_\Lambda^\Gamma X} \leq |M|^X.$$

The map

$$\begin{aligned} \Gamma \times_\Lambda \text{res}_\Lambda^\Gamma X &\longrightarrow X, \\ [g, x] &\longmapsto g \cdot x, \end{aligned}$$

satisfies the hypotheses of Proposition 4.12. Therefore, we obtain the desired estimate $|M|^{\Gamma \times_\Lambda \text{res}_\Lambda^\Gamma X} \leq |M|^X$. \square

Example 4.30. In the situation of Proposition 4.29, in general, equality will *not* hold. For example, we could consider a double covering $S^1 \rightarrow S^1$ and a parameter space for the base manifold consisting of a single point.

We will now complete the proof of Theorem 4.22:

Proof of Theorem 4.22. From Proposition 4.29 we obtain, by taking the infimum over all standard Γ -spaces as parameter spaces for M ,

$$\frac{1}{d} \cdot |N| \leq |M|.$$

Conversely, from Proposition 4.26 we obtain, by taking the infimum over all standard Λ -spaces as parameter spaces for N ,

$$|M| \leq \frac{1}{d} \cdot |N|. \quad \square$$

5. A proportionality principle for integral foliated simplicial volume

In this section, we will provide a proof of the proportionality principle Theorem 1.2 and of Corollary 1.3. We will use the language of measure equivalence of groups and techniques of Bader, Furman, and Sauer [1, Theorem 1.9].

5.1. Measure equivalence. The notion of measure equivalence was originally introduced by Gromov [14, 0.5.E1] as a measure-theoretic analogue of quasi-isometry.

Definition 5.1 (measure equivalence). Two countable groups Γ and Λ are called *measure equivalent* (ME) if there is a standard measure space (Ω, μ) with commuting measure preserving Γ - and Λ -actions, such that each of the actions admits a finite measure fundamental domain X_Γ and X_Λ respectively. The space (Ω, μ) endowed with these actions is called an *ME-coupling* of Γ and Λ . The ratio $c_\Omega = \mu(X_\Lambda)/\mu(X_\Gamma)$ is independent of the chosen fundamental domains and is called the *coupling index* of the ME-coupling Ω .

In our context, the fundamental domains do not need to be strict fundamental domains; it suffices that $\Omega = \bigcup_{\gamma \in \Gamma} \gamma \cdot X_\Gamma$ is a disjoint decomposition up to measure 0. The cocycle in Definition 5.6 then will only be well-defined up to sets of measure 0; however, this poses no problems in the sequel as we will pass to L^1 - and L^∞ -spaces anyway.

Example 5.2 (lattices). A second countable locally compact group G with its Haar measure is an ME-coupling for every pair of lattices Γ and Λ in G . Indeed, the Haar measure on G is bi-invariant (because G contains lattices), and the left actions

$$\begin{aligned} \Gamma \times G &\longrightarrow G, & \Lambda \times G &\longrightarrow G, \\ (\gamma, g) &\longmapsto \gamma g, & (\lambda, g) &\longmapsto g\lambda^{-1}, \end{aligned}$$

of Γ and Λ on G given by multiplication in G commute with each other.

Setup 5.3. Let (Ω, μ) be an ME-coupling of Γ and Λ . We suppose that both the actions are left actions, and we fix a fundamental domain for each of the two actions on the ME-coupling, denoted as X_Γ and X_Λ respectively.

Definition 5.4 (ergodic/mixing ME-coupling). In the situation of Setup 5.3, the ME-coupling is *ergodic* (resp. *mixing*) if the Γ -action on $\Lambda \backslash \Omega$ is ergodic (resp. mixing) and the Λ -action on $\Gamma \backslash \Omega$ is ergodic (resp. mixing).

Remark 5.5. Note that in this situation the Γ -action on $\Lambda \backslash \Omega$ is ergodic if and only if the Λ -action on $\Gamma \backslash \Omega$ is ergodic [12, Lemma 2.2].

Definition 5.6 (ME-cocycle). In the situation of the Setup 5.3, we define the measurable cocycle α_Λ associated to X_Λ as the map

$$\alpha_\Lambda: \Gamma \times X_\Lambda \longrightarrow \Lambda$$

such that $\alpha_\Lambda(\gamma, x)$ is the unique element satisfying $\gamma x \in \alpha_\Lambda(\gamma, x)^{-1} X_\Lambda$ for all $x \in X_\Lambda$ and $\gamma \in \Gamma$. Similarly, we define $\alpha_\Gamma: \Lambda \times X_\Gamma \rightarrow \Gamma$. If we choose another fundamental domain for $\Lambda \backslash \Omega$ then the associated cocycle is measurably cohomologous to α_Λ [12, Section 2] (the same for α_Γ).

With this notation the natural left action of Γ on X_Λ and of Λ on X_Γ is described as follows:

$$\begin{aligned} \Gamma \times X_\Lambda &\longrightarrow X_\Lambda, & \Lambda \times X_\Gamma &\longrightarrow X_\Gamma, \\ (\gamma, x) &\longmapsto \gamma \bullet x := \alpha_\Lambda(\gamma, x)\gamma x, & (\lambda, y) &\longmapsto \lambda \bullet y := \alpha_\Gamma(\lambda, y)\lambda y, \end{aligned} \tag{3}$$

where we write $\gamma \bullet x$ to distinguish it from the action γx of Γ on Ω .

Remark 5.7. In the situation of the Setup 5.3, consider the standard Γ -space $(X_\Lambda, \mu_\Lambda = \mu(X_\Lambda)^{-1}\mu|_{X_\Lambda})$ with Γ -action described above and the standard Γ -space $\Lambda \backslash \Omega$ with the probability measure induced from μ and the left translation Γ -action. Then the map $X_\Lambda \hookrightarrow \Omega \rightarrow \Lambda \backslash \Omega$ is a measure isomorphism.

Definition 5.8 (bounded ME-coupling). In the situation of the Setup 5.3, assume that Λ is finitely generated, and let $l: \Lambda \rightarrow \mathbb{N}$ be the length function associated to some word-metric on Λ . We say that the fundamental domain X_Λ is *bounded* if, for every $\gamma \in \Gamma$, the function $x \mapsto l(\alpha_\Lambda(\gamma, x))$ is in $L^\infty(X_\Lambda, \mathbb{R})$.

Let Γ and Λ be finitely generated. An ME-coupling of Γ and Λ is *bounded* if it admits bounded Γ - and Λ -fundamental domains.

Example 5.9. A connected second countable locally compact group G with its Haar measure is a bounded ME-coupling for every pair of uniform lattices in G [2, p. 321], [31, Corollary 6.12, p. 58].

5.2. Homology of groups. In the aspherical case, we can express integral foliated simplicial volume in terms of group homology. Let Γ be a discrete group. The *bar resolution* of Γ is the $\mathbb{Z}\Gamma$ -chain complex $C_*(\Gamma)$ defined as follows: for each $n \in \mathbb{N}$ let

$$C_n(\Gamma) = \left\{ \sum_{\gamma \in \Gamma^{n+1}} a_\gamma \cdot \gamma_0 \cdot [\gamma_1 | \cdots | \gamma_n] \mid \text{for all } \gamma = (\gamma_0, \dots, \gamma_n) \in \Gamma^{n+1}, a_\gamma \in \mathbb{Z} \right\}$$

with the Γ -action characterized by

$$\bar{\gamma} \cdot (\gamma_0 \cdot [\gamma_1 | \cdots | \gamma_n]) = (\bar{\gamma} \cdot \gamma_0) \cdot [\gamma_1 | \cdots | \gamma_n]$$

for all $\bar{\gamma} \in \Gamma$ and all $\gamma \in \Gamma^{n+1}$. The differential

$$\partial_*: C_*(\Gamma) \longrightarrow C_{*-1}(\Gamma)$$

is defined by

$$\begin{aligned} C_n(\Gamma) &\longrightarrow C_{n-1}(\Gamma), \\ \gamma_0 \cdot [\gamma_1 | \cdots | \gamma_n] &\longmapsto \gamma_0 \cdot \gamma_1 \cdot [\gamma_2 | \cdots | \gamma_n] \\ &\quad + \sum_{j=1}^{n-1} (-1)^j \cdot \gamma_0 \cdot [\gamma_1 | \cdots | \gamma_{j-1} | \gamma_j \cdot \gamma_{j+1} | \gamma_{j+2} | \cdots | \gamma_n] \\ &\quad + (-1)^n \cdot \gamma_0 \cdot [\gamma_1 | \cdots | \gamma_{n-1}]. \end{aligned}$$

Moreover, the bar resolution is a normed chain complex (i.e., the differentials in each degree are bounded operators) with the ℓ^1 -norm given by

$$\left\| \sum_{\gamma \in \Gamma^{n+1}} a_\gamma \cdot \gamma_0 \cdot [\gamma_1 | \cdots | \gamma_n] \right\|_1 = \sum_{\gamma \in \Gamma^{n+1}} |a_\gamma|.$$

We obtain a version of the bar resolution with coefficients using the tensor product. For every normed right $\mathbb{Z}\Gamma$ -module A let

$$C_*(\Gamma, A) := A \otimes_{\mathbb{Z}\Gamma} C_*(\Gamma).$$

Definition 5.10 (group homology). Let Γ be a discrete group and A be a normed right $\mathbb{Z}\Gamma$ -module. Then the *group homology of Γ with coefficients in A* is

$$H_*(\Gamma, A) := H_*(C_*(\Gamma, A)).$$

The ℓ^1 -norm on the chain complex induces an ℓ^1 -semi-norm on the group homology.

Proposition 5.11. *Let M be an aspherical manifold with universal covering \tilde{M} , and fundamental group Γ . Let A be a normed right $\mathbb{Z}\Gamma$ -module. Then there exists a natural chain map*

$$c_\Gamma: C_*(\tilde{M}, A) \longrightarrow C_*(\Gamma, A)$$

that induces an isometric isomorphism

$$c_\Gamma: H_*(M, A) \longrightarrow H_*(\Gamma, A).$$

Proof. It is not difficult to see that the classical mutually inverse $\mathbb{Z}\Gamma$ -chain homotopy equivalences $C_*(\tilde{M}, \mathbb{Z}) \longleftrightarrow C_*(\Gamma, \mathbb{Z})$, defined using a Γ -fundamental domain on \tilde{M} , are norm non-increasing. This gives the desired isometric isomorphism in homology. □

Corollary 5.12. *Let M , Γ , and (X, μ) be as in Definition 4.1. If M is aspherical, there exists an isometric isomorphism*

$$H_*(M, L^\infty(X, \mathbb{Z})) \xrightarrow{\cong} H_*(\Gamma, L^\infty(X, \mathbb{Z})),$$

with respect to the ℓ^1 -semi-norm induced by the ℓ^1 -norm on $L^\infty(X, \mathbb{Z})$.

Lemma 5.13. *Let Γ be a countable group and let (X, μ) be a standard Γ -space as in Definition 4.1. The inclusion $L^\infty(X, \mathbb{Z}) \hookrightarrow L^1(X, \mathbb{Z})$ has dense image with respect to the ℓ^1 -norm and induces an isometric map*

$$H_*(\Gamma, L^\infty(X, \mathbb{Z})) \longrightarrow H_*(\Gamma, L^1(X, \mathbb{Z})).$$

Proof. Inclusions of dense subcomplexes induce isometric maps on homology [29, Lemma 2.9], [18, Proposition 1.7] (the cited proofs also carry over to this integral setting). □

Remark 5.14. Let M , Γ , and (X, μ) be as in Definition 4.1. If M is aspherical, by Corollary 5.12 and Lemma 5.13 we deduce that the X -parametrised simplicial volume of M can be computed via the ℓ^1 -semi-norm on the group homology $H_*(\Gamma, L^1(X, \mathbb{Z}))$.

5.3. Proportionality principle, general case

Proof of Theorem 1.2. Let $n \in \mathbb{N}$. Let Γ and Λ be fundamental groups of oriented closed connected aspherical n -manifolds M and N with positive simplicial volume. Assume that (Ω, μ) is an ergodic bounded ME-coupling of Γ and Λ . Fix a bounded fundamental domain $X_\Gamma \subset \Omega$ (resp. $X_\Lambda \subset \Omega$) of the Γ -action on Ω (resp. of the Λ -action on Ω) and let α_Γ (resp. α_Λ) be the associated cocycle (Definition 5.6).

Using the cocycles associated with the ME-coupling we translate parametrised fundamental cycles of one manifold into parametrised fundamental cycles of the other manifold, while controlling the ℓ^1 -semi-norm. More precisely, consider the standard space $(X_\Gamma, \mu_\Gamma = \mu(X_\Gamma)^{-1} \mu|_{X_\Gamma})$ with the Λ -action defined in equation (3). By Remark 5.14 we use the ℓ^1 -norm on $L^1(X_\Gamma, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} C_*(\Lambda, \mathbb{Z})$ to estimate the X_Γ -parametrised simplicial volume of N . Let $\Lambda^{*+1} \times X_\Gamma$ be endowed with the diagonal Λ -action and which carries the product of the counting measure and μ_Γ . We identify $L^1(X_\Gamma, \mathbb{Z}) \otimes_{\mathbb{Z}\Lambda} C_*(\Lambda, \mathbb{Z})$ with $L^1(\Lambda^{*+1} \times X_\Gamma, \mathbb{Z})_\Lambda^{\text{fin}}$, where the superscript “fin” indicates the submodule of functions $f: \Lambda^{*+1} \times X_\Gamma \rightarrow \mathbb{Z}$ with the property that there is a finite subset F of Λ^{*+1} such that f is supported on $F \times X_\Gamma$, and where the subscript Λ indicates the Λ -co-invariants.

Now, let us consider the measurable, countable-to-one, locally measure preserving (up to a constant factor) map [1, p. 284]

$$\begin{aligned} \varphi_n^{(\alpha_\Gamma)}: \Lambda^{n+1} \times X_\Gamma &\longrightarrow \Gamma^{n+1} \times X_\Lambda, \\ (\lambda_0, \dots, \lambda_n, y) &\longmapsto (\alpha_\Gamma(\lambda_0^{-1}, y)^{-1}, \dots, \alpha_\Gamma(\lambda_n^{-1}, y)^{-1}, \Lambda y \cap X_\Lambda), \end{aligned}$$

and define

$$\begin{aligned}
 (\alpha_\Gamma)_n^{\mathbb{Z}}: L^1(\Lambda^{n+1} \times X_\Gamma, \mathbb{Z})_\Lambda^{\text{fin}} &\longrightarrow L^1(\Gamma^{n+1} \times X_\Lambda, \mathbb{Z})_\Gamma^{\text{fin}}, \\
 f &\longmapsto \left((\gamma, x) \longmapsto (\alpha_\Gamma)_n^{\mathbb{Z}}(f)(\gamma, x) \right) \\
 &:= \sum_{(\lambda, y) \in (\varrho_n^{(\alpha_\Gamma)})^{-1}(\gamma, x)} f(\lambda, y).
 \end{aligned}$$

This map is the restriction to integral chains of the map described by Bader, Furman and Sauer [1, Theorem 5.7]¹ up to rescaling by the coupling index $c_\Omega = \mu(X_\Lambda)/\mu(X_\Gamma)$. The sum on the right hand side is a.e. finite [1, Lemma 5.8], and the boundedness of the coupling guarantees that $(\alpha_\Gamma)_n^{\mathbb{Z}}(f)$ has finite support whenever f has finite support; moreover, a straightforward computation shows that the map is well-defined on the level of co-invariants. Hence, for a bounded ME-coupling of Γ and Λ , the map $(\alpha_\Gamma)_n^{\mathbb{Z}}$ is a well-defined norm non-increasing (up to rescaling by the coupling index) chain map, and $(\alpha_\Gamma)_n^{\mathbb{Z}}$ induces a map

$$H((\alpha_\Gamma)_n^{\mathbb{Z}}): H_n(\Lambda, L^1(X_\Gamma, \mathbb{Z})) \longrightarrow H_n(\Gamma, L^1(X_\Lambda, \mathbb{Z}))$$

of norm at most $1/c_\Omega$.

Let us now consider the diagram in Figure 1. Since M and N are aspheri-

$$\begin{array}{ccc}
 H_n^{\ell^1}(\Lambda, L^1(X_\Gamma, \mathbb{R})) & \xrightarrow{H^{\ell^1}((\alpha_\Gamma)_n^{\mathbb{R}})} & H_n^{\ell^1}(\Gamma, L^1(X_\Lambda, \mathbb{R})) \\
 i_\Lambda \uparrow & & i_\Gamma \uparrow \\
 H_n(\Lambda, L^1(X_\Gamma, \mathbb{R})) & \xrightarrow{H((\alpha_\Gamma)_n^{\mathbb{R}})} & H_n(\Gamma, L^1(X_\Lambda, \mathbb{R})) \\
 j_\Lambda^{\mathbb{R}} \uparrow & & j_\Gamma^{\mathbb{R}} \uparrow \\
 H_n(\Lambda, L^1(X_\Gamma, \mathbb{Z})) & \xrightarrow{H((\alpha_\Gamma)_n^{\mathbb{Z}})} & H_n(\Gamma, L^1(X_\Lambda, \mathbb{Z})) \\
 j_\Lambda \uparrow & & j_\Gamma \uparrow \\
 H_n(\Lambda, \mathbb{Z}) & & H_n(\Gamma, \mathbb{Z}) \\
 c_\Lambda \uparrow & & c_\Gamma \uparrow \\
 H_n(N, \mathbb{Z}) & & H_n(M, \mathbb{Z})
 \end{array}$$

Figure 1. Effect of the cocycle in homology

cal manifolds, Proposition 5.11 ensures that there exist isometric isomorphisms c_Λ and c_Γ . The maps $j_\Lambda, j_\Gamma, j_\Lambda^{\mathbb{R}}, j_\Gamma^{\mathbb{R}}$ are the usual change of coefficients homomorphisms. In particular, by ergodicity of both actions, the maps j_Λ and j_Γ are

¹ In Definition 4.1 we define a right action on $L^\infty(X, \mathbb{Z})$ (and similarly on $L^1(X, \mathbb{Z})$) unlike *op. cit.*, in which $L^\infty(X, \mathbb{Z})$ is endowed with a left action.

(semi)-norm non-increasing isomorphisms. The inclusion of singular chains into ℓ^1 -chains induces the isometric homomorphisms i_Λ and i_Γ . Finally $c_\Omega \cdot H^{\ell^1}((\alpha_\Gamma)_n^{\mathbb{R}})$ is the isometric isomorphism of Bader, Furman and Sauer [1, Theorem 5.7] defined between ℓ^1 -homology groups.

By the diagram in Figure 1 we have that

$$H((\alpha_\Gamma)_n^{\mathbb{R}}) \circ j_\Lambda^{\mathbb{R}} \circ j_\Lambda \circ c_\Lambda([N]_{\mathbb{Z}}) = m \cdot (j_\Gamma^{\mathbb{R}} \circ j_\Gamma \circ c_\Gamma)([M]_{\mathbb{Z}}) \tag{4}$$

for some $m \in \mathbb{Z}$; here, we use that the Γ -action on X_Λ is ergodic, and so $H_n(\Gamma, L^1(X_\Lambda, \mathbb{Z})) \cong \mathbb{Z}$. By $\|N\| > 0$ we have $\|i_\Lambda \circ j_\Lambda^{\mathbb{R}} \circ j_\Lambda \circ c_\Lambda([N]_{\mathbb{Z}})\|_1 > 0$. Since $c_\Omega \cdot H^{\ell^1}((\alpha_\Gamma)_n^{\mathbb{R}})$ is an isometry we deduce that

$$H^{\ell^1}((\alpha_\Gamma)_n^{\mathbb{R}}) \circ i_\Lambda \circ j_\Lambda^{\mathbb{R}} \circ j_\Lambda \circ c_\Lambda([N]_{\mathbb{Z}}) \neq 0.$$

By the commutativity of the top square in Figure 1, we have $|m| \geq 1$ in equation (4), and hence

$$\begin{aligned} \|(j_\Gamma^{\mathbb{R}} \circ j_\Gamma \circ c_\Gamma)([M]_{\mathbb{Z}})\|_1 &\leq |m| \cdot \|(j_\Lambda^{\mathbb{R}} \circ j_\Lambda \circ c_\Lambda)([N]_{\mathbb{Z}})\|_1 \\ &= \|H((\alpha_\Gamma)_n^{\mathbb{R}}) \circ j_\Lambda^{\mathbb{R}} \circ j_\Lambda \circ c_\Lambda([N]_{\mathbb{Z}})\|_1 \\ &\leq c_\Omega^{-1} \cdot \|j_\Lambda^{\mathbb{R}} \circ j_\Lambda \circ c_\Lambda([N]_{\mathbb{Z}})\|_1. \end{aligned}$$

By interchanging the roles of Γ and Λ , we obtain the converse inequality and it turns out that $|m| = 1$.

By the commutativity of the diagram in Figure 1 it follows

$$|M|^{X_\Lambda} = \|j_\Gamma \circ c_\Gamma([M]_{\mathbb{Z}})\|_1 = \|H((\alpha_\Gamma)_n^{\mathbb{Z}}) \circ j_\Lambda \circ c_\Lambda([N]_{\mathbb{Z}})\|_1 \leq c_\Omega^{-1} \cdot |N|^{X_\Gamma},$$

which implies

$$|N|^{X_\Gamma} \geq c_\Omega \cdot |M|^{X_\Lambda}.$$

Similarly we have the other inequality.

Since the map of Remark 5.7 induces an isometric isomorphism between $H_n(\Gamma, L^1(X_\Lambda, \mathbb{Z}))$ and $H_n(\Gamma, L^1(\Lambda \setminus \Omega, \mathbb{Z}))$ (similarly for X_Γ), the previous equality does not depend on the choice of the fundamental domains:

$$|N|^{\Gamma \setminus \Omega} = c_\Omega \cdot |M|^{\Lambda \setminus \Omega}.$$

Let us now suppose that (Ω, μ) is a *mixing* bounded ME-coupling of Γ and Λ . Let (X, μ_X) be an ergodic standard Γ -space, with the structure of Λ -space with the trivial action. Then $(X \times \Omega, \mu_X \otimes \mu)$ is an ergodic bounded ME-coupling of Γ and Λ with respect to the left diagonal Γ - and Λ -actions. Indeed, if F_Γ (resp. F_Λ) is a bounded fundamental domain for the action of Γ on Ω (resp. of Λ) then $X_\Gamma = X \times F_\Gamma$ (resp. $X_\Lambda = X \times F_\Lambda$) is a finite measure fundamental domain of the Γ -action on $X \times \Omega$ (resp. of the Λ -action). We can easily trace back the boundedness of X_Γ (resp. X_Λ) to the one of F_Γ (resp. F_Λ). Finally,

by the property of mixing actions the Γ -action on $\Lambda \backslash (X \times \Omega) \cong X \times \Lambda \backslash \Omega$ is ergodic and $(X \times \Omega, \mu_X \otimes \mu)$ is an ergodic ME-coupling (see Definition 4.8 and Remark 5.5).

Applying now the previous construction to $(X \times \Omega, \mu_X \otimes \mu)$, we have

$$|N|^{\Gamma \backslash (X \times \Omega)} = c_{X \times \Omega} \cdot |M|^{\Lambda \backslash (X \times \Omega)},$$

where $c_{X \times \Omega} = (\mu_X \otimes \mu)(X \times X_\Lambda) / (\mu_X \otimes \mu)(X \times X_\Gamma) = \mu(X_\Lambda) / \mu(X_\Gamma) = c_\Omega$. Using Proposition 4.13 it follows that

$$|N| \leq |N|^{\Gamma \backslash (X \times \Omega)} = c_\Omega \cdot |M|^{X \times (\Lambda \backslash \Omega)} \leq c_\Omega \cdot |M|^X,$$

and by Proposition 4.17 it turns out that

$$|N| \leq c_\Omega \cdot |M|.$$

By interchanging the roles of Γ and Λ we obtain the other inequality. □

5.4. Proportionality principle, locally symmetric spaces of non-compact type.

For the proof of Theorem 1.1 we need a proportionality principle for hyperbolic manifolds. We prove it in the more general setting of locally symmetric spaces of non-compact type (Corollary 1.3).

Proof of Corollary 1.3. Let X be an aspherical symmetric space of non-compact type and let Γ and Λ be uniform lattices in $G := \text{Isom}^0(X)$. The group G with its Haar measure μ is a bounded ME-coupling with respect to Γ and Λ (see Examples 5.2 and 5.9). The ergodicity of the coupling follows from the Moore Ergodicity Theorem [3, Theorem III.2.1]. Clearly, this coupling has coupling index $\text{covol}(\Lambda) / \text{covol}(\Gamma)$. Finally, X/Λ and X/Γ have positive simplicial volume [17].

Notice that the standard Γ -space $\Lambda \backslash G$ (where Λ acts on G as in Example 5.2) is isomorphic to the coset space G/Λ with the probability measure induced from the Haar measure μ and the left translation Γ -action; similarly, the standard Λ -space $\Gamma \backslash G$ with the Λ -action induced from Example 5.2 is isomorphic to the coset space G/Γ with the left translation Λ -action.

Hence, Theorem 1.2 applied to this ergodic ME-coupling implies

$$|X/\Lambda|^{G/\Gamma} = \frac{\text{covol}(\Lambda)}{\text{covol}(\Gamma)} \cdot |X/\Gamma|^{G/\Lambda}.$$

Since the Moore Ergodicity Theorem [3, Theorem III.2.1] ensures also that (G, μ) is a mixing ME-coupling of Γ and Λ , by the second part of Theorem 1.2 it turns out that

$$|X/\Lambda| = \frac{\text{covol}(\Lambda)}{\text{covol}(\Gamma)} \cdot |X/\Gamma|. \quad \square$$

6. Comparing integral foliated simplicial volume and stable integral simplicial volume

We will now compare integral foliated simplicial volume with stable integral simplicial volume. We start with a general sandwich estimate (Proposition 6.1), and we will then give more refined estimates for the hyperbolic case and for special parameter spaces. In particular, we will prove Corollary 1.4 and Theorem 1.5.

6.1. Sandwich estimate for integral foliated simplicial volume. Integral foliated simplicial volume interpolates between ordinary simplicial volume and stable integral simplicial volume:

Proposition 6.1 (comparing simplicial volumes). *Let M be an oriented closed connected manifold. Then*

$$\|M\| \leq |M| \leq \|M\|_{\mathbb{Z}}^{\infty}.$$

Proof. The first inequality is contained in Proposition 4.6. The second inequality follows from Corollary 4.27 (alternatively, from Proposition 4.6 and Theorem 4.22). □

Example 6.2 (integral foliated simplicial volume of surfaces). Let M be an oriented closed connected surface of genus $g(M)$. Then

$$|M| = \begin{cases} 2 & \text{if } M \cong S^2, \\ \|M\| = \|M\|_{\mathbb{Z}}^{\infty} = 4 \cdot g(M) - 4 & \text{otherwise.} \end{cases}$$

Indeed, because S^2 is simply connected, $|S^2| = \|S^2\|_{\mathbb{Z}} = 2$. If $M \not\cong S^2$, then the classical computation of simplicial volume of aspherical surfaces [13, Section 0.2] shows that $\|M\| = \|M\|_{\mathbb{Z}}^{\infty} = 4 \cdot g(M) - 4$, and so the claim follows from Proposition 6.1.

Hence, in the case of finite fundamental group, we have

Corollary 6.3 (finite fundamental group). *Let M be an oriented closed connected manifold with finite fundamental group. Then*

$$|M| = \frac{1}{|\pi_1(M)|} \cdot \|\tilde{M}\|_{\mathbb{Z}} = \|M\|_{\mathbb{Z}}^{\infty},$$

where \tilde{M} is the universal covering of M . In contrast, $\|M\| = 0$.

Proof. The universal covering $\tilde{M} \rightarrow M$ has $|\pi_1(M)|$ sheets; because $\pi_1(M)$ is finite, \tilde{M} indeed is an oriented closed connected manifold. Hence, Theorem 4.22 gives us

$$|M| = \frac{1}{|\pi_1(M)|} \cdot |\tilde{M}|.$$

Moreover, $|\tilde{M}| = \|\tilde{M}\|_{\mathbb{Z}}$ because \tilde{M} is simply connected (Example 4.5). So, in combination with Proposition 6.1, we obtain

$$\|M\|_{\mathbb{Z}}^{\infty} \leq \frac{1}{|\pi_1(M)|} \cdot \|\tilde{M}\|_{\mathbb{Z}} = |M| \leq \|M\|_{\mathbb{Z}}^{\infty}.$$

On the other hand, simplicial volume of manifolds with finite (more generally, amenable) fundamental group is zero [13], and so $\|M\| = 0$. \square

6.2. Proof of Corollary 1.4. We will now prove Corollary 1.4 with help of the proportionality principle for hyperbolic manifolds (Corollary 1.3). More precisely, we will prove the following, slightly more general, statement:

Theorem 6.4. *Let $n \in \mathbb{N}$, and let M and N be oriented closed connected hyperbolic n -manifolds with fundamental groups Γ and Λ , and let $G := \text{Isom}^+(\mathbb{H}^n)$. Let S be a set of representatives of uniform lattices in G , containing a representative for every isometry class of oriented closed connected hyperbolic n -manifolds. The product $\prod_{\Lambda' \in S} G/\Lambda'$ is a standard Γ -space with respect to the diagonal translation action and the product of the probability measures induced by the (bi-invariant) Haar measure on G . Then*

$$|M| \leq |M| \prod_{\Lambda' \in S} G/\Lambda' \leq \frac{\text{vol}(M)}{\text{vol}(N)} \cdot \|N\|_{\mathbb{Z}}^{\infty}.$$

Notice that in every dimension up to isometry there are only countably many different oriented closed connected hyperbolic manifolds. Hence, S in the previous theorem is countable, and so the product $\prod_{\Lambda' \in S} G/\Lambda'$ indeed is a standard Γ -space.

Proof. We can view Γ and Λ as uniform lattices in $G = \text{Isom}^+(\mathbb{H}^n)$ and we have $M = \mathbb{H}^n/\Gamma$ and $\text{covol}(\Gamma) = \text{vol}(M)$ (and similarly for N and its finite coverings).

Let $N' \rightarrow N$ be a finite covering of N , let Λ' be the fundamental group of N' , and let $d := [\Lambda : \Lambda']$ be the number of sheets of this covering. From the proportionality principle for hyperbolic manifolds (Corollary 1.3), Proposition 4.13, and

Proposition 4.6 we obtain

$$\begin{aligned}
 |M| &= |\mathbb{H}^n / \Gamma| \\
 &\leq |\mathbb{H}^n / \Gamma| \prod_{\Lambda'' \in S} G/\Lambda'' \\
 &\leq |\mathbb{H}^n / \Gamma|^{G/\Lambda'} \\
 &= \frac{\text{covol}(\Gamma)}{\text{covol}(\Lambda')} \cdot |\mathbb{H}^n / \Lambda'|^{G/\Gamma} \\
 &= \frac{\text{covol}(\Gamma)}{\text{covol}(\Lambda)} \cdot \frac{1}{[\Lambda : \Lambda']} \cdot |N'|^{G/\Gamma} \\
 &\leq \frac{\text{vol}(M)}{\text{vol}(N)} \cdot \frac{1}{d} \cdot \|N'\|_{\mathbb{Z}}.
 \end{aligned}$$

Taking the infimum over all finite coverings of N finishes the proof. □

This concludes the proof of Corollary 1.4.

6.3. Proof of Theorem 1.5

Definition 6.5. Let Γ be a finitely generated group and let S be a set of finite index subgroups of Γ (hence, S is countable). Then the standard Γ -space $X_{\Gamma,S}$ is given by the set

$$\prod_{\Lambda \in S} \Gamma/\Lambda,$$

equipped with the product measure of the normalised counting measures and the diagonal left Γ -action given by translating cosets.

Notice that in the situation of Definition 6.5 the Γ -action on $X_{\Gamma,S}$ is free if and only if $\bigcap_{\Lambda \in S} \Lambda = \{e\}$, and that the Γ -action is not ergodic in general.

Theorem 6.6 (products of coset spaces as parameter space). *Let M be an oriented closed connected manifold with fundamental group Γ , let S be a set of finite index subgroups of Γ that is stable under finite intersections. For $\Lambda \in S$ we denote the covering space of M associated with the subgroup $\Lambda \subset \Gamma$ by M_{Λ} .*

(1) *Then*

$$|M|^{X_{\Gamma,S}} = \inf_{\Lambda \in S} \frac{1}{[\Gamma : \Lambda]} \cdot \|M_{\Lambda}\|_{\mathbb{Z}}$$

(2) *In particular, if S is the set of all finite index subgroups of Γ , then*

$$|M|^{X_{\Gamma,S}} = \|M\|_{\mathbb{Z}}^{\infty}.$$

The second part is nothing but Theorem 1.5.

Proof. By definition of the stable integral simplicial volume and because finite intersections of finite index subgroups of Γ have finite index in Γ , it suffices to prove the first part.

We first show that the right hand side is an upper bound for the left hand side: For all $\Lambda \in S$ we have

$$|M|^{X_{\Gamma,S}} \leq |M|^{\Gamma/\Lambda} = \frac{1}{[\Gamma : \Lambda]} \cdot \|M_{\Lambda}\|_{\mathbb{Z}}$$

by Proposition 4.13 and Corollary 4.27. Taking the infimum yields

$$|M|^{X_{\Gamma,S}} \leq \inf_{\Lambda \in S} \frac{1}{[\Gamma : \Lambda]} \cdot \|M_{\Lambda}\|_{\mathbb{Z}}.$$

It remains to show that the left hand side also is an upper bound for the right hand side. The main idea is to reduce the parameter space $X_{\Gamma,S}$ to finite products of coset spaces. So, let $F(S)$ be the set of finite subsets of S . For $F \in F(S)$ we write σ_F for the σ -algebra of the finite product $X_{\Gamma,F}$ (which is just the power set of $X_{\Gamma,F}$), we write

$$\pi_F: X_{\Gamma,S} \longrightarrow X_{\Gamma,F}$$

for the canonical projection, and we define

$$L_F := \{f \circ \pi_F \mid f \in L^{\infty}(X_{\Gamma,F}, \mathbb{Z})\} \subset L^{\infty}(X_{\Gamma,S}, \mathbb{Z}).$$

As first step, we will determine $|M|^{X_{\Gamma,F}}$ for $F \in F(S)$: for all $\gamma, \gamma' \in \Gamma$ we have

$$(\gamma \cdot \Lambda = \gamma' \cdot \Lambda, \text{ for all } \Lambda \in F) \iff \gamma \cdot \bigcap_{\Lambda \in F} \Lambda = \gamma' \cdot \bigcap_{\Lambda \in F} \Lambda.$$

Hence, $X_{\Gamma,F} = \prod_{\Lambda \in F} \Gamma/\Lambda$ is, as a Γ -parameter space, a finite convex combination of multiple copies of the coset space $\Gamma/\bigcap_{\Lambda \in F} \Lambda$. In view of Proposition 4.15 we obtain

$$|M|^{X_{\Gamma,F}} = |M|^{\Gamma/\bigcap_{\Lambda \in F} \Lambda}.$$

As second step, we will now show that

$$L := \bigcup_{F \in F(S)} L_F \subset L^{\infty}(X_{\Gamma,S}, \mathbb{Z})$$

is $\|\cdot\|_1$ -dense in $L^{\infty}(X_{\Gamma,S}, \mathbb{Z})$. The σ -algebra σ on the product space $X_{\Gamma,S}$ is the product σ -algebra of the power sets of all factors Γ/Λ with $\Lambda \in S$; i.e., σ is generated by $\bigcup_{F \in F(S)} \pi_F^{-1}(\sigma_F)$. We now consider the system

$$\sigma' := \{A \in \sigma \mid \chi_A \in \bar{L}^{\|\cdot\|_1}\}$$

of subsets of $X_{\Gamma,S}$. In order to show that L is $\|\cdot\|_1$ -dense in $L^\infty(X_{\Gamma,S}, \mathbb{Z})$ it suffices to prove that $\sigma' = \sigma$. It is easy to check that σ' is a σ -algebra on $X_{\Gamma,S}$. Moreover, by definition, $\bigcup_{F \in F(S)} \pi_F^{-1}(\sigma_F) \subset \sigma'$. Hence, $\sigma' = \sigma$, and so L is $\|\cdot\|_1$ -dense in $L^\infty(X_{\Gamma,S}, \mathbb{Z})$.

Therefore, the map

$$H_*(M, L) \longrightarrow H_*(M, L^\infty(X_{\Gamma,S}, \mathbb{Z}))$$

induced by the inclusion $L \hookrightarrow L^\infty(X_{\Gamma,S}, \mathbb{Z})$ of coefficient $\mathbb{Z}\Gamma$ -modules is isometric [29, Lemma 2.9][18, Proposition 1.7] (the cited proofs carry over to this integral setting) and the $X_{\Gamma,S}$ -fundamental class of M is contained in the image; notice that the union $L = \bigcup_{F \in F(S)} L_F$ indeed is a $\mathbb{Z}\Gamma$ -submodule of $L^\infty(X_{\Gamma,S}, \mathbb{Z})$.

Let $c \in C_*(M, L)$ be an L -fundamental cycle of M . By definition of L , there exists an $F \in F(S)$ such that $c \in C_*(M; L_F)$. Thus,

$$\begin{aligned} |c|^L &\geq |M|^{X_{\Gamma,F}} \\ &= |M|^{\Gamma/\bigcap_{\Lambda \in F} \Lambda} \\ &\geq \inf_{\Lambda \in S} |M|^{\Gamma/\Lambda} \\ &= \inf_{\Lambda \in S} \frac{1}{[\Gamma : \Lambda]} \cdot \|M_\Lambda\|_{\mathbb{Z}}; \end{aligned}$$

the first inequality is a consequence of the isometric isomorphism

$$L_F \longrightarrow L^\infty(X_{\Gamma,F}, \mathbb{Z})$$

induced from the projection π_F , the second equality was shown in the first step, the third inequality holds because S is assumed to be closed under finite intersections, and the last equality follows from Corollary 4.27. Taking the infimum over all fundamental cycles and taking the isometry

$$H_*(M; L) \longrightarrow H_*(M; L^\infty(X_{\Gamma,S}, \mathbb{Z}))$$

into account gives the desired estimate

$$|M|^{X_{\Gamma,S}} \geq \inf_{\Lambda \in S} \frac{1}{[\Gamma : \Lambda]} \cdot \|M_\Lambda\|_{\mathbb{Z}}. \quad \square$$

Remark 6.7 (inverse limits). The proof of Theorem 6.6 carries over to the following modification. Instead of the product $\prod_{\Lambda \in S} \Gamma/\Lambda$ we can also consider the inverse limit $I_{\Gamma,S}$ of the system $(\Gamma/\Lambda)_{\Lambda \in S}$ with respect to the canonical projection maps between coset spaces of nested subgroups; we equip $I_{\Gamma,S}$ with the Γ -action induced from the translation action on the coset spaces and we equip $I_{\Gamma,S}$ with the Borel probability space structure given by the discrete σ -algebras on the coset

spaces and the compatible system of normalised counting measures on the coset spaces. We then obtain

$$|M|^{I_{\Gamma,S}} = \inf_{\Lambda \in S} \frac{1}{[\Gamma : \Lambda]} \cdot \|M_\Lambda\|_{\mathbb{Z}}.$$

While this inverse limit space might be harder to visualise than the product space $X_{\Gamma,S}$, it does have the advantage that it is ergodic.

7. Integral foliated simplicial volume of hyperbolic 3-manifolds

We will now complete the proof of Theorem 1.1 by proving the following, more explicit, version:

Theorem 7.1 (integral foliated simplicial volume of hyperbolic 3-manifolds). *Let M be an oriented closed connected hyperbolic 3-manifold, and let S be a set of representatives of uniform lattices in $G := \text{Isom}^+(\mathbb{H}^3)$ containing a representative for every isometry class of oriented closed connected hyperbolic 3-manifolds (see Theorem 6.4). Then*

$$|M| = |M|^{\prod_{\Lambda \in S} G/\Lambda} = \|M\|.$$

Proof. Let M be an oriented closed connected hyperbolic 3-manifold. In view of Theorem 1.6 there exists a sequence $(M_n)_{n \in \mathbb{N}}$ of oriented closed connected hyperbolic 3-manifolds with

$$\lim_{n \rightarrow \infty} \frac{\|M_n\|_{\mathbb{Z}}^\infty}{\|M_n\|} = 1.$$

By Corollary 1.4 (and Theorem 6.4, for the concrete parameter space), for all $n \in \mathbb{N}$ we have

$$\|M\| \leq |M| \leq |M|^{\prod_{\Lambda \in S} G/\Lambda} \leq \frac{\text{vol } M}{\text{vol } M_n} \cdot \|M_n\|_{\mathbb{Z}}^\infty.$$

On the other hand, by the classical proportionality principle for simplicial volume for hyperbolic manifolds we have $\|M_n\| > 0$ and

$$\frac{\text{vol } M}{\text{vol } M_n} = \frac{\|M\|}{\|M_n\|},$$

and so

$$\|M\| \leq |M| \leq |M|^{\prod_{\Lambda \in S} G/\Lambda} \leq \frac{\|M\|}{\|M_n\|} \cdot \|M_n\|_{\mathbb{Z}}^\infty.$$

Because of $\lim_{n \rightarrow \infty} \|M_n\|_{\mathbb{Z}}^\infty / \|M_n\| = 1$ the right hand side converges to $\|M\|$ for $n \rightarrow \infty$. Hence, $\|M\| = |M|^{\prod_{\Lambda \in S} G/\Lambda} = |M|$, as desired. \square

Similarly to Francaviglia, Frigerio, and Martelli [8, Question 6.4], we hence ask:

Question 7.2. *Does the integral foliated simplicial volume of oriented closed connected hyperbolic manifolds of dimension bigger than 3 also coincide with the simplicial volume? Does this even hold for all oriented closed connected aspherical manifolds?*

8. Stable integral simplicial volume and integral foliated simplicial volume of Seifert 3-manifolds

For the sake of completeness, we add also the computation of stable integral simplicial volume and integral foliated simplicial volume of Seifert 3-manifolds:

Proposition 8.1 (Seifert case). *Let M be an oriented compact connected Seifert manifold with $|\pi_1(M)| = \infty$. Then $\|M\|_{\mathbb{Z}}^{\infty} = 0$.*

Proof. We follow step by step the related argument of Francaviglia, Frigerio, and Martelli [8, Proposition 5.11]. A Seifert manifold has a finite covering that is an S^1 -bundle over an orientable surface Σ with some Euler number $e \geq 0$. If the manifold M has boundary, then $e = 0$ and the bundle is a product $S^1 \times \Sigma$. Since $S^1 \times \Sigma$ covers itself with arbitrarily high degree, $S^1 \times \Sigma$ (and hence also M) clearly has stable integral simplicial volume 0.

If M is closed, we denote the covering mentioned above by (Σ, e) . A closed connected orientable S^1 -bundle over an orientable surface either is irreducible or $S^1 \times S^2$ or $\mathbb{RP}^3 \# \mathbb{RP}^3$. Since $\mathbb{RP}^3 \# \mathbb{RP}^3$ admits a double self-covering, in this case the stable integral simplicial volume vanishes. Moreover, we have already considered the case $S^1 \times \Sigma$. Therefore we restrict to irreducible S^1 -bundles. By assumption, (Σ, e) is a closed connected orientable irreducible Seifert manifold with $|\pi_1((\Sigma, e))| = \infty$. Thus, by Remark 3.4, the complexity is equal to the special complexity and it has the following bound [23]:

$$c_S(\Sigma, e) \leq e + 6 \cdot \chi_-(\Sigma) + 6$$

where $\chi_-(\Sigma) = \max\{-\chi(\Sigma), 0\}$.

Let now T be a minimal triangulation of (Σ, e) , and let T' be its first barycentric subdivision. Then T' is a semi-simplicial triangulation that allows us to define an integral fundamental cycle. Since the special complexity of (Σ, e) is equal to the minimal number of tetrahedra in a triangulation and the barycentric subdivision of Δ^3 consists of 24 tetrahedra, we have

$$\|(\Sigma, e)\|_{\mathbb{Z}} \leq 24 \cdot c_S(\Sigma, e) \leq 24 \cdot (e + 6 \cdot \chi_-(\Sigma) + 6).$$

Now for every $d \in \mathbb{N}$ we construct a degree- d^2 covering $(\bar{\Sigma}, e) \rightarrow (\Sigma, e)$ where $\bar{\Sigma}$ is a d -sheeted covering space of Σ as in the argument of Francaviglia, Frigerio, and Martelli [8, Proposition 5.11] in order to conclude. \square

In particular, we obtain:

Corollary 8.2. *Let M be an oriented closed connected Seifert manifold with $|\pi_1(M)| = \infty$. Then*

$$\|M\| = |M| = \|M\|_{\mathbb{Z}}^{\infty} = 0.$$

By Perelman's result a closed Seifert manifold with finite fundamental group admits an elliptic structure (i.e., it is a quotient of S^3 by a finite subgroup of $\text{SO}(4)$ acting by rotations).

Proposition 8.3. *Let $n \in \mathbb{N}_{>0}$, and let M be an oriented closed connected elliptic n -manifold. Then*

$$|M| = \|M\|_{\mathbb{Z}}^{\infty} = \frac{1}{|\pi_1(M)|} \cdot \begin{cases} 1 & \text{if } n \text{ is odd,} \\ 2 & \text{if } n \text{ is even.} \end{cases}$$

In contrast, $\|M\| = 0$.

Proof. Corollary 6.3 shows that

$$|M| = \|M\|_{\mathbb{Z}}^{\infty} = \frac{1}{|\pi_1(M)|} \cdot \|S^n\|_{\mathbb{Z}}.$$

Clearly, $\|S^n\|_{\mathbb{Z}} = 1$ if n is odd, and $\|S^n\|_{\mathbb{Z}} = 2$ if n is even.

On the other hand, the simplicial volume of an elliptic manifold is always zero because simplicial volume is multiplicative under finite coverings [19, Proposition 4.1] and $\|S^n\| = 0$ for all $n \in \mathbb{N}_{>0}$. \square

References

- [1] U. Bader, A. Furman, and R. Sauer, Efficient subdivision in hyperbolic groups and applications. *Groups Geom. Dyn.* **7** (2013), no. 2, 263–292. [Zbl 06190853](#)
[MR 3054570](#)
- [2] U. Bader, A. Furman, and R. Sauer, Integrable measure equivalence and rigidity of hyperbolic lattices. *Invent. Math.* **194** (2013), no. 2, 313–379. [Zbl 1279.22012](#)
[MR 3117525](#)

- [3] M. B. Bekka and M. Mayer, *Ergodic theory and topological dynamics of group actions on homogeneous spaces*. London Mathematical Society Lecture Note Series, 269. Cambridge University Press, Cambridge, 2000. [Zbl 0961.37001](#) [MR 1781937](#)
- [4] M. Bucher-Karlsson, The proportionality constant for the simplicial volume of locally symmetric spaces. *Colloq. Math.* **111** (2008), no. 2, 183–198. [Zbl 1187.53042](#) [MR 2365796](#)
- [5] M. Bucher, M. Burger, and A. Iozzi, A dual interpretation of the Gromov–Thurston proof of Mostow rigidity and volume rigidity for representations of hyperbolic lattices. In M. A. Picardello (ed.), *Trends in harmonic analysis*. Springer INdAM Series, 3. Springer, Milan, 2013, 47–76. [Zbl 1268.53056](#) [MR 3026348](#)
- [6] R. H. Farrell, Representation of invariant measures. *Illinois J. Math.* **6** (1962), 447–467. [Zbl 0108.14203](#) [MR 0150264](#)
- [7] S. Francaviglia, Hyperbolic volume of representations of fundamental groups of cusped 3-manifolds. *Int. Math. Res. Not.* **2004** (2004), no. 9, 425–459. [Zbl 1088.57015](#) [MR 2040346](#)
- [8] S. Francaviglia, R. Frigerio, and B. Martelli, Stable complexity and simplicial volume of manifolds. *J. Topol.* **5** (2012), no. 4, 977–1010. [Zbl 1266.57015](#) [MR 3001317](#)
- [9] R. Frigerio, (Bounded) continuous cohomology and Gromov’s proportionality principle. *Manuscripta Math.* **134** (2011), no. 3–4, 435–474. [Zbl 1220.55003](#) [MR 2765720](#)
- [10] R. Frigerio and C. Pagliantini, The simplicial volume of hyperbolic manifolds with geodesic boundary. *Algebr. Geom. Topol.* **10** (2010), no. 2, 979–1001. [Zbl 1206.53045](#) [MR 2629772](#)
- [11] K. Fujiwara and J. K. Manning, Simplicial volume and fillings of hyperbolic manifolds. *Algebr. Geom. Topol.* **11** (2011), no. 4, 2237–2264. [Zbl 1244.53051](#) [MR 2826938](#)
- [12] A. Furman, Gromov’s measure equivalence and rigidity of higher rank lattices. *Ann. of Math. (2)* **150** (1999), no. 3, 1059–1081. [Zbl 0943.22013](#) [MR 1740986](#)
- [13] M. Gromov, Volume and bounded cohomology. *Inst. Hautes Études Sci. Publ. Math.* **56** (1982), 5–99 (1983). [Zbl 0516.53046](#) [MR 0686042](#)
- [14] M. Gromov, Asymptotic invariants of infinite groups. In G. A. Niblo and M. A. Roller (eds.), *Geometric group theory*. Vol. 2. (Sussex, 1991). London Mathematical Society Lecture Note Series, 182. Cambridge University Press, Cambridge, 1993, 1–295. [Zbl 0841.20039](#) [MR 1253544](#)
- [15] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces*. With appendices by M. Katz, P. Pansu, and S. Semmes. Translated from the French by S. M. Bates. Progress in Mathematics, 152. Birkhäuser Boston, Inc., Boston, MA, 1999. [Zbl 0953.53002](#) [MR 1699320](#)
- [16] A. Kechris, *Classical descriptive set theory*. Graduate Texts in Mathematics, 156. Springer-Verlag, New York, 1995. [Zbl 0819.04002](#) [MR 1321597](#)
- [17] F. Lafont and B. Schmidt, Simplicial volume of closed locally symmetric spaces of non-compact type. *Acta Math.* **197** (2006), no. 1, 129–143. [Zbl 1111.57020](#) [MR 2285319](#)

- [18] C. Löh, ℓ^1 -homology and simplicial volume. Ph.D. thesis, Westfälischen Wilhelms-Universität Münster, Münster, 2007. [Zbl 1152.57304](#)
<http://nbn-resolving.de/urn:nbn:de:hbz:6-37549578216>
- [19] C. Löh, Simplicial volume. *Bull. Man. Atl.* (2011), 7–18.
<http://www.boma.mpim-bonn.mpg.de/data/29print.pdf>
- [20] C. Löh and R. Sauer, Degree theorems and Lipschitz simplicial volume for non-positively curved manifolds of finite volume. *J. Topol.* **2** (2009), no. 1, 193–225.
[Zbl 1187.53043](#) [MR 2499443](#)
- [21] B. Martelli, Complexity of PL manifolds. *Algebr. Geom. Topol.* **10** (2010), no. 2, 1107–1164. [Zbl 1222.57021](#) [MR 2653058](#)
- [22] B. Martelli, Complexity of 3-manifolds. In Y. N. Minsky, M. Sakuma, and C. Series (eds.), *Spaces of Klenian groups*. (Cambridge, 2003) London Mathematical Society Lecture Note Series, 329. Cambridge University Press, Cambridge, 2006, 91–120.
[Zbl 1099.57018](#) [MR 2258746](#)
- [23] B. Martelli and C. Petronio, Complexity of geometric three-manifolds. *Geom. Dedicata* **108** (2004), 15–69. [Zbl 1068.57011](#) [MR 2112664](#)
- [24] S. Matveev, Complexity theory of three-dimensional manifolds. *Acta Appl. Math.* **19** (1990), no. 2, 101–130. [Zbl 0724.57012](#) [MR 1074221](#)
- [25] S. Matveev, *Algorithmic topology and classification of 3-manifolds*. Algorithms and Computation in Mathematics, 9. Springer-Verlag, Berlin, 2003. [Zbl 1048.57001](#)
[MR 1997069](#)
- [26] T. S. Mrowka and P. S. Ozsváth (eds.), *Low dimensional topology*. (Graduate Summer School, Park City, UT, 2006) IAS/Park City Mathematics Series, 15. American Mathematical Society, Providence, R.I., 2009. [Zbl 1161.57010](#) [MR 2499604](#)
- [27] W. D. Neumann and A. W. Reid, Arithmetic of hyperbolic manifolds. In B. Apanasov, W. D. Neumann, A. W. Reid and L. Siebenmann (eds.), *Topology '90* (Columbus, OH, 1990) Ohio State University Mathematical Research Institute Publications, 1. Walter de Gruyter & Co., Berlin, 1992. [Zbl 1184416](#) [MR 1184397](#)
- [28] R. Sauer, Amenable covers, volume and L^2 -Betti numbers of aspherical manifolds. *J. Reine Angew. Math.* **636** (2009), 47–92. [Zbl 1261.53033](#) [MR 2572246](#)
- [29] M. Schmidt, L^2 -Betti numbers of \mathcal{R} -spaces and the integral foliated simplicial volume. Ph.D. Thesis. Westfälische Wilhelms-Universität Münster, Münster, 2005.
[Zbl 1076.57024](#) <http://nbn-resolving.de/urn:nbn:de:hbz:6-05699458563>
- [30] C. Strohm (= C. Löh), *The proportionality principle of simplicial volume*. Diplomarbeit, Westfälische Wilhelms-Universität Münster, Münster, 2004.
[arXiv:math.AT/0504106](http://arxiv.math.AT/0504106) [math.AT]
- [31] M. Stroppel, *Locally compact groups*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2006. [Zbl 1255.22004](#) [MR 2226087](#)
- [32] W. P. Thurston, The geometry and topology of 3-manifolds. Mimeographed notes, 1979. <http://library.msri.org/books/gt3m/PDF/13.pdf>

- [33] V. S. Varadarajan, Groups of automorphisms of Borel spaces. *Trans. Amer. Math. Soc.* **109** (1963), 191–220. [Zbl 0192.14203](#) [MR 0159923](#)
- [34] C. T. C. Wall, *Surgery on compact manifolds*. London Mathematical Society Monographs, 1. Academic Press, London and New York, 1970. [Zbl 0935.57003](#) [MR 0431216](#)

Received April 4, 2014

Clara Löh, Universität Regensburg, Fakultät für Mathematik, Universitätsstraße 31,
93053 Regensburg, Germany

e-mail: clara.loeh@mathematik.uni-r.de

Cristina Pagliantini, ETH Zürich, Department of Mathematics, Rämistrasse 101,
8092 Zürich, Switzerland

e-mail: cristina.pagliantini@math.ethz.ch