



Improved quasi parton distribution through Wilson line renormalization

Jiunn-Wei Chen^{a,b}, Xiangdong Ji^{c,d}, Jian-Hui Zhang^{e,*}

^a Department of Physics, Center for Theoretical Sciences, and Leung Center for Cosmology and Particle Astrophysics, National Taiwan University, Taipei, 106, Taiwan

^b Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^c INPAC, Department of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai, 200240, PR China

^d Maryland Center for Fundamental Physics, Department of Physics, University of Maryland, College Park, MD 20742, USA

^e Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany

Received 12 October 2016; received in revised form 24 November 2016; accepted 4 December 2016

Available online 9 December 2016

Editor: Hong-Jian He

Abstract

Recent developments showed that hadron light-cone parton distributions could be directly extracted from spacelike correlators, known as quasi parton distributions, in the large hadron momentum limit. Unlike the normal light-cone parton distribution, a quasi parton distribution contains ultraviolet (UV) power divergence associated with the Wilson line self energy. We show that to all orders in the coupling expansion, the power divergence can be removed by a “mass” counterterm in the auxiliary z -field formalism, in the same way as the renormalization of power divergence for an open Wilson line. After adding this counterterm, the quasi quark distribution is improved such that it contains at most logarithmic divergences. Based on a simple version of discretized gauge action, we present the one-loop matching kernel between the improved non-singlet quasi quark distribution with a lattice regulator and the corresponding quark distribution in dimensional regularization.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

* Corresponding author.

E-mail address: jianhui.zhang@physik.uni-regensburg.de (J.-H. Zhang).

1. Introduction

One of the most important goals of QCD is to understand the hadron structure from its fundamental degrees of freedom – quarks and gluons. This necessarily goes to the nonperturbative regime of QCD, and it is difficult in general to obtain first principle results directly from the QCD Lagrangian. A powerful tool of obtaining such results is Lattice QCD, which is an approach defined on Euclidean spacetime, and has been used to calculate hadron masses, charges, etc., to a remarkable accuracy. However, it cannot be used to directly access intrinsically Minkowskian quantities such as the parton distribution functions (PDFs). PDFs are defined as the forward hadronic matrix elements of light-cone correlations, and describe the momentum distribution of quarks and gluons inside the hadron. They play a crucial role in understanding the experimental data at high energy hadron colliders such as the LHC. Owing to their real time dependence, Lattice QCD can only be used to indirectly extract the information on the PDFs by calculating their moments, which is limited by technical complications such as operator mixing. Another commonly used approach to determine PDFs is to assume a suitable parametrized form and fit to a large variety of experimental data. Most PDF groups obtained their PDF sets in this way [1–6]. A main drawback of this approach is that it suffers from parametrization uncertainty, and different groups usually produce different results for the same PDF.

Recent developments [7–21] showed that the light-cone observables such as the PDFs can be directly extracted from the large momentum limit of the hadronic matrix element of a space-like correlator, which is known as the quasi observable, using a large momentum effective theory (LaMET) [15] (for other approaches to extract light-cone quantities see e.g. [22–28]). The quasi PDF does not have a real time dependence, and thus can be simulated on the lattice. The infrared (IR) behaviors between the flavor non-singlet quasi-PDF and its corresponding light-cone PDF are shown to be the same at one loop by direct computations, and argued to be the same at all loops in Ref. [9], based on which a factorization formula was also presented.

The factorization in Ref. [9] was given for the bare quasi quark distribution, where all fields and couplings entering the quasi distribution are bare ones. However, as the light-cone distribution, the quasi distribution also contains ultraviolet (UV) divergences and therefore needs renormalization. Ref. [17] explores the renormalization property of the quasi distribution, and shows that it is multiplicatively renormalizable at two-loop order. Also an equivalence was established between the virtual correction of the quasi quark distribution and the correction to the heavy-light quark vector current in heavy quark effective theory, so that the UV divergences in the former can be renormalized as the renormalization of the latter. Dimensional regularization was used in Ref. [17], and the linear divergence present in a cutoff or lattice regularization was ignored. In realistic lattice calculations, one needs to know how to deal with such power divergences. This is one goal of the present paper. We will show that the power divergence in the quasi quark distribution can be removed to all-loop orders by a mass counterterm, which is the same as the renormalization of an open Wilson line. After such a mass renormalization, the quasi quark distribution is improved such that it contains at most logarithmic divergences.

The rest of this paper is organized as follows. In Sec. 2, we discuss the renormalization of power divergences arising from the Wilson line self energy in the quasi quark distribution. In Sec. 3, we present a lattice perturbation theory matching for the quasi quark distribution. We then conclude in Sec. 4.




Fig. 1. One-loop diagrams for quasi quark distribution in Feynman gauge, the conjugate diagrams are not shown.

2. Improved quasi quark distribution through Wilson line renormalization

Let us start by recalling the definition of the quasi quark distribution. For the unpolarized quark density, it is given as [8]

$$\tilde{q}(x, \Lambda, p^z) = \int_{-\infty}^{\infty} \frac{dz}{4\pi} e^{izkz} \langle p | \bar{\psi}(0, 0_{\perp}, z) \gamma^z L(z, 0) \psi(0) | p \rangle, \tag{1}$$

where the quark fields are separated along the spatial z -direction, $L(z, 0)$ is the Wilson line gauge link inserted to ensure gauge invariance, and Λ denotes the UV cutoff.

The one-loop correction to the above quark density has been computed both in the axial gauge [9] and in the Feynman gauge [17]. In Ref. [17], the Wilson line $L(z, 0)$ was separated into two Wilson lines along the z -axis $L(\infty, 0)$ and $L(z, \infty)$, and associated respectively to the quark field to form gauge invariant combinations. Here we keep the gauge link as $L(z, 0)$. The one-loop diagrams are then given in Fig. 1. It is straightforward to check that these diagrams yield the same one-loop result as that in Ref. [17]. The linear divergence comes from the last diagram, which is a Wilson line self energy. It is known that such a linear divergence can be removed by a mass renormalization with the help of an auxiliary z -field formalism [29–31], where the z -field is defined in a one-dimensional parameter space, and the non-local Wilson line can be interpreted as a two-point function of the z -field. In a sense, the auxiliary field $Z(z)$ can be understood as a Wilson line extending between $[z, \infty]$, i.e.

$$Z(z) = L(z, \infty), \tag{2}$$

which satisfies an equation of motion

$$[\partial_z - ig A_z(z)] Z(z) = 0, \tag{3}$$

analogous to a heavy quark field (in the heavy quark limit). The Wilson line

$$L(z, 0) = Z(z) Z^{\dagger}(0) \tag{4}$$

also renormalizes analogously to a heavy quark two point function

$$L^{\text{ren}}(z, 0) = \mathcal{Z}_Z^{-1} e^{-\delta m |z|} L(z, 0), \tag{5}$$

where δm is analogous to the dimension one heavy quark mass counterterm and is linearly divergent, while the dimension zero wave function renormalization constant \mathcal{Z}_Z is logarithmically divergent [29].

To illustrate the cancelation of linear divergence from the mass counterterm, let us determine δm at one-loop level. Recall that the last diagram in Fig. 1 leads to the following linear divergence

$$\lim_{\epsilon \rightarrow 0} \int dk_z \frac{\alpha_s C_F \Lambda}{2\pi} \frac{[\delta(k_z - \bar{x} p_z) - \delta(\bar{x} p_z)] p_z}{k_z^2 + \epsilon^2} \tag{6}$$

with $\bar{x} = 1 - x$. On the other hand, plugging Eq. (5) into Eq. (1), one finds that the mass counterterm yields the following contribution

$$\begin{aligned}
& - \int \frac{dz}{2\pi} p_z e^{i(x-1)p_z z} |z| \delta m = - \lim_{\epsilon \rightarrow 0} \int \frac{dz}{2\pi} p_z e^{-i\bar{x}p_z z} \frac{1 - e^{-\epsilon|z|}}{\epsilon} \delta m \\
& = - \lim_{\epsilon \rightarrow 0} \int \frac{dz}{2\pi} p_z e^{-i\bar{x}p_z z} \int_0^\infty \frac{d\alpha}{\sqrt{\pi\alpha}} (1 - e^{-\frac{z^2}{4\alpha}}) e^{-\alpha\epsilon^2} \delta m \\
& = - \lim_{\epsilon \rightarrow 0} \int \frac{dz}{2\pi} p_z e^{-i\bar{x}p_z z} \int_0^\infty d\alpha \int dk_z \frac{e^{-\alpha(k_z^2 + \epsilon^2)} (1 - e^{ik_z z})}{\pi} \delta m \\
& = - \lim_{\epsilon \rightarrow 0} \int \frac{dz}{2\pi} p_z e^{-i\bar{x}p_z z} \int dk_z \frac{1}{\pi} \frac{(1 - e^{ik_z z})}{k_z^2 + \epsilon^2} \delta m \\
& = - \lim_{\epsilon \rightarrow 0} \int \frac{dk_z}{\pi} p_z \frac{\delta(\bar{x}p_z) - \delta(k_z - \bar{x}p_z)}{k_z^2 + \epsilon^2} \delta m. \tag{7}
\end{aligned}$$

We therefore have

$$\delta m = - \frac{\alpha_s C_F}{2\pi} (\pi \Lambda) \tag{8}$$

at one-loop.

In coordinate space, by expanding the Wilson line to $\mathcal{O}(g^2)$, we have

$$- g^2 C_F \int_0^z dz_1 \int_0^{z_1} dz_2 \frac{1}{4\pi^2} \frac{1}{(z_1 - z_2)^2 + a^2} = - \frac{g^2 C_F}{4\pi^2} \left[\frac{z}{a} \tan^{-1} \left(\frac{z}{a} \right) - \frac{1}{2} \ln \left(1 + \frac{z^2}{a^2} \right) \right], \tag{9}$$

where we have introduced a cutoff a to regularize the short distance singularity in the coordinate space gluon propagator when $z_1 \rightarrow z_2$. By Fourier transforming to momentum space, we have

$$\begin{aligned}
& - \int \frac{dz}{2\pi} p_z e^{i(x-1)p_z z} \frac{g^2 C_F}{4\pi^2} \left[\frac{z}{a} \tan^{-1} \left(\frac{z}{a} \right) - \frac{1}{2} \ln \left(1 + \frac{z^2}{a^2} \right) \right] \\
& = - \lim_{\epsilon \rightarrow 0} \int \frac{dz}{2\pi} p_z e^{-i\bar{x}p_z z} \frac{g^2 C_F}{4\pi^2} \left[\frac{z}{a} \tan^{-1} \left(\frac{z}{a} \right) - \frac{1}{2} \ln \left(1 + \frac{z^2}{a^2} \right) \right] e^{-\epsilon|z|} \\
& = \frac{g^2 C_F}{8\pi^2} \left[\frac{1}{ap_z(1-x)^2} - \frac{1}{|1-x|} + C\delta(1-x) \right], \tag{10}
\end{aligned}$$

where C is a constant that can be determined by demanding the x -integration of the above expression to vanish. We can therefore identify

$$\Lambda = \frac{1}{a}. \tag{11}$$

The mass counterterm diagram of order α_s is shown in Fig. 2. The gauge independence of δm can already be seen from that the one-loop correction in the axial gauge [9] and Feynman gauge [17] contains identical linear divergences. By using the R_ξ gauge, one easily finds that the extra piece $k^\mu k^\nu / k^2$ (where k is the gluon momentum) in the numerator of gluon propagator has no impact on the linear divergence, therefore δm remains the same in different gauges.

One can add more gluons and quarks to the one-loop diagrams in Fig. 1 to form higher loop diagrams. Power counting tells us that power divergence appears only in those diagrams which




Fig. 2. One-loop mass counterterm diagram for quasi quark distribution in Feynman gauge.




Fig. 3. The extra one-loop diagram for quasi quark distribution on the lattice, the conjugate diagram is not shown.

contain the Wilson line self energy as a subdiagram. As shown in the appendix, beyond one-loop the mass counterterm also removes all the power divergence from the Wilson line. Therefore, after including the mass counterterm contribution, there is at most logarithmic divergence in the quasi quark distribution. This indicates that, as far as the power divergence is concerned, the renormalization of a quark bilinear operator like the quasi quark distribution is the same as the renormalization of an open Wilson line. In the discussions above, we showed that the linear divergence is indeed canceled by the mass counterterm at perturbative one-loop level. On the lattice, the mass counterterm δm has to be determined nonperturbatively, which can be done e.g. as in Ref. [32].

3. Matching between lattice and continuum through lattice perturbation theory

In this section, we aim at establishing the one-loop matching connecting the quasi quark distribution on the lattice to the normal quark distribution in the continuum. On the lattice, besides the diagrams in Fig. 1, there exists an extra one-loop diagram (shown in Fig. 3).

The first diagram in Fig. 1 and the diagram in Fig. 3 are nothing but the Feynman gauge quark field renormalization, and have been well understood in lattice QCD. Their contribution can also be obtained from the other diagrams in Fig. 1 through quark number conservation. As shown in Refs. [9] and [17], in the continuum the second and third diagrams in Fig. 1 contain a logarithmic dependence on the hadron momentum instead of a logarithmic divergence. Their contribution is independent of the regularization and thus will be the same in the continuum and on the lattice. One only needs to match the linearly divergent part of the last diagram in Fig. 1 between lattice and continuum. This gives a connection between the lattice cutoff and the linear divergence in the one-loop kernel in the continuum [9] (in general they will differ by a finite term). To obtain this connection, we compute the Wilson line self energy using lattice perturbation theory based on a simple version of discretized gauge action.

The contribution of the Wilson line self energy diagram can be written as

$$\begin{aligned} \delta W &= \frac{\alpha_s C_F}{8\pi^3} \int d^4(k a_L) \frac{1 - \cos(k_z z)}{(1 - \cos(k_z a_L)) 4 \sum_\lambda \sin^2 \frac{k_\lambda a_L}{2}} \\ &= \alpha_s C_F \left[\frac{\pi |z|}{2 a_L} - \frac{1}{\pi} \ln \frac{\pi |z|}{a_L} - \frac{1}{\pi} \right] + \mathcal{O}\left(\frac{a_L}{z}\right), \end{aligned} \tag{12}$$

where a_L denotes the lattice spacing. Comparing this with the expansion of Eq. (9) leads to the following matching between the linear divergence in the continuum and on the lattice

$$\Lambda = \frac{\pi}{a_L}. \quad (13)$$

By replacing Λ in the one-loop kernel of Ref. [9] with the above matching condition, we then obtain the one-loop kernel relating the quasi PDF on the lattice to the normal PDF. By adding a factor of $e^{-\delta m|z|}$ to the integrand of Eq. (1), we obtain the improved quasi-PDF \tilde{q}_{imp} free of power divergence

$$\tilde{q}_{\text{imp}}(x, \Lambda, p^z) = \int_{-\infty}^{\infty} \frac{dz}{4\pi} e^{izk^z - \delta m|z|} \langle p | \bar{\psi}(0, 0_{\perp}, z) \gamma^z L(z, 0) \psi(0) | p \rangle. \quad (14)$$

The matching kernel Z between the improved quasi-PDF \tilde{q}_{imp} and the PDF q is defined as

$$\begin{aligned} \tilde{q}_{\text{imp}}(x, a_L, p^z) &= \int_{-1}^1 \frac{dy}{|y|} Z\left(\frac{x}{y}, p^z a_L, \frac{\mu}{p^z}\right) q(y, \mu) \\ &+ \mathcal{O}\left(\Lambda_{\text{QCD}}^2/(p^z)^2, M^2/(p^z)^2\right). \end{aligned} \quad (15)$$

At one-loop the Z factor can be extracted from Eqs. (15)–(19) of Ref. [9] with the $1/a_L$ contribution subtracted by the counterterm diagram in Fig. 2.

$$Z\left(\xi, p^z a_L, \frac{\mu}{p^z}\right) = \delta(\xi - 1) + \frac{\alpha_s}{2\pi} Z^{(1)}\left(\xi, p^z a_L, \frac{\mu}{p^z}\right) + \dots, \quad (16)$$

where

$$Z^{(1)}/C_F = \begin{cases} \left(\frac{1+\xi^2}{1-\xi}\right) \ln \frac{\xi}{\xi-1} + 1, & \xi > 1, \\ \left(\frac{1+\xi^2}{1-\xi}\right) \ln \frac{(p^z)^2}{\mu^2} + \left(\frac{1+\xi^2}{1-\xi}\right) \ln [4\xi(1-\xi)] - \frac{2\xi}{1-\xi} + 1, & 0 < \xi < 1, \\ \left(\frac{1+\xi^2}{1-\xi}\right) \ln \frac{\xi-1}{\xi} - 1, & \xi < 0, \end{cases} \quad (17)$$

and near $\xi = 1$, one has an extra contribution from the quark self energy

$$Z^{(1)}(\xi) = \delta Z^{(1)}(2\pi/\alpha_s)\delta(\xi - 1) \quad (18)$$

with

$$\delta Z^{(1)} = \frac{\alpha_s C_F}{2\pi} \int dy \begin{cases} -\frac{1+y^2}{1-y} \ln \frac{y}{y-1} - 1, & y > 1, \\ -\frac{1+y^2}{1-y} \ln \frac{(p^z)^2}{\mu^2} - \frac{1+y^2}{1-y} \ln [4y(1-y)] \\ \quad + \frac{2y(2y-1)}{1-y} + 1, & 0 < y < 1, \\ -\frac{1+y^2}{1-y} \ln \frac{y-1}{y} + 1, & y < 0. \end{cases} \quad (19)$$

Given that the power divergence has been removed, the above one-loop matching kernel facilitates a reliable extraction of normal PDFs from lattice data. As already mentioned at the end of Sec. 2, calculating the improved quasi-PDF requires a non-perturbative determination of the mass counterterm δm . One possibility is following the procedure based on the static quark potential for the renormalization of Polyakov loop in Ref. [32], where the static quark potential for a system of a heavy quark and antiquark follows from the asymptotic behavior of a rectangular Wilson loop with two sides along T direction, and the Wilson loop is renormalized such that

the renormalized static quark potential obtains a constant offset proportional to δm . δm is then obtained by imposing an appropriate renormalization condition on the static quark potential. It is also worthwhile to compare the improved quasi PDF with the original one. As shown by the one-loop calculation, δm is negative. The exponential factor $e^{-\delta m|z|}$ then increases the weight of matrix elements with relatively large z , and therefore will increase the contribution at relatively small momentum when Fourier transforming to momentum space.

4. Conclusion

In this paper, we have shown that the power divergence present in the lattice regularization of the quasi PDFs can be removed to all-loop orders by a mass counterterm, which can be interpreted as the mass renormalization for a test particle moving on the Wilson line. After such a mass renormalization, the quasi distribution is improved such that it contains at most logarithmic divergences. We also present the one-loop matching for the quasi quark distribution between lattice and continuum using lattice perturbation theory based on a simple version of discretized gauge action. There are still practical issues in computing quasi PDFs on the lattice. In addition to the typical problems in lattice QCD of using finite spacetime volume and larger-than-physical light quark masses, it is desirable to reduce the lattice spacing such that the nucleon momentum is increased and the errors from finite momentum corrections are reduced, also the Lorentz symmetry, which is important in the LaMET formulation, is better preserved. In the future, the matching factor should be computed with the specific lattice action used in the lattice computation and even using non-perturbative renormalization on the lattice side. Despite these practical issues, our result improves the understanding of the renormalization property of quasi PDFs which is an important step towards a reliable extraction of physical PDFs from lattice data.

5. Note added in proof

While this work is being finalized, a preprint by Ishikawa et al. [33] dealing with the same topic has appeared, where they obtain similar conclusions as ours on the renormalization of power divergence.

Acknowledgements

We thank Will Detmold, David Lin, Andreas Schäfer and Iain Stewart for useful discussions. This work was partially supported by the U.S. Department of Energy via grants DE-FG02-93ER-40762, a grant (No. 11DZ2260700) from the Office of Science and Technology in Shanghai Municipal Government, grants from National Science Foundation of China (No. 11175114, No. 11405104), a Deutsche Forschungsgemeinschaft grant SCHA 458/20-1, the Ministry of Science and Technology, Taiwan under Grant Nos. 105-2112-M-002-017-MY3 and 105-2918-I-002-003 and the CASTS of National Taiwan University.

Appendix A

In this appendix, we present the arguments that the mass counterterm removes the power divergence to all-loop orders.

For simplicity, we start with the one-loop diagrams for the quasi quark distribution in Ref. [17], which can be equivalently drawn as the diagrams in Fig. 4 with a vertex insertion




Fig. 4. One-loop diagrams to illustrate the contribution to quasi quark distribution in Feynman gauge.

(1 for virtual diagrams, and $\gamma^z \delta(x - k^z/p^z)$ for real diagrams to impose momentum constraint). At one-loop, it has been shown [17] that the virtual diagrams are respectively equivalent to the wave function renormalization of a light quark field, the renormalization of a heavy-light vector current, and the wave function renormalization of a heavy quark field. All these can be done to all-loop orders. Moreover, the mass counterterm can be chosen to exactly cancel the linearly divergent contribution from the third diagram in Fig. 4 (both virtual contribution with vertex insertion 1 and real contribution with $\gamma^z \delta(x - k^z/p^z)$). Following Ref. [30], this also holds to all-loop orders. After this mass renormalization, as the real diagrams contain an extra δ -function which effectively reduces the UV degree of divergence by one, they will produce at most logarithmic divergences at higher-loop orders. Therefore, as far as power divergence is concerned, the mass renormalization is sufficient to remove them to all-loop orders.

References

- [1] S. Alekhin, J. Blumlein, S. Moch, *Phys. Rev. D* 86 (2012) 054009, arXiv:1202.2281 [hep-ph].
- [2] J. Gao, M. Guzzi, J. Huston, H.-L. Lai, Z. Li, P. Nadolsky, J. Pumplin, D. Stump, et al., arXiv:1302.6246 [hep-ph].
- [3] H1 Collaboration, ZEUS Collaboration, V. Radescu, PoS ICHEP2010 (2010) 168.
- [4] A.M. Cooper-Sarkar, ZEUS Collaboration, H1 Collaboration, PoS EPS-HEP2011 (2011) 320, arXiv:1112.2107 [hep-ph].
- [5] A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, *Eur. Phys. J. C* 63 (2009) 189, arXiv:0901.0002 [hep-ph].
- [6] R.D. Ball, V. Bertone, S. Carrazza, C.S. Deans, L. Del Debbio, S. Forte, A. Guffanti, N.P. Hartland, et al., *Nucl. Phys. B* 867 (2013) 244, arXiv:1207.1303 [hep-ph].
- [7] X. Ji, J.-H. Zhang, Y. Zhao, *Phys. Rev. Lett.* 111 (11) (2013) 112002, arXiv:1304.6708 [hep-ph].
- [8] X. Ji, *Phys. Rev. Lett.* 110 (26) (2013) 262002, arXiv:1305.1539 [hep-ph].
- [9] X. Xiong, X. Ji, J.H. Zhang, Y. Zhao, *Phys. Rev. D* 90 (2014) 014051, arXiv:1310.7471 [hep-ph].
- [10] H.-W. Lin, J.-W. Chen, S.D. Cohen, X. Ji, arXiv:1402.1462 [hep-ph].
- [11] Y. Hatta, X. Ji, Y. Zhao, *Phys. Rev. D* 89 (2014) 085030, arXiv:1310.4263 [hep-ph].
- [12] Y.-Q. Ma, J.-W. Qiu, arXiv:1404.6860 [hep-ph].
- [13] X. Ji, *Sci. China, Phys. Mech. Astron.* 57 (7) (2014) 1407, arXiv:1404.6680 [hep-ph].
- [14] X. Ji, P. Sun, X. Xiong, F. Yuan, arXiv:1405.7640 [hep-ph].
- [15] X. Ji, J.H. Zhang, Y. Zhao, arXiv:1409.6329 [hep-ph].
- [16] C. Alexandrou, K. Cichy, V. Drach, E. Garcia-Ramos, K. Hadjiyiannakou, K. Jansen, F. Steffens, C. Wiese, *Phys. Rev. D* 92 (2015) 014502, <http://dx.doi.org/10.1103/PhysRevD.92.014502>, arXiv:1504.07455 [hep-lat].
- [17] X. Ji, J.H. Zhang, *Phys. Rev. D* 92 (2015) 034006, <http://dx.doi.org/10.1103/PhysRevD.92.034006>, arXiv:1505.07699 [hep-ph].
- [18] X. Ji, A. Schäfer, X. Xiong, J.H. Zhang, *Phys. Rev. D* 92 (2015) 014039, <http://dx.doi.org/10.1103/PhysRevD.92.014039>, arXiv:1506.00248 [hep-ph].
- [19] X. Xiong, J.H. Zhang, *Phys. Rev. D* 92 (5) (2015) 054037, <http://dx.doi.org/10.1103/PhysRevD.92.054037>, arXiv:1509.08016 [hep-ph].
- [20] H.n. Li, arXiv:1602.07575 [hep-ph].
- [21] J.W. Chen, S.D. Cohen, X. Ji, H.W. Lin, J.H. Zhang, *Nucl. Phys. B* 911 (2016) 246, <http://dx.doi.org/10.1016/j.nuclphysb.2016.07.033>, arXiv:1603.06664 [hep-ph].
- [22] Z. Davoudi, M.J. Savage, *Phys. Rev. D* 86 (2012) 054505, <http://dx.doi.org/10.1103/PhysRevD.86.054505>, arXiv:1204.4146 [hep-lat].
- [23] W. Detmold, C.J.D. Lin, *Phys. Rev. D* 73 (2006) 014501, <http://dx.doi.org/10.1103/PhysRevD.73.014501>, arXiv:hep-lat/0507007.

- [24] K.F. Liu, S.J. Dong, Phys. Rev. Lett. 72 (1994) 1790, <http://dx.doi.org/10.1103/PhysRevLett.72.1790>, arXiv:hep-ph/9306299.
- [25] K.F. Liu, S.J. Dong, T. Draper, D. Leinweber, J.H. Sloan, W. Wilcox, R.M. Woloshyn, Phys. Rev. D 59 (1999) 112001, <http://dx.doi.org/10.1103/PhysRevD.59.112001>, arXiv:hep-ph/9806491.
- [26] K.F. Liu, Phys. Rev. D 62 (2000) 074501, <http://dx.doi.org/10.1103/PhysRevD.62.074501>, arXiv:hep-ph/9910306.
- [27] K.F. Liu, arXiv:1603.07352 [hep-ph].
- [28] V. Braun, D. Mueller, Eur. Phys. J. C 55 (2008) 349, <http://dx.doi.org/10.1140/epjc/s10052-008-0608-4>, arXiv:0709.1348 [hep-ph].
- [29] V.S. Dotsenko, S.N. Vergeles, Nucl. Phys. B 169 (1980) 527, [http://dx.doi.org/10.1016/0550-3213\(80\)90103-0](http://dx.doi.org/10.1016/0550-3213(80)90103-0).
- [30] H. Dorn, Fortschr. Phys. 34 (1986) 11, <http://dx.doi.org/10.1002/prop.19860340104>.
- [31] N.S. Craigie, H. Dorn, Nucl. Phys. B 185 (1981) 204, [http://dx.doi.org/10.1016/0550-3213\(81\)90372-2](http://dx.doi.org/10.1016/0550-3213(81)90372-2).
- [32] B.U. Musch, P. Hagler, J.W. Negele, A. Schafer, Phys. Rev. D 83 (2011) 094507, <http://dx.doi.org/10.1103/PhysRevD.83.094507>, arXiv:1011.1213 [hep-lat].
- [33] T. Ishikawa, Y.Q. Ma, J.W. Qiu, S. Yoshida, arXiv:1609.02018 [hep-lat].