



PAPER • OPEN ACCESS

Towards a quantum time mirror for non-relativistic wave packets

To cite this article: Phillipp Reck *et al* 2018 *New J. Phys.* **20** 033013

View the [article online](#) for updates and enhancements.

Related content

- [Spatial transport of electron quantum states with strong attosecond pulses](#)
M Chovancova, H Agueny, M Førre et al.
- [Effective field theory description of halo nuclei](#)
H-W Hammer, C Ji and D R Phillips
- [A simple formula to predict the influence of the near-field in the optical control of confined electron systems](#)
Takashi Takeuchi, Shinichiro Ohnuki and Tokuei Sako



PAPER

Towards a quantum time mirror for non-relativistic wave packets

Phillipp Reck¹, Cosimo Gorini¹, Arseni Goussev², Viktor Krueckl¹, Mathias Fink³ and Klaus Richter¹¹ Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany² Department of Mathematics, Physics and Electrical Engineering, Northumbria University, Newcastle upon Tyne, NE1 8ST, United Kingdom³ Institut Langevin, ESPCI, CNRS, PSL Research University, 1 rue Jussieu, F-75005, Paris, FranceE-mail: klaus.richter@ur.de**Keywords:** matter waves, quantum wave packets, quantum time mirrors, quantum scattering, nonlinear Schrödinger equation

RECEIVED

21 November 2017

REVISED

9 February 2018

ACCEPTED FOR PUBLICATION

12 February 2018

PUBLISHED

26 March 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Abstract

We propose a method—a quantum time mirror (QTM)—for simulating a partial time-reversal of the free-space motion of a non-relativistic quantum wave packet. The method is based on a short-time spatially homogeneous perturbation to the wave packet dynamics, achieved by adding a nonlinear time-dependent term to the underlying Schrödinger equation. Numerical calculations, supporting our analytical considerations, demonstrate the effectiveness of the proposed QTM for generating a time-reversed echo image of initially localized matter-wave packets in one and two spatial dimensions. We also discuss possible experimental realizations of the proposed QTM.

1. Introduction

The question of how to invert the time evolution of a wave, classical or quantum, in an efficient and controllable way has both fundamental and practical importance. The fundamental aspect of the question is evident from its connection with the problem of unidirectionality of the arrow of time, conceived in a seminal 19th century debate between Loschmidt and Boltzmann [1, 2]. The practical importance is apparent from numerous applications in medicine, telecommunication, material analysis, and, more generally, wave control [3–7].

One fruitful approach to the time inversion of classical wave motion is based on the concept of a time-reversal mirror: an array of receiver-emitter antennas is used to first record an incident wavefront, originating say from a localized source, and then to rebroadcast a time-inverted copy of the recording, thus generating a wave that effectively propagates backward in time and refocuses at the source point. To date, time reversal mirrors have been successfully implemented with acoustic [8, 9], elastic [3], electromagnetic [10], and water waves [11, 12].

The classical procedure at the heart of such time-reversal mirrors, i.e. a continuous measurement and a subsequent reinjection of the signal, cannot be directly applied to quantum systems. The fundamental obstacle here is that any measurement performed on a quantum system is bound to perturb the quantum state and consequently affects its time evolution. (A theoretical scenario in which a time-dependent wave function is measured, recorded and then ‘played back’ by a perfect non-invasive detector-emitter has been analyzed in [13, 14]). An alternative approach to manipulate the propagation of waves relies on non-adiabatic perturbations to the system dynamics, such as an instantaneous change of its boundary conditions [15–22]. Protocols of this kind were considered for time- and space-modulated one-dimensional photonic [23–25] and magnonic crystals [26, 27]. More recently, Bacot *et al* put forward and experimentally realized an instantaneous time mirror for gravity-capillary waves, requiring a sudden but homogenous modulation of water wave celerity [7]. Such approaches bypass the recording procedure, and are thus very appealing for quantum systems. A specific time-reversal protocol, albeit valid in a very narrow momentum range, was indeed devised for a one-dimensional periodically kicked optical lattice [28] and realized in a ⁸⁷Rb Bose–Einstein condensate (BEC) [29]. On the other hand, an instantaneous quantum time mirror (QTM) for Dirac-like systems (exploiting their spinor structure) was recently proposed in [30].

In this paper, we propose and investigate an experimentally realizable method for simulating the time-reversal of the free-space motion, thus mimicking a nonlinear QTM, for a spatially extended (orbital) quantum-mechanical wave function, such as a BEC cloud. The approach relies on generating a short-time spatially uniform perturbation to the wave packet dynamics that corresponds to an additional nonlinear term added in the Schrödinger equation; in a BEC cloud system, such a perturbation can be realized using established experimental techniques allowing to tune the strength of the interaction among the cloud atoms [31, 32]. More precisely, our QTM protocol comprises three stages: (i) a matter-wave packet propagates freely in space for a time $0 < t < t_0$; (ii) at $t = t_0$, a strong nonlinear perturbation is switched on globally for a short period δt , leading to a near-instantaneous modification of the position-dependent phase of the wave function; (iii) the perturbation is switched off, the wave function evolves freely again, and at a time $t_{\text{echo}} > t_0 + \delta t$ an ‘echo’ signal of the original wave packet is observed. From the conceptual viewpoint, the nonlinear QTM proposed in this paper can be viewed as a quantum-mechanical counterpart of the instantaneous time mirror for gravity-capillary waves by Bacot *et al* [7].

The paper is organized as follows. In section 2, we describe the physical principle underlying the proposed QTM. Two concrete scenarios for generating a time-reversed motion of matter waves in one and two spatial dimensions are analyzed in section 3. A summary, concluding remarks, and a discussion of the feasibility of an experimental realization of the proposed QTM are presented in section 4. Technical calculations are deferred to an appendix.

2. Physical principle of a nonlinear QTM

We address the time evolution of a matter-wave packet $\Psi(\mathbf{r}, t)$, subject to the initial condition $\Psi(\mathbf{r}, 0) = \Psi_0(\mathbf{r})$, in accordance with the $(D + 1)$ -dimensional nonlinear Schrödinger equation

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi + \lambda f(t - t_0) |\Psi|^2 \Psi. \quad (1)$$

Here, m is the atomic mass, λ quantifies the nonlinearity strength, and t_0 denotes the time around which the nonlinear term representing interaction effects is switched on. The function $f(\zeta)$ is sharply peaked around $\zeta = 0$ and is chosen to satisfy the normalization condition $\int_{-\infty}^{+\infty} d\zeta f(\zeta) = 1$. We take f to be a δ -function in our analytical calculations and a Gaussian peak $f(\zeta) = \frac{1}{\sqrt{2\pi}\Delta t} e^{-\frac{\zeta^2}{2\Delta t^2}}$ in all numerical simulations. The pulse length $\Delta t \ll t_0$ will be taken as $0.001t_0$ and $0.0025t_0$ in one and two-dimensions, respectively.

Wave packet dynamics in the presence of an infinitesimally short nonlinear kick, $f(\zeta) = \delta(\zeta)$, can be described as follows. Rescaling $t \rightarrow t_0 t$, $\mathbf{r} \rightarrow \sqrt{\frac{\hbar t_0}{m}} \mathbf{r}$, $\Psi \rightarrow \left(\frac{m}{\hbar t_0}\right)^{D/4} \Psi$, and $\lambda \rightarrow \hbar \left(\frac{\hbar t_0}{m}\right)^{D/2} \lambda$, we write the nonlinear Schrödinger equation in a dimensionless form as

$$i \frac{\partial \Psi}{\partial t} = -\frac{1}{2} \nabla^2 \Psi + \lambda \delta(t - 1) |\Psi|^2 \Psi. \quad (2)$$

The evolution of the wave function from $\Psi_0(\mathbf{r})$ at $t = 0$ to its value $\Psi_-(\mathbf{r}) = \Psi(\mathbf{r}, t = 1^-)$ right before the kick is given by

$$\Psi_-(\mathbf{r}) = \int d^D \mathbf{r}' K(\mathbf{r} - \mathbf{r}', t) \Psi_0(\mathbf{r}'), \quad (3)$$

where the integration runs over the infinite D -dimensional space, and $K(\mathbf{q}, t) = (2\pi i t)^{-D/2} \exp[i|\mathbf{q}|^2/(2t)]$ is the free-particle propagator. The nonlinear kick results in an instantaneous change of the wave function from $\Psi_-(\mathbf{r})$ at $t = 1^-$ to

$$\Psi_+(\mathbf{r}) = \Psi_-(\mathbf{r}) e^{-i\lambda |\Psi_-(\mathbf{r})|^2} \quad (4)$$

at $t = 1^+$. Indeed, during the time interval $1^- < t < 1^+$, the wave function transformation is dominated by the second term in the right-hand side of equation (2), and effectively governed by the differential equation $\frac{\partial \ln \Psi(\mathbf{r}, t)}{\partial t} = -i\lambda |\Psi_-(\mathbf{r})|^2 \delta(t - 1)$, the solution of which is given by equation (4). After the kick, the wave function evolves freely, so that $\Psi(\mathbf{r}, t) = \int d^D \mathbf{r}' K(\mathbf{r} - \mathbf{r}', t) \Psi_+(\mathbf{r}')$ for all times $t > 1$.

As evident from equation (4), the instantaneous nonlinear kick alters the phase of the wave function without producing any probability density redistribution, so that $\rho(\mathbf{r}) \equiv |\Psi_+(\mathbf{r})|^2 = |\Psi_-(\mathbf{r})|^2$. The phase change however affects the probability current, whose dimensionless expression reads $\mathbf{j}(\mathbf{r}, t) = \text{Im} [\Psi^*(\mathbf{r}, t) \nabla \Psi(\mathbf{r}, t)]$. A straightforward evaluation of the current right after the kick, $\mathbf{j}_+ = \text{Im} [\Psi_+^* \nabla \Psi_+]$, yields

$$\mathbf{j}_+ = \mathbf{j}_- + \Delta \mathbf{j} \quad \text{with} \quad \Delta \mathbf{j} = -\lambda \rho \nabla \rho, \quad (5)$$

where $\mathbf{j}_- = \text{Im} [\Psi_-^* \nabla \Psi_-]$ is the probability current immediately preceding the kick. This in turn means that, by properly tuning the kicking strength λ , the wave propagation direction can be reversed for those parts of the

matter wave for which the vector $\nabla\rho$ is aligned (or anti-aligned) with \mathbf{j}_- . Below we show that in geometries accessible in atom-optics experiments this reversal effect is robust and well-pronounced.

3. The QTM at work

As our first example, we consider the case of the initial state given by a 1D Gaussian wave packet,

$$\Psi_0(x) = (\pi\sigma^2)^{-1/4} \exp\left(-\frac{x^2}{2\sigma^2} + ikx\right), \quad (6)$$

characterized by the dimensionless real spatial dispersion σ and average momentum k . The corresponding wave function at time $t = 1^-$ is obtained from equation (3) and reads (up to a position-independent phase factor) $\Psi_-(x) = [\pi(\sigma^2 + \sigma^{-2})]^{-1/4} \exp[-\xi^2/2(\sigma^2 + i) + ik\xi]$ with $\xi = x - k$ denoting the distance from the wave packet center. Thus, the probability density at $t = 1$ is $\rho = \exp(-\xi^2/\sigma_1^2)/\sqrt{\pi}\sigma_1$, where $\sigma_1 = \sqrt{\sigma^2 + \sigma^{-2}}$ is the dispersion of the wave packet at the time of the kick. The probability currents before and after the kick are, respectively, $j_- = \left(k + \frac{\xi}{\sigma^2\sigma_1^2}\right)\rho$ and $j_+ = j_- + \Delta j$, with $\Delta j = \frac{2\lambda\xi}{\sigma_1^2}\rho^2$. The minimal kick strength λ_{\min} necessary for reversing the direction of motion of (and effectively reflecting) a part of the wave packet can be estimated by requiring $j_+ = 0$ at $\xi = -\sigma_1$. In the case of a fast moving wave packet, such that $k \gg 1/\sigma^2\sigma_1$, this estimation yields

$$\lambda_{\min} \simeq Ck\left(\sigma^2 + \frac{1}{\sigma^2}\right) \quad (7)$$

with $C = e\sqrt{\pi}/2 \simeq 2.4$. Then, given a kicking strength $\lambda > \lambda_{\min}$, the time t_{echo} at which the reflected part of the wave packet reaches its initial position, leading to a partial echo of the original wave packet, can be evaluated as follows. The velocity of the reflected wave is $k_{\text{rev}} \simeq j_+|_{\xi=-\sigma_1} \simeq k - \frac{\lambda}{C\sigma_1^2} = \left(1 - \frac{\lambda}{\lambda_{\min}}\right)k$, and the revival occurs when $|k_{\text{rev}}|(t_{\text{echo}} - 1) = k$ or, correspondingly, at

$$t_{\text{echo}} = \frac{\lambda}{\lambda - \lambda_{\min}}. \quad (8)$$

The numerical simulations are based on the wave packet propagation algorithm, Time-dependent Quantum Transport [33]. The state is discretized on a square grid and the time evolution is calculated for sufficiently small time steps such that the Hamilton operator can be assumed time independent for each step. We calculate the action of H on ψ in a mixed position and momentum-space representation by the application of Fourier Transforms. With this a Krylov Space is spanned, which can be used to calculate the time evolution using a Lanczos method [34].


The echo strength is quantified by the norm correlation between the initial and the time propagated wave packet defined as [35]

$$\mathcal{N}(t) = \frac{\int d^D\mathbf{r} |\Psi_0(\mathbf{r})|^2 |\Psi(\mathbf{r}, t)|^2}{\sqrt{\int d^D\mathbf{r} |\Psi_0(\mathbf{r})|^4 \int d^D\mathbf{r} |\Psi(\mathbf{r}, t)|^4}}. \quad (9)$$

In contrast to other possible measures of the echo strength, such as fidelity, the norm correlation is easily accessible in experiments on quantum systems with continuous degrees of freedom, e.g., atom optics experiments. Figure 1(a) presents $\mathcal{N}(t)$ for various pulse strengths λ at constant σ and k , demonstrating echo strengths up to 60%. The occurring lower peaks at higher λ for larger times are due secondary peaks of the distorted wave packet, which can be seen in figure 1(b): this panel shows the spatial probability density $\rho = |\psi|^2$ at times $t = 0$ (black dashed curve) and $t = 0.99$ (immediately before pulse, black dashed-dotted curve), as well as the reflected wave packets for different λ 's (color code as in panel (a)), each shown at its peak echo time. In the lower plot, the current density j is shown directly after the pulse $t = 1.01$. Parts of the wave packet with negative current density move backwards leading to the echo. For the parameters used, the estimated value for the minimal pulse strength in (7) is $\lambda_{\min} \simeq 20$ corresponding to the red curve, whose current density exhibits only a vanishing negative part that is insufficient for echo generation, thus verifying the prediction (7).

To explore the parameter space for the possibility of achieving echoes, the peak of the norm correlation (in time) is plotted as a function of λ and σ in figure 1(c) and as a function of λ and k in figure 1(d). The black curve shows the analytic approximation (7) of the minimal pulse strength λ_{\min} . Although it does not fit perfectly, the analytic approximation is in good agreement and still well-suited to approximate the minimal pulse strength required for a time-reversal.

In order to better understand the quality, underlying principles and limitations of the proposed QTM, it is instructive to compare the state Ψ_+ , rendered by the nonlinear kick (see equation (4)), against the desired (perfectly time-reversed) state $\mathcal{C}\Psi_-$, obtained by applying the (anti-unitary) complex conjugation operator \mathcal{C} to



the pre-kick state Ψ_- . In the case of the initial state given by equation (6), we have

$$\Psi_+ = \Psi_- e^{-i\phi}, \quad \phi = \frac{\lambda}{\sqrt{\pi} \sigma_1} e^{-(\xi/\sigma_1)^2} \quad (10)$$

and


$$C\Psi_- = \Psi_- e^{-i(\varphi + \varphi_0)}, \quad \varphi = \left(\frac{\xi}{\sigma_1 \sigma} \right)^2 + 2k\xi, \quad (11)$$

where $\sigma_1 = \sqrt{\sigma^2 + \sigma^{-2}}$ is the width of wave packet at time $t = 1$ when the kick occurs, $\xi = x - k$ is the distance measured from the center of the wave packet, and φ_0 is a constant (position-independent) phase related to the global phase of Ψ_- . While, in general, the two phases, ϕ and φ , have different functional forms, $\phi(\xi)$ may serve as a reasonable approximation to $\varphi(\xi)$, modulo a physically irrelevant constant shift, over a finite position interval. It is the probability density supported by this position interval that makes the main contribution to the time-reversed wave generated by the nonlinear QTM. Figure 2 presents a comparison between the ideal (target) phase $\varphi(\xi)$ and the phase $\phi(\xi)$ imprinted by the proposed QTM. The system parameters are taken to be the same as in figure 1(a), i.e. $\sigma = 1$ and $k = 4$, and the three values of the kicking strength considered are $\lambda = 30, 40$, and 50 .

We further investigate the dynamics of a 2D wave packet, initially given by

$$\Psi_0(\mathbf{r}) = \sqrt{\frac{1}{2\pi^{3/2}R\sigma}} \exp\left[-\frac{(r-R)^2}{2\sigma^2} + ikr\right] \quad (12)$$

with $r = |\mathbf{r}|$ and $k > 0$. For $R \gg \sigma$, the wave function is normalized to one and describes a Gaussian ring of radius R and width σ that spreads radially with the average velocity k . A straightforward, although tedious, calculation shows that, in the parametric regime defined by $1 \ll \sigma \ll R$ and $kR \gg 1$, the wave packet at $t = 1^-$



has the form (up to a spatially uniform phase) (see [appendix](#)):

$$\Psi_{-}(\mathbf{r}) \simeq \sqrt{\frac{1}{2\pi^{3/2}R_1\sigma}} \exp\left[-\frac{(r - R_1)^2}{2\sigma^2} + ikr\right], \quad (13)$$

where $R_1 = R + k$ is the radius of the Gaussian ring at time t^- . (Essentially, equation (13) corresponds to a short-time approximation of the ring-shaped wave packet at the kick time, and is valid in the parametric regime in which the dispersive widening of the ring is negligible.) Thus, the corresponding probability density is given by $\rho = (2\pi^{3/2}R_1\sigma)^{-1} \exp[-(r - R_1)^2/\sigma^2]$, and the probability current at $t = t^+$, reads $\mathbf{j}_+ = \left(k\rho + \frac{2\lambda(r - R_1)}{\sigma^2}\rho^2\right)\frac{\mathbf{r}}{r}$. Then, the evaluation of the minimal kick strength required to trigger a probability density echo proceeds in close analogy with the corresponding 1D calculation, resulting in

$$\lambda_{\min} \simeq 2\pi C(R + k)k\sigma^2. \quad (14)$$

Finally, just as in the 1D case, the echo time is determined by equation (8).


The numerical calculations (figure 3) attest the possibility of pronounced echoes up to 90% also in the 2D setup. Although the parameter range is not in the regime of the analytical approximation, the value of λ_{\min} in equation (14) is still well-suited to estimate the minimal pulse strength λ required (see black curves).

The color plots in figures 1 and 3 seem to imply $\lambda > \lambda_{\min}$ (marked as black lines) to be the only echo requirement for echo generation. However for large λ the wave packet splits into many peaks, as shown in figure 1(b), blue curve for $\lambda = 200$. In such a scenario the norm correlation is still fairly high, but the wave packet might not longer have the desired shape. The effect of many peaks, i.e. very large λ , on the norm correlation can be seen in figure 3(c), where the echo strength moderately declines for $\sigma \approx 1$ and $\lambda > 3000$.

4. Summary and conclusions

In summary, we have proposed a protocol for simulating the time-reversed motion of a localized matter-wave packet evolving in free space. Our method is based on making a near-instantaneous spatially homogeneous perturbation to the wave packet dynamics by externally switching on a nonlinear perturbation for a short time interval. The analytical and numerical considerations presented in our paper demonstrate the efficiency of the proposed QTM in one and two spatial dimensions.

We note that the time reversal protocol proposed in this paper could in principle be employed in different physical systems, such as ultracold atomic clouds, optical pulses, or shallow water waves, as long as the system's time evolution is governed by the nonlinear Schrödinger equation. Here, we further explore the possible connection between the numerical simulations reported in this paper and relevant atom-optics experiments. To this end, we provide an estimate for values of the dimensionless parameters σ and k , defined in equations (6) and (12), for the case of ultracold lithium atoms. The mass of a ^7Li atom is $m = 7.016 \text{ u} = 1.165 \times 10^{-26} \text{ kg}$. Taking the wave packet propagation time until the nonlinear kick to be $t_0 = 10 \text{ ms}$, we see that the wave packet



width range of $10\text{--}50\text{ }\mu\text{m}$ corresponds to $1.05 < \sigma < 5.26$, and the mean velocity range of $2\text{--}10\text{ mm s}^{-1}$ corresponds to $2.1 < k < 10.5$. These parameter ranges coincide with the ones considered in this paper, which strongly suggests that the matter wave reversal effects predicted here can be realized in experiments with lithium BECs.

In order to further facilitate experimental realization of the proposed QTM, we make a rough estimate of the scattering length of condensed lithium atoms required to generate a reflected wave. In the one-dimensional case, the (dimensional) kicking strength $\lambda\hbar\sqrt{\hbar t_0/m}/\Delta t$ (see the discussion preceding equation (2)) is approximately equal to $2N\hbar^2 a_s/(ma_{\perp}^2)$, where N is the number of condensed atoms, a_s is the scattering length, Δt is the kick duration (taken to be $\Delta t = 0.001t_0 = 10\text{ }\mu\text{s}$ in our numerical simulations), and a_{\perp} is the linear length scale of the potential confining the atomic motion in the transverse direction [36]. This yields the estimate $a_s = \lambda a_{\perp}^2 \sqrt{mt_0/\hbar}/(2N\Delta t)$. Taking $N = 10^7$, $a_{\perp} = 10\text{ }\mu\text{m}$, and λ in the range $10\text{--}200$ (see, e.g., figure 1), we find the required scattering length to lie in the range $5\text{ nm} \lesssim a_s \lesssim 105\text{ nm}$. While challenging, the suggested parameter values are not impossible to achieve in modern atom-optics experiments, using, for instance, such novel techniques as optical control of Feshbach resonances [37].

Acknowledgments

The authors thank Ilya Arakelyan for useful discussions. AG acknowledges the support of EPSRC Grant No. EP/K024116/1. CG, VK, KR and PR acknowledge support from Deutsche Forschungsgemeinschaft within SFB 689 and GRK 1570.

Appendix. Free spreading of a Gaussian ring wave packet in 2D: Derivation of equation (13)

Let us consider a two-dimensional wave packet, initially (at $t = t_0$) given by

$$\Psi_0(\mathbf{r}) = C \exp\left(-\frac{(r - R)^2}{2\sigma^2} + ik_0(r - R)\right)$$

with $r = |\mathbf{r}| = \sqrt{x^2 + y^2}$, $\sigma \ll R$, $k_0 > 0$ and $C \simeq 1/\sqrt{2\pi^{3/2}R\sigma}$, so that the probability density is normalized to unity. Let $\Psi(\mathbf{r}, t)$ be the wave packet evolved from $\Psi_0(\mathbf{r})$ in the course of a free-particle evolution through time t . Here, we would like to show that in the parametric regime given by $\sqrt{\hbar t/m} \ll \sigma \ll R$ and $k_0 R \gg 1$ the wave packet Ψ has the same functional dependence on \mathbf{r} as Ψ_0 .

The free-particle propagator in 2D reads

$$K_0(\mathbf{r}, \mathbf{r}', t) = \frac{m}{2\pi i \hbar t} \exp\left(i \frac{m(\mathbf{r} - \mathbf{r}')^2}{2\hbar t}\right).$$

Hence,

$$\begin{aligned} \Psi(\mathbf{r}, t) &= \frac{mC}{2\pi i \hbar t} \int d^2\mathbf{r}' \exp\left(-\frac{(r' - R)^2}{2\sigma^2} + ik_0(r' - R) + i \frac{m(\mathbf{r} - \mathbf{r}')^2}{2\hbar t}\right) \\ &= \frac{mC}{i \hbar t} \exp\left(-\frac{R^2}{2\sigma^2} - ik_0 R + i \frac{m r^2}{2\hbar t}\right) G(r, t), \end{aligned}$$

where

$$G(r, t) = \int_0^\infty dr' r' J_0\left(\frac{m r r'}{\hbar t}\right) \exp\left[-\frac{1}{2}\left(\frac{1}{\sigma^2} - i \frac{m}{\hbar t}\right) r'^2 + \left(\frac{R}{\sigma^2} + ik_0\right) r'\right].$$

Let us investigate the behavior of $G(r, t)$ around the spatial point $R + \frac{\hbar k_0}{m} t$. Taking into account the fact that the main contribution to the integral comes from the region $|r' - R| \lesssim \sigma$, we have

$$\frac{m r r'}{\hbar t} \sim \frac{m R}{\hbar t} \left(R + \frac{\hbar k_0}{m} t\right) = \frac{m R^2}{\hbar t} + k_0 R > k_0 R.$$

Assuming $k_0 R \gg 1$, we see that the argument of the Bessel function is always large compared to one, i.e. $\frac{m r r'}{\hbar t} \gg 1$. This allows us to use the large argument asymptotics,

$$J_0\left(\frac{m r r'}{\hbar t}\right) \simeq \sqrt{\frac{\hbar t}{2\pi m r r'}} [e^{i(\frac{m r r'}{\hbar t} - \frac{\pi}{4})} + e^{-i(\frac{m r r'}{\hbar t} - \frac{\pi}{4})}],$$

in order to write

$$\Psi(\mathbf{r}, t) \simeq \frac{C}{i} \sqrt{\frac{m R}{2\pi \hbar t r}} \exp\left(-\frac{R^2}{2\sigma^2} - ik_0 R + i \frac{m r^2}{2\hbar t}\right) [\Phi_+(r, t) + \Phi_-(r, t)].$$

Here,

$$\begin{aligned} \Phi_\pm(r, t) &= e^{\mp i \frac{\pi}{4}} \int_{-\infty}^{+\infty} dr' \sqrt{\frac{r'}{R}} \exp\left\{-\frac{1}{2}\left(\frac{1}{\sigma^2} - i \frac{m}{\hbar t}\right) r'^2 + \left[\frac{R}{\sigma^2} + i\left(k_0 \pm \frac{m r}{\hbar t}\right)\right] r'\right\} \\ &\simeq e^{\mp i \frac{\pi}{4}} \sqrt{\frac{2\pi}{\frac{1}{\sigma^2} - i \frac{m}{\hbar t}}} \exp\left[\frac{\left[\frac{R}{\sigma^2} + i\left(k_0 \pm \frac{m r}{\hbar t}\right)\right]^2}{2\left(\frac{1}{\sigma^2} - i \frac{m}{\hbar t}\right)}\right]. \end{aligned}$$

Introducing

$$\epsilon = \frac{\hbar t}{m \sigma^2} \quad \text{and} \quad v_0 = \frac{\hbar k_0}{m},$$

we rewrite the previous expression as

$$\Phi_\pm(r, t) = e^{\mp i \frac{\pi}{4}} \sqrt{\frac{2\pi i \hbar t}{m(1 + i\epsilon)}} \exp\left(i \frac{m}{2\hbar t} \frac{[\epsilon R + i(v_0 t \pm r)]^2}{1 + i\epsilon}\right).$$

This leads to

$$\begin{aligned}\Psi(\mathbf{r}, t) &= \frac{C}{i} \sqrt{\frac{mR}{2\pi\hbar t r}} \exp\left(\frac{m}{2\hbar t}[-\epsilon R^2 - 2iRv_0 t + ir^2]\right) [\Phi_+(r, t) + \Phi_-(r, t)] \\ &= C \sqrt{\frac{R}{i(1+i\epsilon)r}} \sum_{\gamma=\pm 1} e^{-i\gamma\frac{\pi}{4}} \\ &\quad \times \exp\left(-\frac{m}{2\hbar t} \frac{\epsilon(R+v_0 t + \gamma r)^2 + i2(R+v_0 t + \gamma r)v_0 t - i[(v_0 t)^2 + \epsilon^2(R+\gamma r)^2]}{1+\epsilon^2}\right).\end{aligned}$$

Assuming further that $\epsilon \ll 1$, we have

$$\begin{aligned}\Psi(\mathbf{r}, t) &\simeq C \sqrt{\frac{R}{ir}} \sum_{\gamma=\pm 1} e^{-i\gamma\frac{\pi}{4}} \exp\left\{-\frac{m}{2\hbar t}[\epsilon(R+v_0 t + \gamma r)^2 + i2(R+v_0 t + \gamma r)v_0 t - i(v_0 t)^2]\right\} \\ &= C \sqrt{\frac{R}{ir}} \sum_{\gamma=\pm 1} e^{-i\gamma\frac{\pi}{4}} \exp\left(-\frac{(R+v_0 t + \gamma r)^2}{2\sigma^2} - ik_0(R+v_0 t + \gamma r) + i\frac{\hbar k_0^2 t}{2m}\right).\end{aligned}$$

Taking into account that the last expression for Ψ is only valid for r close to r_t , where

$$r_t = R + v_0 t = R + \frac{\hbar k_0 t}{m},$$

we see that a contribution of the term with $\gamma = +1$ is negligibly small. Thus we arrive at the sought approximation for the freely propagated wave function:

$$\Psi(\mathbf{r}, t) \simeq \sqrt{\frac{1}{2\pi^{3/2}\sigma r_t}} \exp\left(-\frac{(r-r_t)^2}{2\sigma^2} + ik_0(r-r_t) + i\frac{\hbar k_0^2 t}{2m}\right).$$

This expression is only valid in the parametric regime defined by

$$\sqrt{\frac{\hbar t}{m}} \ll \sigma \ll R \quad \text{and} \quad k_0 R \gg 1.$$

References

- [1] Loschmidt J 1876 Über den Zustand des Wärmegleichgewichts eines Systems von Körpern mit Rücksicht auf die Schwerkraft *Sitzungsberichte Akad. Wiss.* **2** 128
- [2] Boltzmann L 1877 Über die Beziehung eines allgemeinen mechanischen Satzes zum zweiten Hauptsatz der Wärmetheorie *Sitzungsberichte Akad. Wiss.* **2** 67
- [3] Fink M 1997 Time reversed acoustics *Phys. Today* **50** 34
- [4] Fink M 1999 Time-reversed acoustics *Sci. Am.* **281** 91
- [5] Lerosee G, de Rosny J, Tourin A and Fink M 2007 Focusing beyond the diffraction limit with far-field time reversal *Science* **315** 1120
- [6] Mosk A P, Lagendijk A, Lerosee G and Fink M 2012 Controlling waves in space and time for imaging and focusing in complex media *Nat. Photon.* **6** 283
- [7] Bacot V, Labousse M, Eddi A, Fink M and Fort E 2016 Time reversal and holography with spacetime transformations *Nat. Phys.* **12** 972
- [8] Fink M 1992 Time reversal of ultrasonic fields: I. Basic principles *IEEE Trans. Ultrason. Ferroelectr. Freq. Control* **39** 555–66
- [9] Draeger C and Fink M 1997 One-channel time reversal of elastic waves in a chaotic 2d-silicon cavity *Phys. Rev. Lett.* **79** 407
- [10] Lerosee G, de Rosny J, Tourin A, Derode A, Montaldo G and Fink M 2004 Time reversal of electromagnetic waves *Phys. Rev. Lett.* **92** 193904
- [11] Prasadka A, Feat S, Petitjeans P, Pagneux V, Maurel A and Fink M 2012 Time reversal of water waves *Phys. Rev. Lett.* **109** 064501
- [12] Chabchoub A and Fink M 2014 Time-reversal generation of rogue waves *Phys. Rev. Lett.* **112** 124101
- [13] Pastawski H M, Danieli E P, Calvo H L and Foa Torres L E F 2007 Towards a time reversal mirror for quantum systems *Europhys. Lett.* **77** 40001
- [14] Calvo H L and Pastawski H M 2010 Exact time-reversal focusing of acoustic and quantum excitations in open cavities: the perfect inverse filter *Europhys. Lett.* **89** 60002
- [15] Moshinsky M 1952 Diffraction in time *Phys. Rev.* **88** 625
- [16] Gerasimov A S and Kazarnovskii M V 1976 Possibility of observing nonstationary quantum-mechanical effects by means of ultracold neutrons *Sov. Phys.—JETP* **44** 892
- [17] Brukner Č and Zeilinger A 1997 Diffraction of matter waves in space and in time *Phys. Rev. A* **56** 3804
- [18] Mendonça J T and Shukla P K 2002 Time refraction and time reflection: two basic concepts *Phys. Scr.* **65** 160
- [19] del Campo A, Garcia-Calderón G and Muga J G 2009 Quantum transients *Phys. Rep.* **476** 1
- [20] Goussev A 2012 Huygens–Fresnel–Kirchhoff construction for quantum propagators with application to diffraction in space and time *Phys. Rev. A* **85** 013626
- [21] Haslinger P, Dorre N, Geyer P, Rodewald J, Nimmrichter S and Arndt M 2013 A universal matter-wave interferometer with optical ionization gratings in the time domain *Nat. Phys.* **9** 144
- [22] Goussev A 2013 Diffraction in time: an exactly solvable model *Phys. Rev. A* **87** 053621
- [23] Sivan Y and Pendry J B 2011 Time reversal in dynamically tuned zero-gap periodic systems *Phys. Rev. Lett.* **106** 193902

- [24] Sivan Y and Pendry J B 2011 Theory of wave-front reversal of short pulses in dynamically tuned zero-gap periodic systems *Phys. Rev. A* **84** 033822
- [25] Sivan Y and Pendry J B 2011 Broadband time-reversal of optical pulses using a switchable photonic-crystal mirror *Opt. Express* **19** 14502
- [26] Chumak A V, Tiberkevich V S, Karenowska A D, Serga A A, Gregg J F, Slavin A N and Hillebrands B 2010 All-linear time reversal by a dynamic artificial crystal *Nat. Commun.* **1** 141
- [27] Karenowska A D, Gregg J F, Tiberkevich V S, Slavin A N, Chumak A V, Serga A A and Hillebrands B 2012 Oscillatory energy exchange between waves coupled by a dynamic artificial crystal *Phys. Rev. Lett.* **108** 015505
- [28] Martin J, Georgeot B and Shepelyansky D L 2008 Cooling by time reversal of atomic matter waves *Phys. Rev. Lett.* **100** 044106
- [29] Ullah A and Hoogerland M D 2011 Experimental observation of Loschmidt time reversal of a quantum chaotic system *Phys. Rev. E* **83** 046218
- [30] Reck P, Gorini C, Goussev A, Krueckl V, Fink M and Richter K 2017 Dirac quantum time mirror *Phys. Rev. B* **95** 165421
- [31] Inouye S, Andrews M R, Stenger J, Miesner H-J, Stamper-Kurn D M and Ketterle W 1998 Observation of Feshbach resonances in a Bose–Einstein condensate *Nature* **392** 151
- [32] Roberts J L, Claussen N R Jr, Burke J P, Greene C H, Cornell E A and Wieman C E 1998 Resonant magnetic field control of elastic scattering in cold ^{85}Rb *Phys. Rev. Lett.* **81** 5109
- [33] Krückl V 2013 Wave packets in mesoscopic systems: from time-dependent dynamics to transport phenomena in graphene and topological insulators *PhD Thesis* Universität Regensburg, Regensburg, Germany (available at: <http://krueckl.de/#en/tqt.php>)
- [34] Lanczos C 1950 An iteration method for the solution of the eigenvalue problem of linear differential and integral operators *J. Res. Natl Bur. Stand.* **45** 255
- [35] Eckhardt B 2003 Echoes in classical dynamical systems *J. Phys. A: Math. Gen.* **36** 371
- [36] Salasnich L, Parola A and Reatto L 2002 Effective wave equations for the dynamics of cigar-shaped and disk-shaped bose condensates *Phys. Rev. A* **65** 043614
- [37] Clark L W, Ha L, Xu C and Chin C 2015 Quantum dynamics with spatiotemporal control of interactions in a stable Bose–Einstein condensate *Phys. Rev. Lett.* **115** 155301