



Linking Personality Traits to Individual Differences in Affective Spaces

Seth M. Levine^{1*}, Aino L. I. Alahäivälä¹, Theresa F. Wechsler², Anja Wackerle¹, Rainer Rupperecht¹ and Jens V. Schwarzbach¹

¹ Department of Psychiatry and Psychotherapy, University of Regensburg, Regensburg, Germany, ² Department of Clinical Psychology and Psychotherapy, University of Regensburg, Regensburg, Germany

OPEN ACCESS

Edited by:

Anthony P. Atkinson,
Durham University, United Kingdom

Reviewed by:

Luca Passamonti,
University of Cambridge,
United Kingdom
Chujun Lin,
California Institute of Technology,
United States

*Correspondence:

Seth M. Levine
seth.levine@ukr.de

Specialty section:

This article was submitted to
Emotion Science,
a section of the journal
Frontiers in Psychology

Received: 11 December 2019

Accepted: 26 February 2020

Published: 12 March 2020

Citation:

Levine SM, Alahäivälä ALI, Wechsler TF, Wackerle A, Rupperecht R and Schwarzbach JV (2020) Linking Personality Traits to Individual Differences in Affective Spaces. *Front. Psychol.* 11:448. doi: 10.3389/fpsyg.2020.00448

Different individuals respond differently to emotional stimuli in their environment. Therefore, to understand how emotions are represented mentally will ultimately require investigations into individual-level information. Here we tasked participants with freely arranging emotionally charged images on a computer screen according to their subjective emotional similarity (yielding a unique affective space for each participant) and subsequently sought external validity of the layout of the individuals' affective spaces through the five-factor personality model (Neuroticism, Extraversion, Openness to Experience, Agreeableness, Conscientiousness) assessed via the NEO Five-Factor Inventory. Applying agglomerative hierarchical clustering to the group-level affective space revealed a set of underlying affective clusters whose within-cluster dissimilarity, per individual, was then correlated with individuals' personality scores. These cluster-based analyses predominantly revealed that the dispersion of the negative cluster showed a positive relationship with Neuroticism and a negative relationship with Conscientiousness, a finding that would be predicted by prior work. Such results demonstrate the non-spurious structure of individualized emotion information revealed by data-driven analyses of a behavioral task (and validated by incorporating psychological measures of personality) and corroborate prior knowledge of the interaction between affect and personality. Future investigations can similarly combine hypothesis- and data-driven methods to extend such findings, potentially yielding new perspectives on underlying cognitive processes, disease susceptibility, or even diagnostic/prognostic markers for mental disorders involving emotion dysregulation.

Keywords: affective science, Big Five, clustering, emotions, individual differences, personality

INTRODUCTION

Emotions are an aspect of everyday life, and questions regarding their elusive nature have been of interest for millennia. While recent discussions (Hamann, 2012; Lindquist et al., 2012; Adolphs, 2017; Barrett, 2017) and neuroscientific investigations (Grimm et al., 2006; Nielen et al., 2009; Baucom et al., 2012; Goodkind et al., 2012; Chikazoe et al., 2014; Shinkareva et al., 2014; Kragel et al., 2016; Saarimäki et al., 2016) have advanced the field of affective (neuro)science, there still remains no consensus concerning the inherent features of emotions (Ekman, 2016). Alongside this

general interest in understanding affective functionality is the clinical interest in understanding affective dysfunctionality (Gross and Jazaieri, 2014), as emotion dysregulation has been associated with various mental disorders (Mennin et al., 2005; Reimherr et al., 2005; Etkin and Wager, 2007; Amstadter, 2008; Taylor et al., 2012; Carpenter and Trull, 2013). Following the drive for improved translational research in psychiatry (Machado-Vieira, 2012; Knüppel et al., 2013) and personalized medicine (Hamburg and Collins, 2010), we sought to study individual differences in affective information representations by combining data-driven analyses of a behavioral experiment with psychological measures of personality traits, given the well-established link between emotions and personality (Costa and McCrae, 1980; Gross et al., 1998; Kokkonen and Pulkkinen, 2001; Ng and Diener, 2009).

To this end, we employed the multi-arrangement method (Goldstone, 1994) and inverse multidimensional scaling (Kriegeskorte and Mur, 2012), which have recently been utilized in the domain of cognitive neuroscience (Mur et al., 2013; Charest et al., 2014; Bracci et al., 2016; Levine et al., 2018c), to emotionally charged stimuli in order to discern individualized structures (that reflect mental representations) of affective information. Participants freely arranged the stimuli according to their subjective emotional similarity in a continuous space, which resulted in a label-free, unique representation of an “affective space” for each participant. Other recent work has investigated the organizing principles of affect/emotions using psychological (Nummenmaa et al., 2014, 2018; Koch et al., 2016; Cowen and Keltner, 2017; van Tilburg and Igou, 2017) and neuroscientific (Kragel and LaBar, 2015; Skerry and Saxe, 2015; Saarimäki et al., 2018) methods; here we took an approach that focused on how individuals differ in terms of underlying properties of their affective spaces. Being able to distinguish individuals based on affective information (Hamann and Canli, 2004) – and subsequently determine normal variability using a combination of hypothesis-driven and data-driven methods – may offer new routes for social psychology/neuroscience to inform the clinical realm (Cacioppo et al., 2014).

The current study employed data-driven hierarchical clustering to identify clusters of stimuli underlying the affective space. However, to avoid blindly applying unsupervised machine learning methods to the high-dimensional affective spaces, we externally validated the individual differences in affective clustering with differences in personality traits from the five-factor model [“Big Five”: Neuroticism, Extraversion, Openness to Experience, Agreeableness, and Conscientiousness (McCrae and Costa, 1985)], assessed by the NEO-Five Factor Inventory [NEO-FFI (Borkenau and Ostendorf, 2008)]. As prior neuroimaging work has shown that the layout of a representational space can be meaningfully altered by attentional or task-related processes (Brouwer and Heeger, 2013; Nastase et al., 2017) and recent psychological work has shown that clusters within an affective space map onto certain aspects of emotion information (Nummenmaa et al., 2018), we specifically sought to determine whether the *within-cluster distance* of individuals’ affective clusters [which can be seen as the cohesiveness of the concept(s) underlying the pertinent cluster (Iordan et al., 2015)] was linked

to participants’ personality traits. The idea of investigating emotions as they relate to personality follows from decades of evidence that personality is tied to internal or attentional biases of emotion-related processing (Richards et al., 1992; Derryberry and Reed, 1994; Most et al., 2006; Thomsen et al., 2014). Moreover, classic theories of personality would predict that Neuroticism associates with negative information and Extraversion associates with positive information (Eysenck, 1967; Costa and McCrae, 1980; Larsen and Ketelaar, 1989; Rusting and Larsen, 1997). Exploring the relationship between personality and emotional similarity (here, operationalized as cluster dispersion within affective spaces) will allow researchers to investigate how underlying cognitive processes, such as attention or executive control, may interact with personality traits to drive healthy and unhealthy behavior, and whether such interactions are tied to specific brain regions/networks. The findings we present corroborate and extend current knowledge regarding the relationship between affect and personality, validate the non-spurious structure of individualized affective spaces revealed by data-driven methods, and open the door for future research to translate such data-driven paradigms to the clinical domain.

MATERIALS AND METHODS

Participants

One hundred one participants (36 males, 65 females; mean ($\pm \sigma$) age = 24.2 (± 2.59) years) were recruited from the local community via information posters. Participants reported neither a current diagnosis of neurological or mental disorders nor the intake of any psychotropic medication, provided written informed consent before taking part in the study, and were monetarily compensated for their time after completing the experiment. As filling out the NEO-FFI questionnaire was a follow-up procedure to the behavioral experiment, only participants who had initially agreed to be contacted for future studies were asked for their participation in completing the NEO-FFI questionnaire. Of 77 participants contacted, 58 participants (15 males, 43 females; mean ($\pm \sigma$) age = 25.4 (± 2.75) years) ultimately completed the questionnaire, five of whom were compensated with bookstore gift cards following a raffle. All experimental procedures complied with the Declaration of Helsinki and were approved by the local ethics committee at the University of Regensburg.

Apparatus

The experiment was conducted on a 27” Apple iMac using MATLAB R2015b (The MathWorks, Natick, MA, United States). The MATLAB code used to run the experiment was adapted from that described by Kriegeskorte and Mur (2012). All analyses were carried out in MATLAB and SPSS ver. 25 (IBM Corp., Armonk, NY, United States).

Stimuli


Images used in the experiment were obtained from the International Affective Picture System (IAPS) database

(Lang et al., 2008), which contains images with standardized valence and arousal ratings that can be roughly divided into nine categories: animals, people, nature, food, household items, erotic images, accidents, violence, and war. We pseudo-randomly chose six images per category for the experiment, yielding a stimulus set comprising 54 images. If the computer algorithm chose two images that were very visually similar (e.g., two pictures of a dog in a similar position, or two pictures of knives on a table), we manually selected a replacement image with the same (or very similar) valence and arousal values from the corresponding category (see **Supplementary Table 1** for the corresponding image ID's and their respective valence and arousal ratings). Creating the stimulus set was ultimately a trade-off between improved sampling of the emotion space and total experiment duration, as the distance between each pair of images must be computed; the total number of pairwise distances in our case is given by the binomial coefficient $54 \text{ choose } 2$ (i.e., 1,431), which yielded a total experiment duration of, on average, slightly more than one hour per participant.

Trial Protocol

The experiment followed the protocol laid out by Kriegeskorte and Mur (2012), in which participants arrange images on a two-dimensional circular “arena” on a computer screen according to a particular organizing principle. We asked participants to arrange pictures according to how they felt when they viewed each image. That is, images evoking similar emotions should be placed closer together while images evoking dissimilar emotions should be placed further apart; thus the distance between images reflects their relative emotional dissimilarity for the participant.

No more than ten images were shown during a given trial to improve visibility of the pictures (i.e., fitting 54 pictures on the screen at the same time would render them all so small that their content would become unrecognizable). Additionally, we implemented a zoom function, which allowed participants to see the fine-grained detail of each picture. When the participant finished organizing the images, the next trial began with another set of images, for which there remained the least amount of evidence regarding their relative distances to the other images [the lift-the-weakest algorithm (Kriegeskorte and Mur, 2012)]. Thus, some of the images may have been present in two or more successive trials, and these particular images shown on a given trial differed between individuals, depending on how they organized the images throughout the course of the experiment. After distances for all 1,431 pairwise dissimilarities were obtained, inverse multidimensional scaling transformed the two-dimensional distances on the computer screen into a 54×54 dissimilarity matrix (DSM; **Figure 1A**), which represents the high-dimensional dissimilarity structure of the item arrangements (Kriegeskorte and Mur, 2012). However, because the DSM is symmetric across the main diagonal, each participant can be represented by a 1,431-element vector, obtained by vectorizing either the upper or lower triangle of the DSM. We will hereinafter refer to the vector representation of a DSM as a dissimilarity vector (DSV).



NEO-FFI

We used the German version of the NEO-FFI (Borkenau and Ostendorf, 2008) to measure the Big Five personality factors Neuroticism, Extraversion, Openness to Experience, Agreeableness, and Conscientiousness. For each participant, the sum scores for all five factors were assessed and then transferred into a gender- and age-related T-score based on the reference samples reported by Borkenau and Ostendorf (2008), which expresses the manifestation of a Big Five personality factor in an

individual with respect to the manifestation in a gender- and age-corresponding population. For a list of the resulting NEO-FFI scores and their descriptive statistics, see **Supplementary Table 2**.

Data Analysis

Hierarchical Clustering

Because all participants' affective spaces clustered slightly differently, we wanted to ensure that we were comparing clusters across participants based on the same stimuli. To this end, we first scaled each participant's DSV to a range of [0–1] by dividing each DSV by its maximum dissimilarity value. We then computed the median of all 101 scaled DSVs and carried out agglomerative hierarchical clustering using complete linkage (Lance and Williams, 1967) on the group-median DSV (**Figure 1B** and **Supplementary Figure 1**).

We validated the stability of different numbers of clusters underlying the group-median affective space by iteratively cutting the resulting dendrogram at different heights and calculating the silhouette index (Rousseeuw, 1987). This index measures how well a given data point is assigned to a particular cluster (i.e., within-cluster distance) compared to the nearest-neighbor cluster (i.e., between-cluster distance) on a scale from -1 to 1 , with higher values indicating a more appropriate clustering solution (n.b., group-median clusters produced higher silhouette values than group-mean clusters). This procedure yielded a 2-cluster and 4-cluster solution (**Figures 1C–E**), which we then mapped back onto each image pair from the participants' individual DSVs, allowing us to calculate, for each participant, the median (rather than mean, as the distance data tended to be skewed) within-cluster distances (which we normalized by the number of elements in the respective cluster). Formally, the median within-cluster distance D for cluster c was calculated as $\frac{\text{median}(D_c)}{|c|}$, where $|c|$ denotes the cardinality of cluster c .

Following classical multidimensional scaling of the group-median affective space (see **Figure 2** for ease of data visualization only, as the actual cluster distances were computed directly from participants' DSVs), the first dimension (variance explained (VE) = $\sim 48\%$) corresponded to the IAPS valence values (Pearson's $r = 0.94$, $p = 8.5 \times 10^{-25}$), when partialing out the arousal values; as such, we assigned the labels of “positive” and “negative” to the resulting clusters from the 2-cluster solution. To a far lesser degree, the second and third dimensions corresponded to the IAPS arousal values ($r = 0.38$, $p = 0.005$, VE = $\sim 8\%$; $r = 0.50$, $p = 1.5 \times 10^{-4}$, VE = $\sim 5\%$, respectively), while partialing out the valence values. To aid in interpreting the results of the subsequent multiple regression (see next section), we ascribed the following descriptive labels to the four clusters based on the general contents of the clusters: “erotic” (positive sub-cluster), “fear/violence” (negative sub-cluster), “medical” (negative sub-cluster), “nature/people/sports” (positive sub-cluster).

Multiple Linear Regression

For each affective cluster (in each clustering solution), we carried out multiple regression in order to assess the degree to which the Big Five personality factors predicted the median dispersion of the affective clusters. To determine, in an unbiased manner, which regressors were in the optimal model, we took


a combinatorial approach to multiple regression. Specifically, given the five factors (Neuroticism, Extraversion, Openness, Agreeableness, Conscientiousness), there were $2^5 - 1$ possible models (i.e., $\sum_{k=1}^5 \binom{5}{k}$), including only the linear terms), each containing a different combination of the five factors as predictors. To find the optimal model, we performed the regression analysis with each of the 31 models and selected the model with the lowest Bayesian Information Criterion (BIC) (Schwarz, 1978). For each model, the cluster dispersion values and the personality scores were z-scored (using their respective sample means and sample standard deviations) to center the data, and a constant predictor was included. To account for having tested 31 models, we further carried out two Monte Carlo procedures to non-parametrically determine whether the optimal model outperformed models applied to randomized median cluster distances. First, we tested all 31 models on the randomized data and stored the minimum BIC (regardless of the model that yielded this BIC) to control for the inflated family-wise error rate (FWER) from having tested multiple models. Repeating this procedure 1,000 times (per affective cluster) provided us with a null distribution of 1,000 minimum BIC values, from which we calculated empirical p -values for the observed optimal BIC (denoted in the results as $p_{FWER}(\text{all})$: for a comparison against all models when all models were fed random data). Second, we also stored the BIC resulting from the *observed optimal model* when this same model was applied to randomized data 1,000 times. This way, we obtained two empirically derived p -values for our optimal model (one from a minimum-BIC null distribution and one from a same-model-BIC null distribution [denoted in the results as $p(\text{same})$: for a comparison against the same model when fed random data]), which indicate how likely it is to find that the optimal model produces a particular BIC under the null hypotheses of no relationship between the Big Five personality factors and the median cluster distances.

RESULTS

To determine whether the structure of the affective spaces was related to personality dimensions, we sought to predict the individualized median intra-cluster distance (for different affective clusters) from the Big Five personality factors using multiple linear regression. Below are the results from the best-performing regression models for the affective clusters underlying both the 2- and the 4-cluster solutions (see also **Supplementary Table 4**).

Two-Cluster Solution: Larger Negative Cluster Dispersion Corresponds to Higher Neuroticism

Starting with the 2-cluster solution from the hierarchical clustering analysis, the optimal model to predict the negative cluster's dispersion contained the Neuroticism scores as its sole regressor ($\beta = 0.34$, SE = 0.126, $t_{56} = 2.73$, $p = 0.0084$), indicating that a higher Neuroticism score tended to predict a larger cluster



dispersion. Although this model marginally failed to surpass the statistical threshold when compared to all other models in the Monte Carlo procedure [$p_{FWER}(\text{all}) = 0.056$], it did perform better than chance when fed real data compared to randomized data [$p(\text{same}) = 0.01$]. With respect to the positive cluster's dispersion, while the optimal model contained the Agreeableness scores as its sole regressor ($\beta = -0.22$, $SE = 0.130$, $t_{56} = -1.65$, $p = 0.1036$), this model did not perform better than chance would predict [$p_{FWER}(\text{all}) = 0.383$, $p(\text{same}) = 0.11$].


Four-Cluster Solution: Opposite Dispersion Patterns of the Fear/Violence Cluster for Neuroticism and Conscientiousness

With the group-level affective space split into four clusters, the previously negative cluster further subdivided into a cluster whose images encompassed concepts of “fear/violence” and a cluster whose images encompassed “medical” concepts (**Figure 2**). Our combinatorial multiple regression procedure revealed that the optimal model contained Neuroticism ($\beta = 0.280$, $SE = 0.127$, $t_{55} = 2.20$, $p = 0.032$) and Conscientiousness ($\beta = -0.289$, $SE = 0.127$, $t_{55} = -2.28$, $p = 0.0266$) as the two regressors to predict the fear/violence cluster dispersion [$p_{FWER}(\text{all}) = 0.014$, $p(\text{same}) = 0.0009$; see **Supplementary Table 3** for a list of all models that survived the statistical threshold]. Similarly to the results from the 2-cluster analysis, Neuroticism scores tended to increase with increasing cluster dispersion (Pearson's $r = 0.381$). Conscientiousness scores, however, showed the opposite pattern in that they tended

to decrease as cluster dispersion increased (Pearson's $r = -0.388$; see **Figure 3** for a visual depiction of this effect). Additionally, the optimal model to predict the medical cluster's dispersion contained only Neuroticism as the regressor ($\beta = 0.207$, $SE = 0.131$, $t_{56} = 1.58$, $p = 0.12$), but this model did not perform better than chance would predict [$p_{FWER}(\text{all}) = 0.424$, $p(\text{same}) = 0.11$].

Four-Cluster Solution: Increasing Erotic Cluster Dispersion Predicted by Decreasing Openness, Agreeableness, and Conscientiousness

Just as the negative cluster from the 2-cluster solution was further subdivided in the 4-cluster solution, so was the positive cluster, which yielded an “erotic” cluster and a generic positive cluster (labeled as “nature/people/sports”) containing the remaining images that included mainly images related to sports, people, nature, etc. Interestingly, the optimal model to predict dispersion of the erotic cluster was a three-regressor model comprising Openness ($\beta = -0.280$, $SE = 0.120$, $t_{54} = -2.34$, $p = 0.023$; Pearson's $r = -0.253$), Agreeableness ($\beta = -0.291$, $SE = 0.12$, $t_{54} = -2.43$, $p = 0.019$; Pearson's $r = -0.258$), and Conscientiousness ($\beta = -0.314$, $SE = 0.120$, $t_{54} = -2.61$, $p = 0.012$; Pearson's $r = -0.266$). Although this model also marginally failed to surpass the statistical threshold when compared to all other models in the Monte Carlo procedure [$p_{FWER}(\text{all}) = 0.06$], it did perform better than chance would predict when fed real data compared to randomized data [$p(\text{same}) = 0.006$]. Regarding the generic positive cluster of nature/people/sports images, similar



to the 2-cluster solution, the optimal model contained the sole regressor of Agreeableness ($\beta = -0.161$, $SE = 0.132$, $t_{56} = -1.22$, $p = 0.228$), but, once again, this model did not outperform what one would expect from randomness alone [$p_{FWER}(\text{all}) = 0.72$, $p(\text{same}) = 0.24$].

DISCUSSION

In order to explore the underlying structure of individuals' affective spaces, we conducted an experiment in which participants organized emotionally charged images according to their individualized emotional similarity and combined data-driven analyses with psychological measures of personality traits. With this combination of methods, we specifically sought associations between personality traits of the five-factor model (“Big Five”) and the dispersion of affective clusters [which reflect the cohesiveness of the concept(s) underlying the pertinent cluster (Jordan et al., 2015)] that populated individuals' affective spaces. Importantly, this approach demonstrated that the clustering of individuals' affective spaces, obtained through unsupervised machine learning methods, is not spurious, as portions of it can be externally validated with information from individuals' personality traits.

The primary result that survived our statistical controls came from the 4-cluster solution. Multiple regression revealed that increasing dispersion of the fear/violence cluster was linked to increasing Neuroticism and decreasing Conscientiousness (see also **Supplementary Table 5**). Such an opposite relationship between these two personality dimensions is not entirely unexpected, as it has been reported as a general finding when assessing the Big Five personality factors (Costa et al., 1991), linked to mental disorders (Trull and Sher, 1994; Kotov et al., 2010), and tied to such domains as social media activity (Liu et al., 2016), emotional problems in adolescents (Smith et al., 2017), and even physiological mechanisms of inflammation (Sutin et al., 2010). Regarding Neuroticism, it is possible that higher Neuroticism coincides with a greater differentiation in the processing of negative stimuli, thereby leading to a finer-grained categorization (i.e., a more dispersed clustering) of negative information. This notion is backed by higher-Neuroticism individuals exhibiting increased processing of unpleasant information (Gomez et al., 2002; Chan et al., 2007) and tending to describe themselves as anxious (McCrae et al., 1986), as anxious individuals selectively attend to negative stimuli (Broadbent and Broadbent, 1988; MacLeod and Mathews, 1988). Regarding Conscientiousness, higher-Conscientiousness individuals have been described as rigorous and orderly (Costa and McCrae, 1992) and have demonstrated greater emotional control when recovering from negative stimuli (Javaraš et al., 2012). Relatedly, increased clustering and efficiency of the frontoparietal network (Toschi et al., 2018), which is considered a critical brain network for cognitive control (Miller and Cohen, 2001), has been associated with higher Conscientiousness. As such, higher-Conscientiousness individuals may have more control over (or a greater need for) compartmentalizing negative information and appropriately managing their resulting behavior. Given that the layout of information in individuals' representational spaces can be altered by processes such as attention (Nastase et al., 2017), typicality judgments (Jordan et al., 2016), and aversive-learning (Dunsmoor et al., 2014; Levine et al., 2018b), it is possible that these personality traits interact with (or even regulate the effectiveness of) such processes to determine how negative information populates individuals' representational spaces. Thus, the correspondence between personality traits and the degree of differentiation of negative information can be explored as, for example, a biomarker for susceptibility to particular disorders or a further method of classifying disorders involving negative affect.

Our findings also have implications for social and affective neuroscience studies. Recent functional magnetic resonance imaging investigations have employed multivariate pattern analysis (Haxby et al., 2001) in an effort to understand how brain regions or networks represent information related to emotion processing (Bush et al., 2018; Kryklywy et al., 2018; Levine et al., 2018c; Saarimäki et al., 2018). Findings like those we present here can inform such studies by underscoring how the heterogeneity of personality information can influence the distribution of information in affective spaces. As a result, this non-homogeneous affective information across individuals may

impact sensitivity in neuroimaging studies of affect, especially those based on longitudinal or pre-post designs, which need to take affective variability into account (Dauvier et al., 2019). This notion is only emphasized by the recent studies that have utilized data from the Human Connectome Project (Van Essen et al., 2013) to link individuals' personality data to resting-state functional networks (Dubois et al., 2018; Mulders et al., 2018; Nostro et al., 2018; Toschi et al., 2018; Passamonti et al., 2019), as certain aspects of the functions underlying these networks may also vary with differences in personality. In some cases, such heterogeneity (in both healthy and clinical populations) may be exactly the variable of interest when translating findings and methods from cognitive neuroscience to (precision) psychiatry (Feczko et al., 2019).

In terms of limitations, it is possible that specific facets of the personality dimensions underlie our results. However, we cannot accurately relate facet-level information to particular aspects of individuals' affective spaces because we administered the NEO-FFI rather than the NEO-PI-R (Costa and McCrae, 1992), which is a longer five-factor model personality questionnaire allowing researchers to investigate the six distinct facets of each of the personality factors. Our approach is therefore less sensitive to exploring facet-level information, which could be a critical component in further specifying the value of such findings. Thus, future investigations following this direction could make use of more detailed questionnaires. An additional, generic drawback is that the regression analyses reported here come from only 58 participants (~57% of the total sample), as we did not have questionnaire data from all participants. This reduction in the sample size consequently reduced the statistical power of our analyses (*cf.* **Supplementary Figure 2**).

CONCLUSION

In conclusion, we carried out a behavioral experiment based on label-free, subjective, emotional similarity judgments and combined data-driven analyses with the classical Big Five personality traits to investigate cluster dispersion in individualized affective spaces. Our combination of methods revealed, primarily, a relationship between how individuals tended to judge the emotional distance of negatively charged stimuli and the personality dimensions of Neuroticism and Conscientiousness. These findings demonstrate that there is non-spurious structure underlying affective spaces revealed with unsupervised machine learning methods. Such assumption-free techniques may help bridge affective sciences with the clinical domain by providing objective measures of normal affective variability across individuals and characterizing how such variability may interact with personality traits and relate to mental conditions. Future investigations can scrutinize such

REFERENCES

Adolphs, R. (2017). How should neuroscience study emotions? By distinguishing emotion states, concepts, and experiences. *Soc. Cogn. Affect. Neurosci.* 12, 24–31. doi: 10.1093/scan/nsw153

findings, for example, for their longitudinal value in identifying subtypes of particular patient populations. Such directions offer clinical psychology and psychiatry the opportunity to adapt psychological and neuroscientific methods to develop new markers for critical issues such as disease susceptibility and treatment response.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, to any qualified researcher upon reasonable request. A subset of the data presented in this manuscript were previously made public in a preprint at PsyArXiv (Levine et al., 2018a).

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Ethics Committee at the University of Regensburg. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

SL and JS designed the study. RR contributed to the experiment resources. SL, AA, and AW acquired the data. SL and TW analyzed the data. SL, AA, and TW drafted the manuscript. All authors revised and approved the final version of the manuscript.

FUNDING

AA was supported by a scholarship from the Osk. Huttunen Foundation.

ACKNOWLEDGMENTS

We would like to thank Angelika Lingnau for contributing experimental code for data acquisition and Viola Wagner for assistance with data analysis.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsyg.2020.00448/full#supplementary-material>

Amstadter, A. B. (2008). Emotion regulation and anxiety disorders. *J. Anxiety Disord.* 22, 211–221. doi: 10.1016/j.janxdis.2007.02.004

Barrett, L. F. (2017). The theory of constructed emotion: an active inference account of interoception and categorization. *Soc. Cogn. Affect. Neurosci.* 12, 1–23. doi: 10.1093/scan/nsw154

- Baucom, L. B., Wedell, D. H., Wang, J., Blitzer, D. N., and Shinkareva, S. V. (2012). Decoding the neural representation of affective states. *Neuroimage* 59, 718–727. doi: 10.1016/j.neuroimage.2011.07.037
- Borkenau, P., and Ostendorf, F. (2008). *NEO-FFI NEO-Fünf-Faktoren-Inventar nach Costa und McCrae (NEO-FFI) (2., neu normierte und vollständig überarbeitete Auflage)*. Göttingen: Hogrefe.
- Bracci, S., Op, and de Beeck, H. (2016). Dissociations and associations between shape and category representations in the two visual pathways. *J. Neurosci.* 36, 432–444. doi: 10.1523/JNEUROSCI.2314-15.2016
- Broadbent, D., and Broadbent, M. (1988). Anxiety and attentional bias: state and trait. *Cogn. Emot.* 2, 165–183. doi: 10.1080/02699938808410922
- Brouwer, G. J., and Heeger, D. J. (2013). Categorical clustering of the neural representation of color. *J. Neurosci.* 33, 15454–15456. doi: 10.1523/jneurosci.2472-13.2013
- Bush, K. A., Privratsky, A., Gardner, J., Zielinski, M. J., and Kilts, C. D. (2018). Common functional brain states encode both perceived emotion and the psychophysiological response to affective stimuli. *Sci. Rep.* 8:15444. doi: 10.1038/s41598-018-33621-6
- Cacioppo, J. T., Cacioppo, S., Dulawa, S., and Palmer, A. A. (2014). Social neuroscience and its potential contribution to psychiatry. *World Psychiatry* 13, 131–139. doi: 10.1002/wps.20118
- Carpenter, R. W., and Trull, T. J. (2013). Components of emotion dysregulation in borderline personality disorder: a review. *Curr. Psychiatry Rep.* 15, 335.
- Chan, S. W. Y., Goodwin, G. M., and Harmer, C. J. (2007). Highly neurotic never-depressed students have negative biases in information processing. *Psychol. Med.* 37, 1281–1291. doi: 10.1017/S0033291707000669
- Charest, I., Kievit, R. A., Schmitz, T. W., Deca, D., and Kriegeskorte, N. (2014). Unique semantic space in the brain of each beholder predicts perceived similarity. *Proc. Natl. Acad. Sci. U.S.A.* 111, 14565–14570. doi: 10.1073/pnas.1402594111
- Chikazoe, J., Lee, D. H., Kriegeskorte, N., and Anderson, A. K. (2014). Population coding of affect across stimuli, modalities and individuals. *Nat. Neurosci.* 17, 1114–1122. doi: 10.1038/nn.3749
- Costa, P. T., and McCrae, R. R. (1980). Influence of extraversion and neuroticism on subjective well-being: happy and unhappy people. *J. Pers. Soc. Psychol.* 38, 668–678. doi: 10.1037/0022-3514.38.4.668
- Costa, P. T., and McCrae, R. R. (1992). *Revised NEO Personality Inventory (NEO-PI-R) and NEO Five-Factor Inventory (NEO-FFI) professional manual*. Odessa, FL: Psychological Assessment Resources.
- Costa, P. T., McCrae, R. R., and Dye, D. A. (1991). Facet scales for agreeableness and conscientiousness: a revision of the NEO personality inventory. *Pers. Individ. Differ.* 12, 887–888. doi: 10.1016/0191-8869(91)90177-D
- Cowen, A. S., and Keltner, D. (2017). Self-report captures 27 distinct categories of emotion bridged by continuous gradients. *Proc. Natl. Acad. Sci. U.S.A.* 114, E7900–E7909. doi: 10.1073/pnas.1702247114
- Dauvier, B., Pavani, J.-B., Le Vigouroux, S., Kop, J.-L., and Congard, A. (2019). The interactive effect of neuroticism and extraversion on the daily variability of affective states. *J. Res. Pers.* 78, 1–15. doi: 10.1016/j.jrp.2018.10.007
- Derryberry, D., and Reed, M. A. (1994). Temperament and attention: orienting toward and away from positive and negative signals. *J. Pers. Soc. Psychol.* 66, 1128–1139. doi: 10.1037/0022-3514.66.6.1128
- Dubois, J., Galdi, P., Han, Y., Paul, L. K., and Adolphs, R. (2018). Resting-State functional brain connectivity best predicts the personality dimension of openness to experience. *Personal. Neurosci.* 1, 1–21. doi: 10.1017/pen.2018.8
- Dunsmoor, J. E., Kragel, P. A., Martini, A., and LaBar, K. S. (2014). Aversive learning modulates cortical representations of object categories. *Cereb. Cortex* 24, 2859–2872. doi: 10.1093/cercor/bht138
- Ekman, P. (2016). What scientists who study emotion agree about. *Perspect. Psychol. Sci.* 11, 31–34. doi: 10.1177/1745691615596992
- Etkin, A., and Wager, T. D. (2007). Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am. J. Psychiatry* 164, 1476–1488. doi: 10.1176/appi.ajp.2007.07030504
- Eysenck, H. J. (1967). *The Biological Basis of Personality*. Springfield, IL: Charles C. Thomas.
- Feczko, E., Miranda-Dominguez, O., Marr, M., Graham, A. M., Nigg, J. T., and Fair, D. A. (2019). The heterogeneity problem: approaches to identify psychiatric subtypes. *Trends Cogn. Sci.* 23, 584–601. doi: 10.1016/j.tics.2019.03.009
- Goldstone, R. (1994). An efficient method for obtaining similarity data. *Behav. Res. Methods Instrum. Comput.* 26, 381–386. doi: 10.3758/BF03204653
- Gomez, R., Gomez, A., and Cooper, A. (2002). Neuroticism and extraversion as predictors of negative and positive emotional information processing: comparing Eysenck's, Gray's, and Newman's theories. *Eur. J. Pers.* 16, 333–350. doi: 10.1002/per.459
- Goodkind, M. S., Sollberger, M., Gyurak, A., Rosen, H. J., Rankin, K. P., Miller, B., et al. (2012). Tracking emotional valence: the role of the orbitofrontal cortex. *Hum. Brain Mapp.* 33, 753–762. doi: 10.1002/hbm.21251
- Grimm, S., Schmidt, C. F., Bermpohl, F., Heinzel, A., Dahlem, Y., Wyss, M., et al. (2006). Segregated neural representation of distinct emotion dimensions in the prefrontal cortex—an fMRI study. *Neuroimage* 30, 325–340. doi: 10.1016/j.neuroimage.2005.09.006
- Gross, J. J., and Jazaieri, H. (2014). Emotion, emotion regulation, and psychopathology. *Clin. Psychol. Sci.* 2, 387–401. doi: 10.1177/2167702614536164
- Gross, J. J., Sutton, S. K., and Ketelaar, T. (1998). Relations between affect and personality: support for the affect-level and affective-reactivity views. *Personal. Soc. Psychol. Bull.* 24, 279–288. doi: 10.1177/0146167298243005
- Hamann, S. (2012). Mapping discrete and dimensional emotions onto the brain: controversies and consensus. *Trends Cognit. Sci.* 16:458. doi: 10.1016/j.tics.2012.07.006
- Hamann, S., and Canli, T. (2004). Individual differences in emotion processing. *Curr. Opin. Neurobiol.* 14, 233–238. doi: 10.1016/j.conb.2004.03.010
- Hamburg, M. A., and Collins, F. S. (2010). The path to personalized medicine – perspective. *N. Engl. J. Med.* 363, 301–304. doi: 10.1056/NEJMp1006304
- Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., and Pietrini, P. (2001). Distributed and overlapping representations of faces and objects in ventral temporal cortex. *Science* 293, 2425–2430. doi: 10.1126/science.1063736
- Iordan, M. C., Greene, M. R., Beck, D. M., and Fei-Fei, L. (2015). Basic level category structure emerges gradually across human ventral visual cortex. *J. Cognit. Neurosci.* 27, 1427–1446. doi: 10.1162/jocn_a_00790
- Iordan, M. C., Greene, M. R., Beck, D. M., and Fei-Fei, L. (2016). Typicality sharpens category representations in object-selective cortex. *Neuroimage* 134, 170–179. doi: 10.1016/j.neuroimage.2016.04.012
- Javaras, K. N., Schaefer, S. M., van Reekum, C. M., Lapate, R. C., Greischar, L. L., Bachhuber, D. R., et al. (2012). Conscientiousness predicts greater recovery from negative emotion. *Emotion* 12, 875–851. doi: 10.1037/a0028105
- Knüppel, H., Metz, C., Meerpohl, J. J., and Strech, D. (2013). How psychiatry journals support the unbiased translation of clinical research. A cross-sectional study of editorial policies. *PLoS ONE* 8:e75995. doi: 10.1371/journal.pone.0075995
- Koch, A., Alves, H., Krüger, T., and Unkelbach, C. (2016). A general valence asymmetry in similarity: good is more alike than bad. *J. Exp. Psychol. Learn. Mem. Cognit.* 42, 1171–1192. doi: 10.1037/xlm0000243
- Kokkonen, M., and Pulkkinen, L. (2001). Extraversion and neuroticism as antecedents of emotion regulation and dysregulation in adulthood. *Eur. J. Pers.* 15, 407–424. doi: 10.1002/per.425
- Kotov, R., Gamez, W., Schmidt, F., and Watson, D. (2010). Linking “big” personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychol. Bull.* 136, 768–821. doi: 10.1037/a0020327
- Kragel, P. A., Knodt, A. R., Hariri, A. R., and LaBar, K. S. (2016). Decoding spontaneous emotional states in the human brain. *PLoS Biol.* 14:2000106. doi: 10.1371/journal.pbio.2000106
- Kragel, P. A., and LaBar, K. S. (2015). Multivariate neural biomarkers of emotional states are categorically distinct. *Soc. Cogn. Affect. Neurosci.* 10, 1437–1448. doi: 10.1093/scan/nsv032
- Kriegeskorte, N., and Mur, M. (2012). Inverse MDS: inferring dissimilarity structure from multiple item arrangements. *Front. Psychol.* 3:245. doi: 10.3389/fpsyg.2012.00245
- Kryklywy, J. H., Macpherson, E. A., and Mitchell, D. G. V. (2018). Decoding auditory spatial and emotional information encoding using multivariate versus univariate techniques. *Exp. Brain Res.* 236, 945–953. doi: 10.1007/s00221-018-5185-7
- Lance, G. N., and Williams, W. T. (1967). A general theory of classificatory sorting strategies: I. Hierarchical systems. *Comput. J.* 9, 373–380. doi: 10.1093/comjnl/9.4.373

- Lang, P. J., Bradley, M. M., and Cuthbert, B. N. (2008). International affective picture system (IAPS): affective ratings of pictures and instruction manual.
- Larsen, R. J., and Ketelaar, T. (1989). Extraversion, neuroticism and susceptibility to positive and negative mood induction procedures. *Pers. Individ. Differ.* 10, 1221–1228. doi: 10.1016/0191-8869(89)90233-X
- Levine, S. M., Alahäivälä, A. L. I., Wackerle, A., Rupperecht, R., and Schwarzbach, J. V. (2018a). Identifying emotion subtypes from individualized emotion space. *PsyArXiv* [Preprint]. doi: 10.31234/osf.io/4u6vn
- Levine, S. M., Pfäller, M., Reichenberger, J., Shibani, Y., Mühlberger, A., Rupperecht, R., et al. (2018b). Relating experimentally-induced fear to pre-existing phobic fear in the human brain. *Soc. Cogn. Affect. Neurosci.* 13, 164–172. doi: 10.1093/scan/nsx147
- Levine, S. M., Wackerle, A., Rupperecht, R., and Schwarzbach, J. V. (2018c). The neural representation of an individualized relational affective space. *Neuropsychologia* 120, 35–42. doi: 10.1016/j.neuropsychologia.2018.10.008
- Lindquist, K. A., Wager, T. D., Kober, H., Bliss-Moreau, E., and Barrett, L. F. (2012). The brain basis of emotion: a meta-analytic review. *Behav. Brain Sci.* 35, 121–143. doi: 10.1017/S0140525X11000446
- Liu, L., Preotiu-Pietro, D., Samani, Z. R., Moghaddam, M. E., and Ungar, L. (2016). “Analyzing personality through social media profile picture choice,” in *Proceedings of the 10th International AAAI Conference on Web and Social Media* (Menlo Park, CA: AAAI).
- Machado-Vieira, R. (2012). Tracking the impact of translational research in psychiatry: state of the art and perspectives. *J. Transl. Med.* 10:175. doi: 10.1186/1479-5876-10-175
- MacLeod, C., and Mathews, A. (1988). Anxiety and the allocation of attention to threat. *Q. J. Exp. Psychol. Sect. A* 40, 653–670. doi: 10.1080/14640748808402292
- McCrae, R. R., and Costa, P. T. (1985). Updating norman’s “adequate taxonomy”: intelligence and personality dimensions in natural language and in questionnaires. 49, 710–721. doi: 10.1037/0022-3514.49.3.710
- McCrae, R. R., Costa, P. T., and Busch, C. M. (1986). Evaluating comprehensiveness in personality systems: the California Q-Set and the five-factor model. *J. Pers.* 4, 430–446. doi: 10.1111/j.1467-6494.1986.tb00403.x
- Mennin, D. S., Heimberg, R. G., Turk, C. L., and Fresco, D. M. (2005). Preliminary evidence for an emotion dysregulation model of generalized anxiety disorder. *Behav. Res. Ther.* 43, 1281–1310. doi: 10.1016/j.brat.2004.08.008
- Miller, E. K., and Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annu. Rev. Neurosci.* 24, 167–202. doi: 10.1146/annurev.neuro.24.1.167
- Most, S. B., Chun, M. M., Johnson, M. R., and Kiehl, K. A. (2006). Attentional modulation of the amygdala varies with personality. *Neuroimage* 31, 934–944. doi: 10.1016/j.neuroimage.2005.12.031
- Mulders, P., Llera, A., Tendolkar, I., van Eijndhoven, P., and Beckmann, C. (2018). Personality profiles are associated with functional brain networks related to cognition and emotion. *Sci. Rep.* 8:13874. doi: 10.1038/s41598-018-32248-x
- Mur, M., Meys, M., Bodurka, J., Goebel, R., Bandettini, P. A., and Kriegeskorte, N. (2013). Human object-similarity judgments reflect and transcend the primate-IT object representation. *Front. Psychol.* 4:128. doi: 10.3389/fpsyg.2013.00128
- Nastase, S. A., Connolly, A. C., Oosterhof, N. N., Halchenko, Y. O., Guntupalli, J. S., Visconti, et al. (2017). Attention selectively reshapes the geometry of distributed semantic representation. *Cereb. Cortex* 27, 4277–4291. doi: 10.1093/cercor/bhx138
- Ng, W., and Diener, E. (2009). Personality differences in emotions. *J. Individ. Differ.* 30, 100–106. doi: 10.1027/1614-0001.30.2.100
- Nielen, M. M. A., Heslenfeld, D. J., Heinen, K., Van Strien, J. W., Witter, M. P., Jonker, C., et al. (2009). Distinct brain systems underlie the processing of valence and arousal of affective pictures. *Brain Cognit.* 71, 387–396. doi: 10.1016/j.bandc.2009.05.007
- Nostro, A. D., Müller, V. I., Varikuti, D. P., Pläschke, R. N., Hoffstaedter, F., Langner, R., et al. (2018). Predicting personality from network-based resting-state functional connectivity. *Brain Struct. Funct.* 223, 2699–2719. doi: 10.1007/s00429-018-1651-z
- Nummenmaa, L., Glerean, E., Hari, R., and Hietanen, J. K. (2014). Bodily maps of emotions. *Proc. Natl. Acad. Sci. U.S.A.* 111, 646–651. doi: 10.1073/pnas.1321664111
- Nummenmaa, L., Hari, R., Hietanen, J. K., and Glerean, E. (2018). Maps of subjective feelings. *Proc. Natl. Acad. Sci. U.S.A.* 115, 9198–9203. doi: 10.1073/pnas.1807390115
- Passamonti, L., Riccelli, R., Indovina, I., Duggento, A., Terracciano, A., and Toschi, N. (2019). Time-resolved connectome of the five-factor model of personality. *Sci. Rep.* 9:15066. doi: 10.1038/s41598-019-51469-2
- Reimherr, F. W., Marchant, B. K., Strong, R. E., Hedges, D. W., Adler, L., Spencer, T. J., et al. (2005). Emotional dysregulation in adult ADHD and response to atomoxetine. *Biol. Psychiatry* 58, 125–131. doi: 10.1016/j.biopsych.2005.04.040
- Richards, A., French, C. C., Johnson, W., Naparstek, J., and Williams, J. (1992). Effects of mood manipulation and anxiety on performance of an emotional Stroop task. *Br. J. Psychol.* 83, 479–491. doi: 10.1111/j.2044-8295.1992.tb02454.x
- Rousseeuw, P. J. (1987). Silhouettes: a graphical aid to the interpretation and validation of cluster analysis. *J. Comput. Appl. Mathemat.* 20, 53–65. doi: 10.1016/0377-0427(87)90125-7
- Rusting, C. L., and Larsen, R. J. (1997). Extraversion, neuroticism, and susceptibility to positive and negative affect: a test of two theoretical models. *Pers. Individ. Differ.* 22, 607–612. doi: 10.1016/S0191-8869(96)00246-2
- Saarimäki, H., Ejtehadian, L. F., Glerean, E., Jääskeläinen, I. P., Vuilleumier, P., Sams, M., et al. (2018). Distributed affective space represents multiple emotion categories across the human brain. *Soc. Cogn. Affect. Neurosci.* 13, 471–482. doi: 10.1093/scan/nsy018
- Saarimäki, H., Gotsopoulos, A., Jääskeläinen, I. P., Lampinen, J., Vuilleumier, P., Hari, R., et al. (2016). Discrete neural signatures of basic emotions. *Cereb. Cortex* 26, 2563–2573. doi: 10.1093/cercor/bhv086
- Schwarz, G. (1978). Estimating the dimension of a model. *Ann. Stat.* 6, 461–464. doi: 10.1214/aos/1176344136
- Shinkareva, S. V., Wang, J., Kim, J., Facciani, M. J., Baucom, L. B., and Wedell, D. H. (2014). Representations of modality-specific affective processing for visual and auditory stimuli derived from functional magnetic resonance imaging data. *Hum. Brain Mapp.* 35, 3558–3568. doi: 10.1002/hbm.22421
- Skerry, A. E., and Saxe, R. (2015). Neural representations of emotion are organized around abstract event features. *Curr. Biol.* 25, 1945–1954. doi: 10.1016/j.cub.2015.06.009
- Smith, K. A., Barstead, M. G., and Rubin, K. H. (2017). Neuroticism and conscientiousness as moderators of the relation between social withdrawal and internalizing problems in adolescence. *J. Youth Adolesc.* doi: 10.1007/s10964-016-0594-z
- Sutin, A. R., Terracciano, A., Deiana, B., Naitza, S., Ferrucci, L., Uda, M., et al. (2010). High neuroticism and low conscientiousness are associated with interleukin-6. *Psychol. Med.* 40, 1485–1493. doi: 10.1017/S0033291709992029
- Taylor, S. F., Kang, J., Brege, I. S., Tso, I. F., Hosanagar, A., and Johnson, T. D. (2012). Meta-analysis of functional neuroimaging studies of emotion perception and experience in schizophrenia. *Biol. Psychiatry* 71, 136–145. doi: 10.1016/j.biopsych.2011.09.007
- Thomsen, D. K., Olesen, M. H., Schnieber, A., and Tønnesvang, J. (2014). The emotional content of life stories: positivity bias and relation to personality. *Cognit. Emot.* 28, 260–277. doi: 10.1080/02699931.2013.815155
- Toschi, N., Riccelli, R., Indovina, I., Terracciano, A., and Passamonti, L. (2018). Functional connectome of the five-factor model of personality. *Personal. Neurosci.* 1, 1–10. doi: 10.1017/pen.2017.2
- Trull, T. J., and Sher, K. J. (1994). Relationship between the five-factor model of personality and axis I disorders in a nonclinical sample. *J. Abnorm. Psychol.* 103, 350–360. doi: 10.1037/0021-843X.103.2.350
- Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., and Ugurbil, K. (2013). The WU-minn human connectome project: an overview. *Neuroimage* 80, 62–79. doi: 10.1016/j.neuroimage.2013.05.041
- van Tilburg, W. A. P., and Igou, E. R. (2017). Boredom begs to differ: differentiation from other negative emotions. *Emotion* 17, 309–322. doi: 10.1037/emo0000233

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Levine, Alahäivälä, Wechsler, Wackerle, Rupperecht and Schwarzbach. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.