







## STUDY PROTOCOL

# The evidence for interventions in early childhood allergy prevention – towards a living systematic review: protocol

## [version 1; peer review: awaiting peer review]

Uwe Matterne <sup>1</sup>, Christina Tischer<sup>2</sup>, Jiancong Wang <sup>1</sup>, Helge Knüttel <sup>3</sup>,  
Jon Genuneit<sup>4</sup>, Michael Perkin<sup>5</sup>, Christian Apfelbacher <sup>1,6</sup>

<sup>1</sup>Institute of Social Medicine and Health Systems Research, Otto von Guericke University Magdeburg, Leipziger Str. 44, Magdeburg, 39120, Germany

<sup>2</sup>Institute for Health Resort Medicine and Health Promotion, State Institute of Health, Bavarian Health and Food Safety Authority, Prinzregentenstraße 6, Bad Kissingen, 97688, Germany

<sup>3</sup>University Library, University of Regensburg, Universitätsstraße 31, Regensburg, 93053, Germany

<sup>4</sup>Pediatric Epidemiology, Department of Pediatrics, Medical Faculty, Leipzig University, Liebigstraße 20a, Leipzig, 04103, Germany

<sup>5</sup>Population Health Research Institute, St George's, University of London, Cranmer Terrace, London, SW17 0RE, UK

<sup>6</sup>Family Medicine and Primary Care, Lee Kong Chian School of Medicine, Nanyang Technological University Singapore, 11 Mandalay Road, 308232, Singapore

**V1** First published: 24 Mar 2021, 10:235  
<https://doi.org/10.12688/f1000research.51490.1>

Latest published: 24 Mar 2021, 10:235  
<https://doi.org/10.12688/f1000research.51490.1>

### Abstract

**Background:** Research in early childhood allergy prevention (ECAP) is flourishing and new intervention strategies have proven to be promising. Due to the dynamic nature of ECAP, gaps between what is known and how guidelines inform practice are likely. A living systematic review (LSR) can narrow this gap by incorporating new evidence as it becomes available. No efficacy comparisons across various ECAP interventions for similar outcomes have been carried out. Networks of randomised clinical trials can be evaluated in the context of a network meta-analysis (NMA). We aim to establish a LSR on the efficacy and safety of any intervention investigated in randomised controlled trials (RCT) to prevent the occurrence of allergic sensitisation (AS), symptoms or diagnoses of allergic diseases in infancy and early childhood (0-3 years).

**Methods:** A baseline SR will synthesise the evidence from existing SRs of RCTs as well as RCTs not yet considered in these. After completion of the baseline SR we propose to conduct a LSR. Using this methodology, we aim to undertake constant evidence surveillance, three-monthly search updates, and review updates every three months, should new evidence emerge.

**Conclusions:** The ECAP evidence landscape has undergone dramatic transformations and this process is likely to continue. As a response to this, a LSR offers the potential to allow more timely synthesis of new evidence as it emerges. Long gaps between updates of SRs makes it

### Open Peer Review

**Reviewer Status** AWAITING PEER REVIEW

Any reports and responses or comments on the article can be found at the end of the article.

harder for guidelines and recommendations to be up to date. Users of information, such as parents, may be confused if they encounter new evidence that is not part of a trusted guideline. A LSR approach allows us to continuously search the literature and update the evidence-base of existing ECAP interventions resulting in a decreased timespan from evidence accrual to informing clinical practice.

### Keywords

Early childhood allergy prevention, randomised controlled trial, living systematic review



This article is included in the [Living Evidence](#) collection.

**Corresponding author:** Uwe Matterne ([uwe.matterne@med.ovgu.de](mailto:uwe.matterne@med.ovgu.de))

**Author roles:** **Matterne U:** Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Tischer C:** Writing – Review & Editing; **Wang J:** Writing – Review & Editing; **Knüttel H:** Methodology, Software, Validation, Writing – Review & Editing; **Genuneit J:** Methodology, Validation, Writing – Review & Editing; **Perkin M:** Methodology, Validation, Writing – Review & Editing; **Apfelbacher C:** Conceptualization, Funding Acquisition, Methodology, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing

**Competing interests:** No competing interests were disclosed by UM, CT, JW, HK and MP. CA is spokesperson of the Health Literacy in Early Childhood Allergy Prevention (HELICAP) research group, and has received consultancy fees from Sanofi Genzyme and LEOPharma. JG was the project manager at Leipzig University on unrestricted research grants from Danone Nutricia Research to Ulm University and to Leipzig University for research into human milk composition within the Ulm SPATZ Health Study and the Ulm Birth Cohort Study. While the grant periods have expired, work with the data generated as part of these grants continues.

**Grant information:** We acknowledge the financial support of this project by the German Research Foundation (DFG): AP 235/3-1, funded as subproject of DFG research group FOR 2959 (HELICAP), AP 235/4-1. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2021 Matterne U *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Matterne U, Tischer C, Wang J *et al.* **The evidence for interventions in early childhood allergy prevention – towards a living systematic review: protocol [version 1; peer review: awaiting peer review]** F1000Research 2021, 10:235 <https://doi.org/10.12688/f1000research.51490.1>

**First published:** 24 Mar 2021, 10:235 <https://doi.org/10.12688/f1000research.51490.1>

## Introduction

### Rationale

Allergy in children is common. Frequent food allergies (FA) in children include hen's egg, cow's milk and peanut.<sup>1</sup> Around 10% of children are affected by FA and the incidence is still rising in developing countries.<sup>1</sup> Food allergy impacts quality of life.<sup>2-4</sup> Allergic diseases (eczema,<sup>5-7</sup> asthma and hay fever/allergic rhinitis)<sup>8,9</sup> are also highly prevalent and associated with decreased health-related quality of life (HRQOL).<sup>10</sup>

To counteract the large number of allergies and reduce their burden, a major shift from merely managing manifest allergy to preventing its occurrence has taken place. Previous prevention efforts revolved around avoidance of potential allergens (particularly in at risk individuals), while more recently a new paradigm has been embraced whose focus is on early allergen exposure to induce immune tolerance. This new paradigm has been informing study designs for food allergy prevention. So far it has been established that oral tolerance induction is allergen specific and efficacious in single introduction trials of peanut and egg.<sup>11</sup>

There is also research activity revolving around environmentally derived prevention paradigms (i.e. exposure to rural environments, cowshed pill, or unpasteurised milk).<sup>12</sup> However, one trial found no evidence that an orally applied bacterial lysate affected allergy development in at risk infants.<sup>13</sup> It is however, expected that more research in this field will emerge soon.

In general, the design of preventive strategies has been informed by several hypotheses regarding the aetiology of allergy and allergic disease. The original hygiene hypothesis states that lack of exposure to common infections causes allergy.<sup>14</sup> However, not all infections protect from allergy, resulting in criticism of the hypothesis.<sup>15</sup> Further research has led to modifications of this hypothesis, suggesting that exposure to specific pathogens, commensals and symbionts protects against development of allergy.<sup>16,17</sup> The dual-allergen-exposure hypothesis proposes that exposure to food allergens through the skin leads to allergy, while early consumption of these foods induces tolerance.<sup>18</sup> While in the past it was advocated that avoidance of allergenic food would reduce the onset of allergy,<sup>19</sup> several recent trials have lent support to the dual-allergen-exposure hypothesis.<sup>20-22</sup> The vitamin D hypothesis suggests that low vitamin D levels increase the risk of developing food allergy.<sup>23,24</sup> There are, however, conflicting data on the relationship between vitamin D and the development of food allergy.<sup>25</sup>

Research in early childhood allergy prevention (ECAP) is flourishing and more than 50 systematic reviews (SR) that exclusively or partly reviewed RCTs<sup>26-81</sup> have been published, some of which provide useful insights. However, not all include a standardised evaluation of the quality of the evidence (e.g. Grading of Recommendations Assessment, Development and Evaluation (GRADE))<sup>82</sup> for randomised controlled trials (RCT).<sup>83</sup> Risk of bias (RoB) assessment was done by a variety of different tools. Some used the National Institute for Health and Care Excellence (NICE) methodological checklist<sup>84</sup> or the Strength of Recommendation Taxonomy (SORT)<sup>85</sup> criteria. Thus, a homogenous RoB assessment and consistent evidence grading approach for each study type would be desirable.

### Research challenges

As intervention effects may vary depending on whether at risk, not at-risk or children with manifest allergy are investigated, studies will be grouped accordingly. ECAP in the latter would still be considered primary prevention, rather than secondary or tertiary prevention, since we would look only at the prevention of new allergies.

Many of the above SRs included observational studies as well as clinical trials. Studies examining the effect of breastfeeding versus no breastfeeding on allergy development cannot use an RCT design; however, for most other ECAP strategies, there are no ethical or other constraints that prevent the use of the gold standard RCT. Hence, we will only consider RCTs in our SR.

Not all ECAP findings may yet be part of allergy prevention guidelines. As outlined above, due to the wealth of studies and new ECAP paradigms, more than 50 SRs have attempted to synthesise the evidence accumulated within the various approaches to ECAP. It is likely that there are gaps between what we know from the best available research and what happens in healthcare practice due to the dynamic nature of ECAP research. The incorporation of new findings into existing reviews or their updates is time-consuming and not always feasible. Hence, adapting prevention guidelines in light of new findings and providing health care providers and other information users with up-to-date evidence may be impeded. Recent advances in the presentation of systematically reviewed (qualitative and meta-analytic) data have led to the concept of a living systematic review (LSR). A LSR is defined as "a systematic review that is continually updated, incorporating relevant new evidence as it becomes available".<sup>86</sup>

Furthermore, no efficacy comparisons across various ECAP strategies for similar outcomes have been carried out. Networks of randomised clinical trials can be evaluated in the context of a network meta-analysis (NMA). NMA refers to a procedure that allows inferences about the comparative effectiveness of interventions that may or may not have been evaluated directly against each other.<sup>87–89</sup>

### Objective

We aim to establish an LSR on the efficacy and safety of any intervention to prevent the occurrence of allergic sensitisation (AS), symptoms or diagnoses of allergic diseases in infancy and early childhood (0-3 years).

The specific objectives are:

1. To identify all individual-level interventions using the oral, skin, environmental or pharmaceutical route for the prevention of allergy in allergy-free children or the prevention of new allergies in children with manifest allergy
2. To identify all community-level interventions (such as community programmes promoting dietary and environmental diversity in early life) which have thus far been investigated in RCTs to prevent the occurrence of allergy in infancy.
3. To summarise the evidence regarding the effects (efficacy and safety) of these interventions in preventing the occurrence of allergy in infancy and early childhood.
4. To judge the quality of this evidence.
5. To provide a corresponding plain language summary (PLS) accessible for consumers.
6. To develop a workflow for an LSR, which ensures that the evidence synthesis is continuously kept up-to-date.

### Methods

The review will be undertaken according to the methods outlined in the Cochrane Handbook for Systematic Reviews of Interventions,<sup>90</sup> PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses)<sup>91</sup> and its extension for application to NMA.<sup>92</sup> A protocol registration will be made at PROSPERO.<sup>93</sup> Any updates to the protocol will be made through PROSPERO. The protocol was developed closely considering the Prisma-P<sup>94</sup> checklist.<sup>95</sup> A baseline SR will synthesise the evidence available so far and then transformed into a LSR by regular updates.

### Eligibility criteria

Studies will be eligible for inclusion in the review should they meet the following (PICO) criteria:

- Date of publication: 1980 onward
- Types of study: Randomised controlled trials,
- Population: expectant and/or breastfeeding mothers of and/or children 0 - 3 years (at time of intervention), at risk children (0 - 3 years) (at least one parent with known allergic disease), not at risk children (0-3 years) (no parental allergic disease), children (0-3 years) with manifest allergy (only if intervention aims at preventing a new allergy or study reports this outcome)
- Interventions: any aimed at the individual or community level at preventing the onset of new allergy or allergic disease or allergic sensitisation
  - o Oral route (supplements (e.g. vitamins, minerals, pro-, pre-, symbiotic, gut bacteria), time and presence of allergenic food introduction (e.g. peanut, egg protein, cow's milk, fish), variation in condition (raw, cooked, pasteurised, fermented) and/or amount of complimentary food introduction, avoidance of potential allergens
  - o Skin route (e.g. emollients, treated water for washing)
  - o Environmental interventions (exposure to natural outdoor environments, green space, outdoor spaces, exposure to farm environments (cow- and other animal-shed bacteria, farming associated bacteria and

- o microbes, bacterial lysate, acinetobacter, microbiome, mucous membrane, microbiota), avoidance of chemicals or allergens (e.g. mattress protector (mites))
- o Pharmaceutical prevention (e.g. allergen-immuno-therapy (AIT), Bacillus Calmette-Guérin (BCG) vaccination))
- Comparator: inactive comparator such as placebo, no intervention or usual care
- Primary outcomes:
  - o Physician-diagnosed or parent-reported incidence of allergic asthma (AA), allergic rhinitis (AR), atopic eczema (AE), food allergy (FA)
  - o Physician-diagnosed or parent-reported recurrent symptoms of sneeze, wheeze, cough, itch, flexural eczema, or FA
- Secondary outcomes:
  - o incidence of AS measured by in-vivo tests such as skin-prick test (SPT)
  - o incidence of AS measured by in-vitro tests such as fluorescently labelled anti-IgE antibody, enzyme-linked immunosorbent assay (ELISA), or Radio-Allergo-Sorbent-Test (RAST)
  - o adverse events (AE), severe adverse events (SAE), withdrawals

ECAP has revolved around antenatal and postnatal strategies targeting the mother and strategies targeting the child after birth (often during a critical period). Strategies that have been explored in RCTs are illustrated in [Table 1](#).

**Exclusion criteria**

Any observational research (cross-sectional, case-control, case-series, prospective/retrospective cohort studies) or quasi-experimental studies (matched controlled designs) will be excluded. We will not consider interventions aimed at treating allergy unless the intervention is also hypothesised to reduce the onset of a new allergic manifestation.

**Table 1. Overview of primary ECAP strategies investigated thus far or in progress in RCTs**

Intervention route	Example	Mother		Infant/Child		
		Prenatal	If breastfeeding postnatal	At risk	Not at risk	One or more AS, AD
Oral route	Diet/nutrient supplementation	✓ <sup>109</sup>	✓ <sup>40</sup>	✓ <sup>110</sup>	✓ <sup>67</sup>	✓ <sup>111</sup>
	Diet/nutrient avoidance	✓ <sup>112</sup>	✓ <sup>112</sup>	✓ <sup>113</sup>	None found	✓ <sup>22</sup>
	Early allergenic food introduction	n.a.	n.a.	✓ <sup>20</sup>	✓ <sup>108</sup>	✓ <sup>22</sup>
Skin route	Skincare	n.a.	n.a.	✓ <sup>39</sup>	✓ <sup>39</sup>	✓ <sup>114</sup>
Environmental route	House dust mite avoidance	n.a.	n.a.	✓ <sup>31</sup>	None found	
Pharmaceutical route	Allergen immuno-therapy (AIT)	n.a.	n.a.	✓ <sup>115</sup>	None found	✓ <sup>116</sup>
	BCG vaccination	n.a.	n.a.	✓ <sup>117</sup>	✓ <sup>118</sup>	
	Oral H1-antihistamines	n.a.	n.a.	n.a.	n.a.	✓ <sup>119</sup>

ECAP: early childhood allergy prevention, RCT, randomised controlled trial, AD: allergic disease, AS: allergic sensitisation, superscript numbers indicate an example reference.

## Search methods

We will use five approaches for the identification of studies for the baseline SR:

### 1) Topic-based searches in databases and registries

We will search for all relevant RCTs regardless of publication status (published, unpublished, in press, or ongoing) in the following bibliographic databases: MEDLINE (Ovid), Embase (Ovid), CENTRAL (Cochrane Library), Science Citation Index Expanded & Social Sciences Citation Index (Web of Science), Cochrane Skin Group Specialized Register, GREAT (The Global Resource of Eczema Trials, Centre of Evidence Based Dermatology), in clinical trial registries (U.S. National Institutes of Health [ClinicalTrials.gov](https://clinicaltrials.gov), ISRCTN registry, Australian New Zealand Clinical Trials Registry, World Health Organization International Clinical Trials Registry Platform, EU Clinical Trials Register).

We will search the conference proceedings (European Academy of Dermatology and Venereology (EADV), European Academy of Allergy and Clinical Immunology (EAACI), American Academy of Dermatology (AAD), American Academy of Allergy, Asthma & Immunology (AAAI), Asia Pacific Association of Allergy, Asthma and Clinical Immunology (APAAACI) and Asia Pacific Association of Pediatric Allergy, Respiriology & Immunology (APAPARI), World Allergy Organisation (WAO)) and Sociedad Latinoamericana de Alergia, Asma e Inmunología, Asunción (SLAAI). Additionally, we will review published documents from health technology assessment agencies (e.g. NICE, IQWiG). No restriction on status or year of publication will be applied. The search will be restricted to publications in English, German, French, Italian or Spanish. An initial draft search strategy for MEDLINE was developed by a medical librarian experienced in comprehensive searches for systematic reviews ([Box 1](#)). The performance of his strategy will be checked against the growing set of known relevant records. The final search strategy will be peer reviewed. The search strategy is composed of the components Population, Intervention, Outcome and Study filter that will be intersected with the Boolean AND operator. For each of the components relevant terms from text fields and controlled vocabulary were used in order to achieve high sensitivity. This strategy will be adapted to the other databases as appropriate.

### 2) Searches by trial registry numbers

Registry numbers of eligible trials will be collected to be used in follow-up searches with the aim to identify additional trial reports.

### 3) Trials included in relevant SRs

#### **Box 1:** Initial draft search strategy for MEDLINE

- 1 exp infant/ or Child, Preschool/ or (child or children).ti,ab,kf. or (pre-school\$ or preschool\$).ti,ab,kf. or Nurseries/ or (nursery or nurseries).ti,ab,kf. or exp Parents/ or (parent or parents or mother or mothers).ti,ab,kf. or (infant or infants).ti,ab,kf. or infancy.ti,ab,kf. or toddler?.ti,ab,kf. or (baby or babies).ti,ab,kf. or newborn\$.ti,ab,kf. or neonat\$.ti,ab,kf. or Pediatrics/ or (pediatric\$ or paediatric\$).ti,ab,kf. or early childhood.ti,ab,kf. or (Pregnant Women/ or Pregnancy/ or Prenatal Nutritional Physiological Phenomena/) or pregnan\$.ti,ab,kf. or Prenatal Exposure Delayed Effects/or Maternal Exposure/or ((maternal or prenatal) adj1 exposure\$).ti,ab,kf. or (fetus or fetuses or fetal or foetus or foetuses or foetal).ti,ab,kf. or Fetus/ (Population)
- 2 exp Preventive Health Services/ or Preventive Medicine/or "prevention control".fs. or prevent\$.ti,ab,kf. or prophyla\$.ti,ab,kf. or Infant Formula/or (formula or supplement\$).ti,ab,kf. or ((risk or protect\$ or development or avoidance or exposure or introduction) adj6 (allerg\$ or hypersensitiv\$ or atopy or atopic or dermatitis or neurodermatitis or asthma)).ti,ab,kf. (Intervention)
- 3 exp Hypersensitivity/ or Allergens/ or allerg\$.ti,ab,kf. or hypersensitiv\$.ti,ab,kf. or prick test\$.ti,ab,kf. or exp asthma/ or Dyspnea/or (asthma\$ or dyspnea or wheezing).ti,ab,kf. or (difficult\$ adj1 breathing).ti,ab,kf. or rhinoconjunctivitis.ti,ab,kf. or (atopic adj1 (dermatit\$ or neurodermatit\$ or eczema or disease)).ti,ab,kf. or Diaper Rash/or ((infant or infantile or diaper) adj1 (rash or rashes or eczema or dermatit\$)).ti,ab,kf. or Disseminated Neurodermat\$.ti,ab,kf. (Outcome)
- 4 ((randomized controlled trial or controlled clinical trial).pt. or randomized.ab. or placebo.ab. or drug therapy.fs. or randomly.ab. or trial.ab. or groups.ab.) not (exp animals/not humans.sh.) (Cochrane Highly Sensitive Search Strategy for identifying randomized trials in MEDLINE: sensitivity-maximizing version (2008 revision))
- 5 1 and 2 and 3 (Combined concepts: Patients AND Intervention AND Outcome)
- 6 5 and 4 (Combined concepts: Patients AND Intervention AND Outcome AND RCTs)

We will search for relevant SRs in order to identify additional trials in these SRs.

4) Reference lists of included trials

We will screen the reference lists of all included study reports. To aid in this process the reference lists will be downloaded from Web of Science when available.

5) Existing systematic reviews

For the creation of the baseline SR previous SRs on ECAP will be systematically searched for.

### Study selection, data extraction, analyses and syntheses

Database search results will be imported into [EppiReviewer](#) (version 4.11.5.3)<sup>96</sup> for deduplication and study selection. Assessment of eligibility, data extraction, risk of bias (RoB) evaluation and quality of the evidence assessment will be carried out by at least two researchers independently. The latter will be done according to the GRADE<sup>82</sup> recommendations.

The following data will be extracted:

Study characteristics: Author, year of publication, geographical region, study design, type of intervention, type of control, number of participants in intervention group and control group, study duration, time points of assessment, follow-up period, type of primary and secondary outcomes and safety indicators.

Participant characteristics: Person in whom intervention took place (mother (prenatal, postnatal), child, both), age, sex, ethnicity, allergic risk status, parental atopy, presence of allergic sensitisation and/or condition.

Study outcomes: Efficacy outcomes (unadjusted/adjusted): Incidence of allergic sensitization and/or allergic disease; safety outcomes: proportions of AEs, SAEs, withdrawal due to AEs

Existing systematic reviews will be incorporated into the baseline SR. Assessment of the quality of these will be carried out by A MeaSurement Tool to Assess systematic Reviews-2 (AMSTAR-2).<sup>97</sup> AMSTAR-2 is a critical appraisal tool for SRs that include randomised or non-randomised studies of healthcare interventions, or both.<sup>97</sup> Findings of SRs whose quality has been judged adequate will be summarised and studies of the same intervention and outcome not considered in these SR will be individually assessed for RoB and quality of the evidence.

### Meta-analyses and network-meta-analysis (NMA)

If no obvious qualitative heterogeneity within studies exists, we will perform meta-analyses across similar applications of interventions. Pair-wise meta-analyses between two intervention conditions (provided at least two eligible studies exist that are not too heterogeneous) for each different endpoint/outcome (FA, AS, AE, asthma, AR) will be conducted. These are to be updated in the LSR as relevant new studies emerge. To guard against potential type I error inflation and occurrence of type II errors which are a function of the number of analyses done with the same data (as new data is incorporated with each update) we will follow the recommendations outlined by Simmonds *et al.*<sup>98</sup>

Pairwise meta-analysis is a statistical technique for quantitatively synthesising similar studies in a systematic review. Useful in its own right, it is, however, limited in that it can only compare two interventions simultaneously, and only those evaluated directly in head-to-head trials.<sup>89</sup> Pairwise meta-analysis allows for comparisons between pairs of interventions (an experimental intervention and a comparator intervention) for a specific outcome in a particular population or setting. It is, however, often the case that a variety of different interventions are available for any given condition. A single SR that includes all relevant interventions and presents their comparative effectiveness and potential for harm would help people to decide between alternative interventions. NMA affords an analysis option for such a review.<sup>88</sup> A network of interventions consists of any set of studies that links three or more interventions via direct comparisons. Within a network, there can be numerous ways to make indirect comparisons between interventions. They refer to comparisons that have not been made directly within studies. Mathematical combinations of the available direct intervention effect sizes are used to estimate indirect effects. In NMA direct and indirect estimates across a network of interventions are combined in a single analysis.<sup>88</sup>

We also aim to conduct NMA to compare different interventions for the same endpoint/outcome (PO, SO and AE), should assumptions for NMA be met. We will also perform a full evaluation of the confidence in the results from NMA by using

the web application **CINeMA** (Confidence in Network Meta-Analysis).<sup>99</sup> This web application simplifies the evaluation of confidence in the findings from NMA and has evolved out of the GRADE approach. The GRADE<sup>82</sup> approach offers an assessment of the confidence in the results from systematic reviews and meta-analyses. Many organisations, for example the World Health Organisation, have adopted the GRADE approach.<sup>100</sup> Based on GRADE, two systems have been proposed to evaluate the credibility of results from NMAs,<sup>101,102</sup> but the complexity of the methods and lack of suitable software have limited their wide adoption.<sup>103</sup> CINeMA is based on six domains: within-study bias (referring to the impact of RoB in the included studies), across-studies bias (publication or reporting bias), indirectness (relevance to the research question and transitivity), imprecision (comparing the range of treatment included in the 95% confidence interval with the range of equivalence), heterogeneity (predictive intervals), and incoherence (if estimates from direct and indirect evidence disagree).<sup>111</sup> Judgements across the six domains are then summarised to obtain four levels of confidence for each relative treatment, corresponding to the usual GRADE<sup>82</sup> approach: very low, low, moderate or high.


If possible, we will also perform subgroup analysis (sex, atopic predisposition as a marker of high risk to develop allergic sensitization or allergic disease).

### Transformation into a LSR

This review will continuously evaluate the role of interventions for the prevention in early childhood (ECAP). A living systematic review is a cumulative synthesis that is updated regularly as new evidence becomes available.<sup>86</sup> According to Cochrane’s Living Evidence Network<sup>104</sup> transformation into a LSR is justified when the review question is a particular priority for decision-making, there is an important level of uncertainty in the existing evidence and there is likely to be emerging evidence that will affect the conclusions of the LSR. The review question is of particular priority for decision making because one SR on ECAP<sup>105</sup> is among the Cochrane Priority Reviews and the rate of publications on ECAP interventions has substantially increased over the past years (Figure 1) and is expected to continue so. The level of uncertainty remains an issue. A recent SR on interventions for pregnant or breastfeeding women and/or infants concluded that while dietary avoidance of food allergens, vitamin supplements, fish oil, probiotics, prebiotics, synbiotics, and emollients may have little to no effect on preventing food allergy, the evidence was judged as very uncertain.<sup>43</sup> New evidence is expected to emerge on variations of the induction of tolerance paradigm as paved by previous trials such as EAT<sup>20</sup> or LEAP.<sup>22,106</sup>

An artificial intelligence algorithm for automated searches will be developed in collaboration with information specialists and software engineers. We will gradually incorporate more and more intelligent software tools for eligibility tests, data extraction, and analysis and synthesis presentation as they prove to be reliable in the subsequent update processes.

This evolving search algorithm is to be run at three-monthly intervals and will be set up to notify the author team about studies with a high likelihood of being eligible. At the same interval manual searches will be conducted until the



**Figure 1.** Number of publications derived from Web of Science Core Collection (allergy AND prevention AND childhood) per year since 2002 (performed 26/02/2021).



automated search algorithm is sufficiently reliable. The author team will review the search results, decide upon inclusion, and update the living SR's web version after a new evidence synthesis is deemed necessary every three months. At the same time references of included studies and the corresponding tables and figures will be updated. Every three months the date of each subsequent search, the number of new included studies, and new summary of findings tables will be published on the LSR's website along with an updated plain language summary.

### Discontinuation of the 'living' aspect of the LSR

The LSR will be maintained in its 'living' form until no new evidence is likely to arise and/or so long as the certainty of the evidence for particular ECAP interventions remains unsettled. The necessary future funding will be sought.

### Dissemination of findings

The baseline SR will be published in a peer-reviewed journal and indexed publication platform that allows the linkage of review updates through versioning of the review publication. A plain language summary will be provided on a project website. This website is intended to also contain graphical and other information for health care providers and the public.

For the LSR, we will consider resubmission to the journal in which the baseline SR is published should the direction of the effect on the critical outcomes change or a substantial modification to the evidence's certainty occur.

### Study status

At submission of this protocol the final search strategy is being peer-reviewed.

### Discussion

The prevention of allergy in early childhood is important given the high personal burden and societal costs of allergic diseases. The ECAP evidence landscape has undergone dramatic transformations and this process is likely to continue. As a response to this, a LSR offers the potential to undertake a more timely synthesise of new evidence as it emerges. Long gaps in between updates of SRs may make it harder for guidelines and recommendations to be current and up to date. Users of information, such as parents may be confused if they encounter new evidence that is not part of a trusted guideline. A LSR approach allows us to continuously search the literature and update the evidence-base of existing ECAP interventions resulting in a decreased timespan from evidence accrual to informing clinical practice.

It is also crucially important to assess whether an ECAP intervention is associated with harm. For instance, there is evidence suggesting that early egg white powder introduction at 4 to 6 months of age for the prevention of egg allergy in children from the general population may increase the occurrence of allergy.<sup>108</sup>

Further, a consistent assessment of the certainty of the evidence in line with the GRADE recommendation will be carried out across all included studies. This will allow a high level of concordance between the certainty of the evidence and future guideline recommendations.

Whilst our approach of including all interventions for the prevention of all allergies in early childhood may seem ambitious, none of the individual systematic reviews being undertaken at present (e.g.<sup>33,34</sup>) will be able to conduct a comparison of multiple interventions (network meta-analysis). The latter provides information regarding the relative treatment effect and the ranking order for multiple treatments for a particular outcome irrespective of whether they were conducted in the same or different trials.

### Research, clinical and policy implications

The baseline and subsequent LSR will identify research gaps needing to be addressed by future research. The clinical implications of the LSR will be the provision of up-to-date information that can dynamically inform national, international clinical and public health guidelines and influence the practice of allergists, paediatricians, ENT specialists, other primary care providers and public health authorities in terms of evidence-based ECAP strategies. The provision of a regularly updated plain language summary will benefit parents as they can easily access a widely visible website with information provided in lay terms. The LSR will include research evidence from across the world and we hope will have a multinational benefit.

### Data availability

#### Underlying data

No underlying data are associated with this article.

## Reporting guidelines

Figshare: PRISMA-P checklist for “The evidence for interventions in early childhood allergy prevention – towards a living systematic review: protocol” <https://doi.org/10.6084/m9.figshare.14135450.v1>.<sup>95</sup>

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/) (CC-BY 4.0).

## References

- Loh W, Tang MLK: **The Epidemiology of Food Allergy in the Global Context.** *Int J Environ Res Public Health.* 2018; **15**.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dunn Galvin A, Dubois AEJ, Flokstra-de Blok BMJ, et al.: **The effects of food allergy on quality of life.** *Chem Immunol Allergy.* 2015; **101**: 235–52.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Warren CM, Otto AK, Walkner MM, et al.: **Quality of Life Among Food Allergic Patients and Their Caregivers.** *Curr Allergy Asthma Rep.* 2016; **16**: 38.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dunn Galvin A, Hourihane JO: **Health-related quality of life in food allergy: Impact, correlates, and predictors.** *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 2016; **59**: 841–8.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Deckers IAG, McLean S, Linssen S, Mommers M, et al.: **Investigating international time trends in the incidence and prevalence of atopic eczema 1990-2010: a systematic review of epidemiological studies.** *PLoS One.* 2012; **7**: e39803.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garg N, Silverberg JI: **Epidemiology of childhood atopic dermatitis.** *Clin Dermatol.* 2015; **33**: 281–8.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Odhiambo JA, Williams HC, Clayton TO, et al.: **Global variations in prevalence of eczema symptoms in children from ISAAC Phase Three.** *J Allergy Clin Immunol.* 2009; **124**: 1251–8. e23.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- McNeill G, Tagiyeva N, Aucott L, et al.: **Changes in the prevalence of asthma, eczema and hay fever in pre-pubertal children: a 40-year perspective.** *Paediatr Perinat Epidemiol.* 2009; **23**: 506–12.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Small P, Keith PK, Kim H: **Allergic rhinitis.** *Allergy Asthma Clin Immunol.* 2018; **14**: 51.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Matterne U, Schmitt J, Diepgen TL, et al.: **Children and adolescents health-related quality of life in relation to eczema, asthma and hay fever: results from a population-based cross-sectional study.** *Qual Life Res.* 2011; **20**: 1295–305.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Krawiec M, Fisher H, duToit G, et al.: **Overview of oral tolerance induction for prevention of food allergy - where are we now?** *Allergy.* 2021.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pechlivanis S, von Mutius E: **Effect of Farming on Asthma.** *Acta Med Acad.* 2020; **49**: 144–55.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Roßberg S, Keller T, Icke K, et al.: **Orally applied bacterial lysate in infants at risk for atopy does not prevent atopic dermatitis, allergic rhinitis, asthma or allergic sensitization at school age: Follow-up of a randomized trial.** *Allergy.* 2020.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Strachan DP: **Hay fever, hygiene, and household size.** *BMJ.* 1989; **299**: 1259–60.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lambrecht BN, Hammad H: **The immunology of the allergy epidemic and the hygiene hypothesis.** *Nat Immunol.* 2017; **18**: 1076–83.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ege MJ: **The Hygiene Hypothesis in the Age of the Microbiome.** *Ann Am Thorac Soc.* 2017; **14**: S348–S353.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bach J-F: **The hygiene hypothesis in autoimmunity: the role of pathogens and commensals.** *Nat Rev Immunol.* 2018; **18**: 105–20.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lack G: **Update on risk factors for food allergy.** *J Allergy Clin Immunol.* 2012; **129**: 1187–97.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Prescott SL, Bouygue GR, Videky D, et al.: **Avoidance or exposure to foods in prevention and treatment of food allergy?** *Curr Opin Allergy Clin Immunol.* 2010; **10**: 258–66.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Perkin MR, Logan K, Bahnsen HT, et al.: **Efficacy of the Enquiring About Tolerance (EAT) study among infants at high risk of developing food allergy.** *J Allergy Clin Immunol.* 2019; **144**: 1606–1614. e2.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tan JW, Valerio C, Barnes EH, et al.: **A randomized trial of egg introduction from 4 months of age in infants at risk for egg allergy.** *J Allergy Clin Immunol.* 2017; **139**: 1621–1628. e8.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Du Toit G, Roberts G, Sayre PH, et al.: **Randomized trial of peanut consumption in infants at risk for peanut allergy.** *N Engl J Med.* 2015; **372**: 803–13.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Du Toit G, Foong R-X, Lack G: **Prevention of food allergy - Early dietary interventions.** *Allergol Int.* 2016; **65**: 370–7.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Camargo CA JR, Clark S, Kaplan MS, et al.: **Regional differences in EpiPen prescriptions in the United States: the potential role of vitamin D.** *J Allergy Clin Immunol.* 2007; **120**: 131–6.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Giannetti A, Bernardini L, Cangemi J, et al.: **Role of Vitamin D in Prevention of Food Allergy in Infants.** *Front Pediatr.* 2020; **8**: 447.  
[PubMed Abstract](#) | [Free Full Text](#) | [Free Full Text](#)
- Al-Saud B, Sigurdardottir ST: **Early Introduction of Egg and the Development of Egg Allergy in Children: A Systematic Review and Meta-Analysis.** 2018; **177**: 350–9.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Azad MB, Coneys JG, Kozyrskij AL, et al.: **Probiotic supplementation during pregnancy or infancy for the prevention of asthma and wheeze: systematic review and meta-analysis.** 2020; **347**: f6471.  
[PubMed Abstract](#) | [Free Full Text](#) | [Free Full Text](#)
- Barfield A, Brown H, Pernel P, et al.: **Effectiveness of Emollient Therapy in Pediatric Patients With Atopic Dermatitis.** *J Dermatol Nurse Assoc.* 2017; **9**: 123–8.  
[Publisher Full Text](#)
- Best KP, Gold M, Kennedy D, et al.: **Omega-3 long-chain PUFA intake during pregnancy and allergic disease outcomes in the offspring: a systematic review and meta-analysis of observational studies and randomized controlled trials.** *Am J Clin Nutr.* 2016; **103**: 128–43.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Boyle RJ, Ierodiakonou D, Khan T, et al.: **Hydrolysed formula and risk of allergic or autoimmune disease: systematic review and meta-analysis.** *BMJ.* 2016; **352**: i974.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bremner SF, Simpson EL: **Dust mite avoidance for the primary prevention of atopic dermatitis: A systematic review and meta-analysis.** *Pediatr Allergy Immunol.* 2015; **26**: 646–54.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Burgess JA, Dharmage SC, Allen K, et al.: **Age at introduction to complementary solid food and food allergy and sensitization: A systematic review and meta-analysis.** *Clin Exp Allergy.* 2019; **49**: 754–69.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Alexander DD, Schmitt DF, Tran NL, et al.: **Partially hydrolyzed 100% whey protein infant formula and atopic dermatitis risk reduction: a systematic review of the literature.** *Nutr Rev.* 2010;

- 68: 232–45.  
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Anandan C, Nurmatov U, Sheikh A: **Omega 3 and 6 oils for primary prevention of allergic disease: systematic review and meta-analysis.** *Allergy*. 2009; **64**: 840–8.  
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Arnoldussen DL, Linehan M, Sheikh A: **BCG vaccination and allergy: a systematic review and meta-analysis.** *J Allergy Clin Immunol*. 2011; **127**: 246–53, 253.e1–21.  
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Baumgartner M, Brown CA, Exl B-M, *et al.*: **Controlled trials investigating the use of one partially hydrolyzed whey formula for dietary prevention of atopic manifestations until 60 months of age: An overview using meta-analytical techniques.** *Nutri Res*. 1998; **18**: 1425–42.  
[Publisher Full Text](#) | [Reference Source](#)
37. Chang Y-S, Trivedi MK, Jha A, *et al.*: **Synbiotics for Prevention and Treatment of Atopic Dermatitis: A Meta-analysis of Randomized Clinical Trials.** *JAMA Pediatr*. 2016; **170**: 236–42.  
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Chmielewska A, Piescik-Lech M, Shamir R, *et al.*: **Systematic review: Early infant feeding practices and the risk of wheat allergy; 9.**  
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Cooke A, Bedwell C, Campbell M, *et al.*: **Skin care for healthy babies at term: A systematic review of the evidence.**  
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Cuello-García C, Fiocchi A, Pawankar R, *et al.*: **Prebiotics for the prevention of allergies: A systematic review and meta-analysis of randomized controlled trials.** *Clin Exp Allergy*. 2017; **47**: 1468–77.  
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Cuello-García CA, Brozek JL, Fiocchi A, *et al.*: **Probiotics for the prevention of allergy: A systematic review and meta-analysis of randomized controlled trials.** *J Allergy Clin Immunol*. 2015; **136**.  
[PubMed Abstract](#) | [Publisher Full Text](#)
42. de Silva D, Geromi M, Halken S, *et al.*: **Primary prevention of food allergy in children and adults: systematic review.** *Allergy*. 2014; **69**: 581–9.  
[PubMed Abstract](#) | [Publisher Full Text](#)
43. de Silva D, Halken S, Singh C, *et al.*: **Preventing food allergy in infancy and childhood: Systematic review of randomised controlled trials.** *Pediatr Allergy Immunol*. 2020; **31**: 813–26.  
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Doege K, Grajecki D, Zyriax B-C, *et al.*: **Impact of maternal supplementation with probiotics during pregnancy on atopic eczema in childhood—a meta-analysis.** *Br J Nutr*. 2012; **107**: 1–6.  
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Dogaru CM, Nyffenegger D, Pescatore AM, *et al.*: **Breastfeeding and childhood asthma: systematic review and meta-analysis.** *Am J Epidemiol*. 2014; **179**: 1153–67.  
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Garcia-Larsen V, Ierodiakonou D, Jarrold K, *et al.*: **Diet during pregnancy and infancy and risk of allergic or autoimmune disease: A systematic review and meta-analysis.** *PLoS Med*. 2018; **15**: e1002507.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Gunaratne AW, Makrides M, Collins CT: **Maternal prenatal and/or postnatal n-3 long chain polyunsaturated fatty acids (LCPUFA) supplementation for preventing allergies in early childhood.** *Cochrane Database Syst Rev*. 2015; **7**: CD010085.  
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Güngör D, Nadaud P, LaPergola CC, *et al.*: **Infant milk-feeding practices and food allergies, allergic rhinitis, atopic dermatitis, and asthma throughout the life span: a systematic review.** 2019; **109**: 772S–799S.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Hays T, Wood RA: **A systematic review of the role of hydrolyzed infant formulas in allergy prevention.** *Arch Pediatr Adolesc Med*. 2005; **159**: 810–6.  
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Hosseini B, Berthon BS, Wark P, *et al.*: **Effects of Fruit and Vegetable Consumption on Risk of Asthma, Wheezing and Immune Responses: A Systematic Review and Meta-Analysis.** *Nutrients*. 2017; **9**.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Ierodiakonou D, Garcia-Larsen V, Logan A, *et al.*: **Timing of Allergenic Food Introduction to the Infant Diet and Risk of Allergic or Autoimmune Disease: A Systematic Review and Meta-analysis.** *JAMA*. 2016; **316**: 1181–92.  
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Kramer MS, Kakuma R: **Maternal dietary antigen avoidance during pregnancy or lactation, or both, for preventing or treating atopic disease in the child.** *Cochrane Database Syst Rev*. 2012; **12**.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Klemens CM, Berman DR, Mozurkewich EL: **The effect of perinatal omega-3 fatty acid supplementation on inflammatory markers and allergic diseases: a systematic review.** *BJOG*. 2011; **118**: 916–25.  
[PubMed Abstract](#) | [Publisher Full Text](#)
54. Kremmyda L-S, Vlachava M, Noakes PS, *et al.*: **Atopy risk in infants and children in relation to early exposure to fish, oily fish, or long-chain omega-3 fatty acids: a systematic review.** *Clin Rev Allergy Immunol*. 2011; **41**: 36–66.  
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Kristiansen M, Dhimi S, Netuveli G, *et al.*: **Allergen immunotherapy for the prevention of allergy: A systematic review and meta-analysis.** **28**: 18–29.  
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Kuang L, Jiang Y: **Effect of probiotic supplementation in pregnant women: a meta-analysis of randomised controlled trials.** *Br J Nutr*. 2020; **123**: 870–80.  
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Larson K, McLaughlin J, Stonehouse M, *et al.*: **Introducing Allergenic Food into Infants' Diets: Systematic Review, 2.**  
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Macdonald C, Sternberg A, Hunter P: **A systematic review and meta-analysis of interventions used to reduce exposure to house dust and their effect on the development and severity of asthma.** *Cien Saude Colet*. 2008; **13**: 1907–15.  
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Mansfield JA, Bergin SW, Cooper JR, *et al.*: **Comparative probiotic strain efficacy in the prevention of eczema in infants and children: a systematic review and meta-analysis.** *Mil Med*. 2014; **179**: 580–92.  
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Netting MJ, Middleton PF, Makrides M: **Does maternal diet during pregnancy and lactation affect outcomes in offspring? A systematic review of food-based approaches.** *Nutrition*. 2014; **11-12**: 1225–41.  
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Newberry SJ, Chung M, Booth M, *et al.*: **Omega-3 Fatty Acids and Maternal and Child Health: An Updated Systematic Review.** *Evid Rep Technol Assess*. 2016: 1–826.  
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Obbagy JE, English LK, Wong YP, *et al.*: **Complementary feeding and food allergy, atopic dermatitis/eczema, asthma, and allergic rhinitis: a systematic review.** *Suppl*. 7.  
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Panduru M, Panduru NM, Salavastru CM, *et al.*: **Probiotics and primary prevention of atopic dermatitis: a meta-analysis of randomized controlled studies.** *J Eur Acad Dermatol Venereol*. 2015; **29**: 232–42.  
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Osborn DA, Sinn JK, Jones LJ: **Infant formulas containing hydrolysed protein for prevention of allergic disease.** *Cochrane Database Syst Rev*. 2018; **10**.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Pelucchi C, Chatenoud L, Turati F, *et al.*: **Probiotics supplementation during pregnancy or infancy for the prevention of atopic dermatitis: a meta-analysis.** *Epidemiology (Cambridge, Mass.)* 2012; **23**: 402–14.  
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Seyedrezaazadeh E, Moghaddam MP, Ansarin K, *et al.*: **Fruit and vegetable intake and risk of wheezing and asthma: a systematic review and meta-analysis.** *Nutr Rev*. 2014; **72**: 411–28.  
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Schindler T, Sinn JK, Osborn DA: **Polyunsaturated fatty acid supplementation in infancy for the prevention of allergy.** *Cochrane Database Syst Rev*. 2016; **10**.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Szajewska H, Horvath A: **A partially hydrolyzed 100% whey formula and the risk of eczema and any allergy: an updated meta-analysis.** *World Allergy Organ J*. 2017; **10**: 27.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Szajewska H, Horvath A: **Lactobacillus rhamnosus GG in the Primary Prevention of Eczema in Children: A Systematic Review and Meta-Analysis.** *Nutrients*. 2018; **10**: pii: E1319.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Thompson RL, Miles LM, Lunn J, *et al.*: **Peanut sensitisation and allergy: influence of early life exposure to peanuts.** 2010; **103**: 1278–86.  
[PubMed Abstract](#) | [Publisher Full Text](#)

71. Vahdaninia M, Mackenzie H, Dean T, *et al.*: **omega-3 LCPUFA supplementation during pregnancy and risk of allergic outcomes or sensitization in offspring: A systematic review and meta-analysis.** *Ann Allergy Asthma Immunol.* 2019; **122**.  
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Vahdaninia M, Mackenzie H, Helps S, *et al.*: **Prenatal Intake of Vitamins and Allergic Outcomes in the Offspring: A Systematic Review and Meta-Analysis.** *J Allergy Clin Immunol Pract.* 2017; **5**: 771–8.  
[PubMed Abstract](#) | [Publisher Full Text](#)
73. Venter C, Agostoni C, Arshad SH, *et al.*: **Dietary factors during pregnancy and atopic outcomes in childhood: a systematic review from the European Academy of Allergy and Clinical Immunology.** *Pediatr Allergy Immunol.* 2020.  
[PubMed Abstract](#) | [Publisher Full Text](#)
74. Waidyatillake NT, Dharmage SC, Allen KJ, *et al.*: **Association between the age of solid food introduction and eczema: A systematic review and a meta-analysis.** *Clin Exp Allergy.* 2018; **48**: 1000–15.  
[PubMed Abstract](#) | [Publisher Full Text](#)
75. Wei X, Jiang P, Liu J, *et al.*: **Association between probiotic supplementation and asthma incidence in infants: a meta-analysis of randomized controlled trials.** *J Asthma.* 2020; **57**: 167–78.  
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Yang H, Xun P, He K: **Fish and fish oil intake in relation to risk of asthma: a systematic review and meta-analysis.** *PLoS One.* 2013; **8**: e80048.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
77. Zuccotti G, Meneghin F, Aceti A, *et al.*: **Probiotics for prevention of atopic diseases in infants: systematic review and meta-analysis.** 2015; **70**: 1356–71.  
[PubMed Abstract](#) | [Publisher Full Text](#)
78. Zhang G-Q, Liu B, Li J, *et al.*: **Fish intake during pregnancy or infancy and allergic outcomes in children: A systematic review and meta-analysis; 2.**  
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Yuan M, Tan M, Moore D, *et al.*: **Timing of Cow's Milk or Cow's Milk Formula Introduction to the Infant Diet and Atopic Risk in Children: a Systematic Review and Meta-analysis.** *Clin Rev Allergy Immunol.* 2019.  
[PubMed Abstract](#) | [Publisher Full Text](#)
80. Yepes-Nunez JJ, Brozek JL, Fiocchi A, *et al.*: **Vitamin D supplementation in primary allergy prevention: Systematic review of randomized and non-randomized studies.** 2018; **73**.  
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Kelleher MM, Cro S, Cornelius V, *et al.*: **Skin care interventions in infants for preventing eczema and food allergy.** *Cochrane Database Syst Rev.* 2021; **2**: CD013534.  
[PubMed Abstract](#) | [Publisher Full Text](#)
82. Schünemann H, Brozek J, Guyatt G, *et al.*, editors. **GRADE handbook for grading quality of evidence and strength of recommendations.** The GRADE Working Group; 2013.  
[Reference Source](#)
83. Wells G, Shea B, O'Connell D, *et al.*: **The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in metaanalyses.** Cited 8 Jul 2020.  
[Reference Source](#)
84. National Institute for Health and Care Excellence: **Developing NICE guidelines: the manual.** Cited 8 Jul 2020.  
[Reference Source](#)
85. Ebell MH, Siwek J, Weiss BD, *et al.*: **Strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature.** *Am Fam Physician.* 2004; **69**: 548–56.  
[PubMed Abstract](#)
86. Elliott JH, Synnot A, Turner T, *et al.*: **Living systematic review: 1. Introduction—the why, what, when, and how.** *J Clin Epidemiol.* 2017; **91**: 23–30.  
[PubMed Abstract](#) | [Publisher Full Text](#)
87. Mills EJ, Thorlund K, Ioannidis JPA: **Demystifying trial networks and network meta-analysis.** *BMJ.* 2013; **346**: f2914.  
[PubMed Abstract](#) | [Publisher Full Text](#)
88. Chaimani A, Caldwell DM, Li T, *et al.*: **Chapter 11: Undertaking network meta-analyses.** In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions version 6.1 (updated September 2020)*. Cochrane; 2020 Cited 20 Jan 2021.  
[Reference Source](#)
89. Rouse B, Chaimani A, Li T: **Network meta-analysis: an introduction for clinicians.** *Intern Emerg Med.* 2017; **12**: 103–11.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
90. Higgins JPT, Thomas J, Chandler J, *et al.*: *Cochrane Handbook for Systematic Reviews of Interventions.* 2nd ed. Chichester (UK): John Wiley & Sons; 2019.
91. Moher D, Liberati A, Tetzlaff J, *et al.*: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement.** *Journal of clinical epidemiology.* 2009; **62**: 1006–12.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. Hutton B, Salanti G, Caldwell DM, *et al.*: **The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations.** *Ann Intern Med.* 2015; **162**: 777–84.  
[PubMed Abstract](#) | [Publisher Full Text](#)
93. National Institute for Health Research: **PROSPERO International prospective register of systematic reviews.** Cited 8 Jul 2020.  
[Reference Source](#)
94. Moher D, Shamseer L, Clarke M, *et al.*: **Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement.** *Syst Rev.* 2015; **4**: 1.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
95. Mattrer U, Tischer C, Wang J, *et al.*: **Prisma-P checklist; 2021.**  
[Reference Source](#)
96. Thomas J, Graziosi S, Brunton J, *et al.*: **EPPI-Reviewer: advanced software for systematic reviews, maps and evidence synthesis: EPPI-Centre Software.** UCL Social Research Institute; 2020.
97. Shea BJ, Reeves BC, Wells G, *et al.*: **AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both.** *BMJ.* 2017; **358**: j4008.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
98. Simmonds M, Salanti G, McKenzie J, *et al.*: **Living systematic reviews: 3. Statistical methods for updating meta-analyses.** *J Clin Epidemiol.* 2017; **91**: 38–46.  
[PubMed Abstract](#) | [Publisher Full Text](#)
99. Institute of Social and Preventive Medicine, University of Bern: **CINeMA: Confidence in Network Meta-Analysis [Software].** 2017.  
[Reference Source](#)
100. World Health Organisation: **WHO handbook for guideline development.** 2012 Cited 20 Jan 2021.  
[Reference Source](#)
101. Puhan MA, Schünemann HJ, Murad MH, *et al.*: **A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis.** *BMJ.* 2014; **349**: g5630.  
[PubMed Abstract](#) | [Publisher Full Text](#)
102. Salanti G, Del Giovane C, Chaimani A, *et al.*: **Evaluating the quality of evidence from a network meta-analysis.** *PLoS One.* 2014; **9**: e99682.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
103. Nikolakopoulou A, Higgins JPT, Papakonstantinou T, *et al.*: **CINeMA: An approach for assessing confidence in the results of a network meta-analysis.** *PLoS Med.* 2020; **17**: e1003082.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
104. Brooker J, Synnot A, McDonald S, *et al.*: **Guidance for the production and publication of Cochrane living systematic reviews: Cochrane Reviews in living mode.** Version December 2019 Cited 25 Feb 2021.  
[Reference Source](#)
105. Best CM, Xu J, Patchen BK, *et al.*: **Vitamin D supplementation in pregnant or breastfeeding women or young children for preventing asthma.** *Cochrane Database Syst Rev.* 2019.  
[Publisher Full Text](#) | [Free Full Text](#)
106. Du Toit G, Sayre PH, Roberts G, *et al.*: **Allergen specificity of early peanut consumption and effect on development of allergic disease in the Learning Early About Peanut Allergy study cohort.** *J Allergy Clin Immunol.* 2018; **141**: 1343–53.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
107. Schmidt L, Olorisade BK, McGuinness LA, *et al.*: **Data extraction methods for systematic review (semi)automation: A living review protocol.** *F1000Res* 2020; **9**: 210.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
108. Bellach J, Schwarz V, Ahrens B, *et al.*: **Randomized placebo-controlled trial of hen's egg consumption for primary prevention in infants.** *J Allergy Clin Immunol.* 2017; **139**: 1591–1599. e2.  
[PubMed Abstract](#) | [Publisher Full Text](#)
109. Litonjua AA, Carey VJ, Laranjo N, *et al.*: **Effect of Prenatal Supplementation With Vitamin D on Asthma or Recurrent Wheezing in Offspring by Age 3 Years: The VDAART Randomized Clinical Trial.** *JAMA.* 2016; **315**: 362–70.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

110. Cabana MD, McKean M, Caughey AB, *et al.*: **Early Probiotic Supplementation for Eczema and Asthma Prevention: A Randomized Controlled Trial.** *Pediatrics*. 2017; **140**.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
111. Gore C, Custovic A, Tannock GW, *et al.*: **Treatment and secondary prevention effects of the probiotics *Lactobacillus paracasei* or *Bifidobacterium lactis* on early infant eczema: randomized controlled trial with follow-up until age 3 years.** *Clin Exp Allergy*. 2012; **42**: 112–22.  
[PubMed Abstract](#) | [Publisher Full Text](#)
112. Zeiger RS, Heller S: **The development and prediction of atopy in high-risk children: follow-up at age seven years in a prospective randomized study of combined maternal and infant food allergen avoidance.** *J Allergy Clin Immunol*. 1995; **95**: 1179–90.  
[PubMed Abstract](#) | [Publisher Full Text](#)
113. Tachimoto H, Imanari E, Mezawa H, *et al.*: **Effect of Avoiding Cow's Milk Formula at Birth on Prevention of Asthma or Recurrent Wheeze Among Young Children: Extended Follow-up From the ABC Randomized Clinical Trial.** *JAMA Netw Open*. 2020; **3**: e2018534.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
114. Yamamoto-Hanada K, Kobayashi T, Williams HC, *et al.*: **Early aggressive intervention for infantile atopic dermatitis to prevent development of food allergy: a multicenter, investigator-blinded, randomized, parallel group controlled trial (PACI Study)-protocol for a randomized controlled trial.** *Clin Transl Allergy*. 2018; **8**: 47.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
115. Zolkipli Z, Roberts G, Cornelius V, *et al.*: **Randomized controlled trial of primary prevention of atopy using house dust mite allergen oral immunotherapy in early childhood.** *J Allergy Clin Immunol* 2015; **136**: 1541–1547. e11.  
[PubMed Abstract](#) | [Publisher Full Text](#)
116. Valovirta E, Petersen TH, Piotrowska T, *et al.*: **Results from the 5-year SQ grass sublingual immunotherapy tablet asthma prevention (GAP) trial in children with grass pollen allergy.** *J Allergy Clin Immunol*. 2018; **141**: 529–538. e13.  
[PubMed Abstract](#) | [Publisher Full Text](#)
117. Steenhuis TJ, van Aalderen WMC, Bloksma N, *et al.*: **Bacille-Calmette-Guerin vaccination and the development of allergic disease in children: a randomized, prospective, single-blind study.** *Clin Exp Allergy*. 2008; **38**: 79–85.  
[PubMed Abstract](#) | [Publisher Full Text](#)
118. Thøstesen LM, Kjaer HF, Pihl GT, *et al.*: **Neonatal BCG has no effect on allergic sensitization and suspected food allergy until 13 months.** *Pediatr Allergy Immunol*. 2017; **28**: 588–96.  
[PubMed Abstract](#) | [Publisher Full Text](#)
119. Warner JO: **A double-blinded, randomized, placebo-controlled trial of cetirizine in preventing the onset of asthma in children with atopic dermatitis: 18 months' treatment and 18 months' posttreatment follow-up.** *J Allergy Clin Immunol*. 2001; **108**: 929–37.  
[PubMed Abstract](#) | [Publisher Full Text](#)

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact [research@f1000.com](mailto:research@f1000.com)

**F1000Research**