

# Sacubitril/valsartan for the treatment of non-obstructive hypertrophic cardiomyopathy: An open label randomized controlled trial (SILICOFCM)

Lazar Velicki<sup>1,2</sup>, Dejana Popovic<sup>3</sup>, Nduka C. Okwose<sup>4,5</sup> , Andrej Preveden<sup>1,2</sup>, Milorad Tesic<sup>3</sup>, Maria Tafelmeier<sup>6</sup>, Sarah J. Charman<sup>4</sup>, Fausto Barlocco<sup>7</sup>, Guy A. MacGowan<sup>4,8</sup>, Petar M. Seferovic<sup>3</sup>, Nenad Filipovic<sup>9,10</sup>, Arsen Ristic<sup>3</sup>, Iacopo Olivetto<sup>7</sup>, Lars S. Maier<sup>6</sup>, and Djordje G. Jakovljevic<sup>4,5,8\*</sup> , on behalf of SILICOFCM Investigators<sup>†</sup>

<sup>1</sup>Faculty of Medicine, University of Novi Sad, Novi Sad, Serbia; <sup>2</sup>Institute of Cardiovascular Diseases of Vojvodina, Sremska Kamenica, Serbia; <sup>3</sup>Cardiology Department, Faculties of Medicine and Pharmacy, University of Belgrade, Serbia; <sup>4</sup>Cardiovascular Research, Translational and Clinical Research and Biosciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK; <sup>5</sup>Research Centre for Health and Life Sciences, Institute of Health and Wellbeing, Faculty of Health and Life Science, Coventry University, Coventry, UK; <sup>6</sup>Department of Internal Medicine II, University Medical Centre Regensburg, Regensburg, Germany; <sup>7</sup>Careggi University Hospital, University of Florence, Florence, Italy; <sup>8</sup>Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK; <sup>9</sup>Bioengineering Research and Development Center, BioIRC, Kragujevac, Serbia; and <sup>10</sup>Faculty of Engineering, University of Kragujevac, Kragujevac, Serbia

Received 11 May 2023; revised 12 April 2024; accepted 30 April 2024; online publish-ahead-of-print 27 May 2024

## Aim

Sacubitril/valsartan treatment reduces mortality and hospitalizations in heart failure with reduced ejection fraction but has limited application in hypertrophic cardiomyopathy (HCM). The aim of this study was to evaluate the effect of sacubitril/valsartan on peak oxygen consumption ( $\text{VO}_2$ ) in patients with non-obstructive HCM.

## Methods and results

This is a phase II, randomized, open-label multicentre study that enrolled adult patients with symptomatic non-obstructive HCM (New York Heart Association class I–III) who were randomly assigned (2:1) to receive sacubitril/valsartan (target dose 97/103 mg) or control for 16 weeks. The primary endpoint was a change in peak  $\text{VO}_2$ . Secondary endpoints included echocardiographic measures of cardiac structure and function, natriuretic peptides and other cardiac biomarkers, and Minnesota Living with Heart Failure quality of life. Between May 2018 and October 2021, 354 patients were screened for eligibility, 115 patients (mean age 58 years, 37% female) met the study inclusion criteria and were randomly assigned to sacubitril/valsartan ( $n = 79$ ) or control ( $n = 36$ ). At 16 weeks, there was no significant change in peak  $\text{VO}_2$  from baseline in the sacubitril/valsartan (15.3 [4.3] vs. 15.9 [4.3] ml/kg/min,  $p = 0.13$ ) or control group ( $p = 0.47$ ). No clinically significant changes were found in blood pressure, cardiac structure and function, plasma biomarkers, or quality of life.

## Conclusion

In patients with HCM, a 16-week treatment with sacubitril/valsartan was well tolerated but had no effect on exercise capacity, cardiac structure, or function.

## Keywords

Hypertrophic cardiomyopathy • Exercise capacity • Sacubitril/valsartan

\*Corresponding author. Research Centre for Health and Life Sciences, Institute of Health and Wellbeing, Faculty of Health and Life Sciences, Coventry University, 20 Whitefriars Street, Coventry CV1 2DS, UK. Tel: +44 (0) 24 7765 8075. Email: djordje.jakovljevic@coventry.ac.uk

<sup>†</sup>SILICOFCM Investigators are listed in Appendix A.

## Introduction

Hypertrophic cardiomyopathy (HCM) is one of the most common genetic heart diseases characterized by left ventricular (LV) hypertrophy<sup>1</sup> with a prevalence ranging from 1:200 to 1:500.<sup>2</sup> It is caused by a pathological mutation of the cardiac sarcomeric protein genes and transmitted in an autosomal dominant manner.<sup>3</sup> Core pathophysiological features of HCM include increase in myocardial wall thickness, diastolic abnormalities, and dynamic LV outflow tract obstruction.<sup>4–6</sup>

There is limited evidence on the reliability of pharmacological treatments to alter disease progression in HCM. The VANISH trial showed significant improvements in cardiac structure and function in early stage HCM after 24-month administration of the angiotensin II receptor blocker valsartan, indicating an opportunity to attenuate disease progression.<sup>7</sup> However, most therapies focus on symptoms relief and the prophylactic implantation of cardioverter-defibrillators to reduce the risk of sudden cardiac death.<sup>6,8</sup> Moreover, hypotension and a reduction in LV ejection fraction (LVEF) associated with some therapies undermine the benefits of continued administration.<sup>9</sup>

Sacubitril/valsartan, a first-in-class angiotensin receptor–neprilysin inhibitor, was approved for the treatment of heart failure with reduced ejection fraction based on its superiority over angiotensin-converting enzyme inhibitors (ACEi) in reducing cardiovascular mortality and heart failure hospitalizations.<sup>10</sup> Neprilysin degrades biologically active atrial and B-type natriuretic peptides, secreted by cardiomyocytes in response to increased myocardial wall tension and thus may play a cardio-protective role from volume and pressure overload.<sup>11</sup> Neprilysin inhibition increases generation of myocardial cyclic guanosine monophosphate, which improves myocardial relaxation and reduces hypertrophy.<sup>11</sup> There is evidence that sacubitril/valsartan improves myocardial morphology and physiology in heart failure<sup>12–15</sup> and can enhance exercise capacity.<sup>16</sup> However, the effects of sacubitril/valsartan on clinical and physiological phenotype in patients with HCM have not yet been evaluated. The aim of the present study was to determine the effect of sacubitril/valsartan on functional capacity, cardiac structure and function, cardiac biomarkers, and quality of life in patients with non-obstructive HCM. To the best of our knowledge this is the first study in human participants to report on the effect of sacubitril/valsartan on clinical phenotype in HCM.

## Methods

### Study design

This was a prospective, multicentre, open-label, randomized, controlled phase II clinical trial which was registered with [ClinicalTrials.gov](https://clinicaltrials.gov) (NCT03832660), and protocol published previously.<sup>17</sup> Data were collected between May 2018 and February 2022. The inclusion and exclusion criteria prioritized safety and included a patient population which was reflective of real-world non-obstructive HCM. Eligible patients were aged  $\geq 18$  years with a diagnosis of non-obstructive HCM (LV hypertrophy with maximal LV wall thickness of  $\geq 15$  mm [or  $\geq 13$  mm if familial HCM]); LVEF  $\geq 55\%$ ; and New York Heart Association (NYHA)


class I–III. Patients had to be able to safely perform maximal cardiopulmonary exercise testing (CPET) and be willing/able to return for follow-up visit. Key exclusion criteria included less than 3 months post-septal reduction therapy (surgery or catheter-based intervention); resting LV outflow tract gradient  $>30$  mmHg; renal insufficiency with a glomerular filtration rate  $<30$  ml/min/1.73 m<sup>2</sup>; life expectancy  $<12$  months; history of exercise-induced syncope or sustained ventricular arrhythmias; concomitant treatment with aliskiren-containing drugs; history of ACEi- or angiotensin receptor blocker (ARB)-induced angioedema or history of hereditary or idiopathic angioedema; evidence of hepatic disease; history of malignancy of any organ system (other than localized basal or squamous cell carcinoma of the skin or localized prostate cancer), treated or untreated, within the past 2 years, regardless of whether there was evidence of local recurrence or metastases; life-threatening or uncontrolled dysrhythmia, including symptomatic or sustained ventricular tachycardia and atrial fibrillation or flutter with a resting ventricular rate  $>110$  bpm. The study was in accordance with the Declaration of Helsinki and Good Clinical Practice guidelines and was approved by the Research Ethics Committees and/or Institutional Review Boards of the study sites. All patients provided written informed consent.

### Study procedures

All patients underwent baseline assessments which included medical history, physical examination and anthropometrics, serum biomarker analysis, genetic testing (if not performed earlier), 12-lead electrocardiogram, transthoracic echocardiography, quality of life assessment and CPET. Biomarkers of cardio-metabolic function were evaluated using venous blood. Echocardiographic images were acquired in the standard parasternal (long-axis, short-axis) and apical (apical 4-chamber, apical 2-chamber, and apical long-axis) views, for which three cardiac cycles were recorded. Parasternal short-axis views were acquired at three levels: basal (at mitral valve level), mid-papillary, and apical (minimum cavity distal to papillary muscle level). LV diameter and fractional shortening, interventricular septum, and LV posterior wall thickness, and left atrial dimensions were measured using M-mode and two-dimensional echocardiography according to current recommendations.<sup>18</sup> Peak velocity of the LV outflow tract was recorded from the apical 5-chamber view by pulse Doppler, and pressure gradient was calculated. The sites and maximal extent of ventricular hypertrophy were assessed and measured in end-diastole. LV volumes and LVEF and right ventricular systolic function were assessed from the apical 4-chamber view, using the biplane method of discs. Left atrial volumes were measured in systole just before the mitral valve opening, using a monoplane area–length method. Transmitral Doppler imaging was used to assess diastolic function.

Patients completed an individualized symptom-limited graded CPET using cycle ergometer, according to established guidelines.<sup>19</sup> Pharmacologic therapy was maintained during CPET. Exercise commenced with 3 min of unloaded cycling followed by increased workloads of 10 W/min until maximum exertion was achieved. Bike cadence was maintained at 60–70 rpm throughout the test.

Simultaneous 12-lead electrocardiography and an automated blood pressure were recorded. Oxygen consumption (VO<sub>2</sub>), minute ventilation (VE), carbon dioxide production (VCO<sub>2</sub>), and other CPET variables were acquired breath by breath, and expressed in 10 s intervals. Peak VO<sub>2</sub> and peak respiratory exchange ratio (RER) were expressed as the highest 10 s averaged sample obtained during the last 20 s of testing. VE and VCO<sub>2</sub> values, acquired from the beginning of exercise to peak, were input into spreadsheet software (Microsoft Excel, Microsoft



Corp., Bellevue, WA, USA) to calculate the  $VE/VCO_2$  slope via least squares linear regression ( $y = mx + b$ ,  $m = \text{slope}$ ). Test termination criteria consisted of symptoms (i.e. dyspnoea and/or fatigue), ventricular tachycardia,  $>2$  mm of horizontal or downsloping ST-segment depression, a drop of systolic blood pressure  $>20$  mmHg during progressive exercise, or  $RER \geq 1.10$ .

## Randomization

Patients were randomly assigned (2:1) using computer-generated numbers to receive orally administered sacubitril/valsartan or control group (optimal standard therapy). Randomization was stratified by functional capacity, NYHA class, current beta-blocker use (yes or no).

## Pharmacological intervention

Patients allocated to the pharmacological intervention group began an initial dose of 24/26 mg or 49/51 mg sacubitril/valsartan, followed by up-titration every 14 days until the target dose of 97/103 mg twice daily was reached. Sacubitril/valsartan was administered orally for 16 weeks. Patients previously on ACEi or ARB therapy had a 36 h washout period before initiation of sacubitril/valsartan to reduce the risk of angioedema. Doses were adjusted based on overall safety and tolerability. Hypotension was defined as investigator-reported hypotension with a systolic blood pressure  $<100$  mmHg. Pre-specified

criteria for temporary discontinuation of study drug, including LVEF  $<50\%$ , have been previously described.<sup>17</sup>

## Optimal standard therapy (control group)

Patients in the control group received optimal standard therapy for HCM according to current guidelines.<sup>8</sup> All patients were asked to return for re-evaluation 16 weeks after enrolment and underwent the same procedures and assessments as at the baseline visit.

## Outcomes

The primary study endpoint was a between-group change in exercise tolerance (i.e. peak  $VO_2$ ) from baseline to week 16. Secondary endpoints were between-group change from baseline to follow-up visit in (a) indices of cardiorespiratory fitness, (b) quality of life measures (i.e. Minnesota Living with Heart Failure Questionnaire [MLHFQ] [range, 0–105; higher scores indicate worse quality of life; minimal clinically important difference, 5 points]), (c) cardiac remodelling measures from echocardiography (i.e. magnitude or distribution of cardiac hypertrophy, degree of LV outflow obstruction, LV chamber dimensions, systolic and diastolic function, and LV wall motion), (d) serum concentrations of creatine kinase, creatine kinase-myocardial band, and N-terminal pro-B-type natriuretic peptide (NT-proBNP), low-density lipoprotein,

high-density lipoprotein, total cholesterol, electrolytes. All assessments for secondary endpoints were controlled for type I error in hierarchical order upon achieving significance in the primary endpoint (with two-tailed  $p < 0.05$  required to proceed). Pre-specified safety endpoints included frequency and severity of treatment-emergent adverse events and serious adverse events.

## Statistical analysis

Existing literature suggests that between-group change of 1.27 ml/kg/min in peak  $\text{VO}_2$  is clinically significant.<sup>20</sup> Assuming similar effects, 120 patients were required on a 2:1 randomization basis, to achieve 90% power to detect a significant difference in peak  $\text{VO}_2$  between groups, at a two-tailed significance level of  $\leq 0.05$ .

Statistical analysis was performed using SPSS (version 28 IBM). Data are presented as mean and standard deviation. Individual changes in outcome measures from baseline to follow-up were determined by either a paired *t*-test for normally distributed data or a Wilcoxon signed-rank test for non-normally distributed data. Between-group differences from baseline to follow-up were assessed by independent *t*-test for normally distributed data or a Mann–Whitney U test for non-normally distributed data. A two-sided *p*-value  $< 0.05$  was considered as statistically significant difference. All analyses were done on data from the modified intention-to-treat population. Missing data occurred at random and were replaced with the mean for the sample at either baseline or follow-up.<sup>21</sup>

## Results

From May 2018 to October 2021, 354 adults with non-obstructive HCM were screened for eligibility and 115 patients met the study inclusion criteria who were randomized into sacubitril/valsartan ( $n = 79$ ) or control ( $n = 36$ ) (Figure 1). Before randomization, 43 patients (37%) were taking ACEi and 17 patients (15%) were taking ARBs. We recruited a typical population of HCM patients from an outpatient HCM clinic. Our sample was diverse with 43% testing genotype-positive for HCM mutation and 10% having implanted cardioverter-defibrillators. No patient was treated via surgical or interventional septal reduction therapy. Baseline demographic characteristics were not significantly different between sacubitril/valsartan and control groups (Table 1).

## Primary outcome

All patients achieved peak level of exertion at baseline and follow-up demonstrated by mean RER of 1.08 both in sacubitril/valsartan and control groups. At 16 weeks, there was no significant change in peak  $\text{VO}_2$  from baseline in the sacubitril/valsartan (15.3 [4.3] vs. 15.9 [4.3] ml/kg/min,  $p = 0.13$ ), or control group (15.2 [3.8] vs. 15.6 [3.9] ml/kg/min,  $p = 0.47$ ) (Table 2).

## Secondary outcomes

There were no significant changes from baseline in anaerobic threshold or peak heart rate in sacubitril/valsartan (9.9 [3.5] vs. 10.7 [3.1] ml/kg/min,  $p = 0.12$ ; 118 [19] vs. 115 [18] bpm,  $p = 0.09$ ) or control group (11.1 [2.9] vs. 11.2 [3.4] ml/kg/min,  $p = 0.90$ ; 120 [18] vs. 118 [16] bpm,  $p = 0.53$ ) (Table 2). There were no

**Table 1** Baseline characteristics of the study population

| Variables                                | Sacubitril/valsartan ( $n = 79$ ) | Control ( $n = 36$ ) |
|------------------------------------------|-----------------------------------|----------------------|
| Demographic characteristics              |                                   |                      |
| Age (years)                              | 59 (10)                           | 57 (12)              |
| Female sex, $n$ (%)                      | 29 (37)                           | 14 (39)              |
| Body weight (kg)                         | 84.8 (17.4)                       | 80.9 (15.1)          |
| BMI ( $\text{kg/m}^2$ )                  | 28.1 (5)                          | 26.8 (4)             |
| NYHA functional class, $n$ (%)           |                                   |                      |
| I                                        | 28 (35)                           | 21 (55)              |
| II                                       | 43 (55)                           | 14 (39)              |
| III                                      | 7 (10)                            | 1 (6)                |
| Systolic blood pressure (mmHg)           | 137 (21)                          | 131 (18)             |
| Diastolic blood pressure (mmHg)          | 82 (11)                           | 77 (10)              |
| LA dimension (mm)                        | 42.6 (5.6)                        | 41.5 (5.4)           |
| LA volume (ml)                           | 42.6 (11.9)                       | 37.7 (10.3)          |
| Atrial fibrillation, $n$ (%)             | 7 (9)                             | 1 (3)                |
| ICD, $n$ (%)                             | 8 (10)                            | 4 (11)               |
| Genetic mutation <sup>a</sup> , $n$ (%)  |                                   |                      |
| MYBPC3                                   | 34 (43)                           | 15 (42)              |
| MYH7                                     | 15 (18)                           | 5 (14)               |
| MYL3                                     | 1 (1)                             | 0 (0)                |
| TNNT2                                    | 12 (15)                           | 6 (17)               |
| TNNI3                                    | 10 (13)                           | 2 (6)                |
| TPM1                                     | 1 (1)                             | 0 (0)                |
| Negative                                 | 26 (33)                           | 9 (25)               |
| N/A                                      | 8 (10)                            | 5 (14)               |
| Medications, $n$ (%)                     |                                   |                      |
| Angiotensin-converting enzyme inhibitors | 5 (6)                             | 2 (6)                |
| Angiotensin receptor blockers            | 2 (3)                             | 1 (3)                |
| Beta-adrenergic blocking agent           | 61 (77)                           | 29 (81)              |
| Calcium channel blocking agent           | 25 (32)                           | 12 (33)              |
| Diuretics                                | 28 (35)                           | 13 (36)              |
| Diabetes                                 | 6 (8)                             | 1 (3)                |
| Anti-inflammatory                        | 32 (41)                           | 13 (36)              |
| Statins                                  | 31 (39)                           | 15 (42)              |
| Anticoagulants                           | 19 (24)                           | 10 (28)              |
| Family history, $n$ (%)                  |                                   |                      |
| Hypertrophic cardiomyopathy              | 14 (18)                           | 5 (14)               |
| Dilated cardiomyopathy                   | 2 (3)                             | 2 (6)                |
| Sudden cardiac death                     | 27 (34)                           | 12 (33)              |
| Comorbidities, $n$ (%)                   |                                   |                      |
| Diabetes                                 | 7 (9)                             | 2 (6)                |
| Thyroid disease                          | 4 (5)                             | 3 (8)                |
| Renal dysfunction                        | 2 (3)                             | 0 (0)                |
| Hepatic dysfunction                      | 1 (1)                             | 2 (6)                |
| Chronic obstructive pulmonary disease    | 4 (5)                             | 2 (6)                |
| Anaemia                                  | 2 (3)                             | 0 (0)                |

Values are given as mean (standard deviation), unless otherwise indicated.

BMI, body mass index; ICD, implantable cardioverter-defibrillator; LA, left atrial; MYBPC3, myosin-binding protein C; MYH,  $\beta$ -myosin heavy chain; MYL, myosin light chain; N/A, not available; NYHA, New York Heart Association; TNNT, troponin T; TNNI, troponin I; TPM, tropomyosin.

<sup>a</sup>Some patients express more than one gene mutation.

**Table 2** Changes in exercise measures from baseline to follow-up

| Variables                        | Sacubitril/valsartan |            |             | Control    |            |              | Between group<br>$\Delta$ (95% CI) | p-value |
|----------------------------------|----------------------|------------|-------------|------------|------------|--------------|------------------------------------|---------|
|                                  | Baseline             | Follow-up  | $\Delta$    | Baseline   | Follow-up  | $\Delta$     |                                    |         |
| Peak VO <sub>2</sub> (ml/kg/min) | 15.3 (4.3)           | 15.9 (4.3) | 0.65 (3.6)  | 15.2 (3.8) | 15.6 (3.9) | 0.46 (3.8)   | 0.19 (−1.28–1.67)                  | 0.80    |
| VO <sub>2</sub> AT (ml/kg/min)   | 9.90 (3.5)           | 10.7 (3.1) | 0.80 (3.7)  | 11.1 (2.9) | 11.2 (3.4) | 0.09 (3.8)   | 0.71 (−1.00–2.43)                  | 0.41    |
| Peak HR (bpm)                    | 118 (19)             | 115 (18)   | −3.41 (17)  | 120 (18)   | 118 (16)   | −1.97 (18.6) | −1.44 (−8.53–5.65)                 | 0.69    |
| Peak VE/VCO <sub>2</sub>         | 30.1 (7.0)           | 28.4 (4.9) | −1.65 (6.4) | 27.6 (3.5) | 29.1 (4.7) | 1.44 (4.8)   | −3.10 (−5.61 to −0.58)             | 0.02    |

Data are given as mean (standard deviation).

AT, anaerobic threshold; CI, confidence interval; HR, heart rate; VE/VCO<sub>2</sub>, minute ventilation to carbon dioxide output slope (ventilatory efficiency); VO<sub>2</sub>, oxygen consumption.

**Table 3** Changes in secondary outcomes from baseline to follow-up

| Variables                       | Sacubitril/valsartan |                |             | Control        |                |              | Between group<br>Δ (95% CI) | p-value |
|---------------------------------|----------------------|----------------|-------------|----------------|----------------|--------------|-----------------------------|---------|
|                                 | Baseline             | Follow-up      | Δ           | Baseline       | Follow-up      | Δ            |                             |         |
| Cardiac structure and function  |                      |                |             |                |                |              |                             |         |
| SBP (mmHg)                      | 137 (21)             | 136 (20)       | −1 (19)     | 131 (18)       | 129 (22)       | −2 (21)      | −1.12 (−9.4–7.2)            | 0.79    |
| DBP (mmHg)                      | 82 (11)              | 80 (12)        | −2 (13)     | 77 (10)        | 79 (12)        | −2 (12)      | 0.44 (−2.0–9.2)             | 0.20    |
| LVOT gradient (rest) (mmHg)     | 12 (14)              | 12 (14)        | 0.08 (11)   | 12 (14)        | 14 (13)        | 2 (6)        | −1.92 (−5.7–1.9)            | 0.191   |
| IVSd (mm)                       | 18.4 (3.5)           | 17.9 (3.2)     | −0.46 (2.2) | 18.7 (4.3)     | 18.5 (3.7)     | −0.22 (2.5)  | −0.25 (−1.2–0.7)            | 0.277   |
| PLWd (mm)                       | 12.9 (3.4)           | 12.7 (2.4)     | −0.15 (2.5) | 11.9 (3.0)     | 11.6 (2.2)     | −0.27 (2.2)  | 0.12 (−0.8–1.1)             | 0.797   |
| E/e'                            | 12.2 (5.2)           | 12.7 (4.7)     | 0.54 (4.8)  | 11.7 (3.9)     | 11.0 (3.4)     | −0.65 (3.2)  | 1.20 (−0.6–3.0)             | 0.143   |
| LVEF (%)                        | 64 (7)               | 64 (6)         | −0.5 (4.5)  | 66 (8)         | 68 (7)         | 2.3 (8)      | −2.79 (−5.2 to −0.4)        | 0.023   |
| LVOT gradient (Valsalva) (mmHg) | 23 (23)              | 21 (20)        | −2.5 (18.6) | 21 (25)        | 19.0 (17)      | −2.05 (17)   | −0.45 (−8.0–7.1)            | 0.41    |
| LAVi (ml)                       | 43 (12)              | 43 (12)        | 0.59 (10)   | 38 (10)        | 36 (8)         | −1.49 (10)   | 2.08 (−2.3–6.5)             | 0.349   |
| Blood biomarkers                |                      |                |             |                |                |              |                             |         |
| CK (μ/L)                        | 120 (64)             | 107 (51)       | −13 (60)    | 128 (53)       | 133 (54)       | 5 (59)       | −18 (−43.0–6.5)             | 0.075   |
| CK-MB (μ/L)                     | 15 (5.5)             | 17 (7)         | 2.23 (5.8)  | 18 (4)         | 17 (5.7)       | −1.33 (6)    | 3.6 (−0.9–8.0)              | 0.131   |
| NT-proBNP (pg/ml)               | 779 (258–1285)       | 986 (300–1204) |             | 951 (568–1428) | 916 (315–1030) |              | 0.02 (−0.2–0.2)*            | 0.841*  |
| Sodium (mmol/L)                 | 140 (2.5)            | 140 (2.7)      | 0.00 (3.30) | 141 (1.8)      | 139 (2.5)      | −1.67 (2.9)  | 1.67 (0.4–2.9)              | 0.015   |
| Potassium (mmol/L)              | 4.6 (0.4)            | 4.5 (0.5)      | −0.01 (0.6) | 4.4 (0.3)      | 4.5 (0.5)      | 0.06 (0.5)   | −0.07 (−0.3–0.1)            | 0.527   |
| AST (μmol/L)                    | 24 (7.7)             | 23 (6.2)       | −1.23 (7.6) | 25 (6.9)       | 25 (8.1)       | 0.13 (6.9)   | −1.36 (−4.5–1.8)            | 0.525   |
| Troponin (ng/L)                 | 28 (27)              | 26 (25)        | −2.51 (27)  | 23 (22)        | (20) (14)      | −2.43 (15)   | −0.08 (−11.5–11.4)          | 0.47    |
| Total protein (g/L)             | 73 (4.6)             | 72 (4.7)       | −1.23 (5.0) | 72 (5.8)       | 73 (4.9)       | 0.32 (5.3)   | −1.55 (−3.6–0.5)            | 0.138   |
| LDL (mmol/L)                    | 3.0 (1.0)            | 2.7 (0.8)      | −0.27 (0.9) | 3.0 (1.0)      | 2.8 (1.0)      | −0.2 (1.1)   | −0.06 (−0.5–0.3)            | 0.752   |
| HDL (mmol/L)                    | 1.2 (0.4)            | 1.2 (0.3)      | −0.01 (0.3) | 1.4 (0.3)      | 1.3 (0.4)      | −0.02 (0.3)  | 0.00 (−0.1–0.1)             | 0.955   |
| Total cholesterol (mmol/L)      | 5.0 (1.3)            | 4.9 (1.1)      | −0.15 (0.9) | 5.0 (1.2)      | 4.8 (1.3)      | −0.19 (1.5)  | 0.04 (−0.5–0.6)             | 0.896   |
| Triglycerides (mmol/L)          | 1.7 (0.8)            | 1.6 (0.6)      | −0.08 (0.8) | 1.4 (0.9)      | 1.4 (0.4)      | −0.07 (0.96) | −0.01 (−0.4–0.3)            | 0.441   |
| Quality of life                 |                      |                |             |                |                |              |                             |         |
| MLHFQ                           | 27 (7–41)            | 22 (8–29)      |             | 27 (19–36)     | 24 (11–35)     |              | −4.88 (−12–3)               | 0.2     |

Values are given as mean (standard deviation, or median (interquartile range)).

AST, aspartate transaminase; CI, confidence interval; CK, creatine kinase; CK-MB, creatine kinase myocardial band; DBP, diastolic blood pressure; E/e', ratio of mitral peak velocity of early filling to early diastolic mitral annular velocity; HDL, high-density lipoprotein; IVSd, interventricular septal diameter; LAVi, left atrial volume index; LDL, low-density lipoprotein; LVEF, left ventricular ejection fraction; LVOT, left ventricular outflow tract; MLHFQ, Minnesota Living with Heart Failure Questionnaire; NT-proBNP, N-terminal pro-B-type natriuretic peptide; PLWd, posterior left ventricular wall diameter; SBP, systolic blood pressure.

<sup>a</sup>Calculated from log transformed data.

clinically relevant changes in other outcome measures including ventilatory efficiency, systolic or diastolic function, cardiac chamber dimensions and thickness, NT-proBNP and quality of life (Table 3).

## Adverse events

Treatment with sacubitril/valsartan was generally well tolerated and there were no observations of hypotension, non-sustained ventricular arrhythmias, atrial fibrillation, or other arrhythmias suggesting that it could be used for relevant indications in patients with HCM. All patients reached the target dose of 97/103 mg.

The use of sacubitril/valsartan was not associated with any serious adverse events. During the study period, three patients in the sacubitril/valsartan group died from sudden cardiac death, ischaemic stroke, and decompensated heart failure due to infective endocarditis, whilst one patient in the control group died from sudden cardiac death.

## Discussion

This is the first study to evaluate the effect of sacubitril/valsartan in patients with non-obstructive HCM. Our results show that

treatment with sacubitril/valsartan for 16 weeks was well tolerated, but had no significant effect on exercise capacity, cardiac structure or function. Analyses of other outcomes of exercise performance and cardiac remodelling revealed no significant differences between baseline and follow-up for either sacubitril/valsartan or control group. Our results are significant as this is the first study to report findings on the effects of angiotensin receptor–neprilysin inhibitor in HCM patients.

Reduced peak  $\text{VO}_2$  has been shown to correlate with clinical decompensation and mortality in HCM.<sup>22,23</sup> As disease progresses, the majority of HCM patients develop dyspnoea on exertion, thus limiting exercise capacity, an important predictor of mortality.<sup>24</sup> Heart failure symptoms are frequently experienced despite normal ejection fraction due to LV outflow obstruction or diastolic dysfunction<sup>25</sup> caused by impaired cardiac energetics which can manifest in genotype-positive individuals before the development of hypertrophy.<sup>26</sup> Whilst pilot studies have shown that sacubitril/valsartan can improve exercise capacity in heart failure patients,<sup>16,27</sup> our results, like several randomized controlled trials including the PARALLAX<sup>28</sup> and ACTIVITY-HF<sup>29</sup> studies, failed to show significant benefits of sacubitril/valsartan over placebo or control in improving exercise capacity.<sup>28–31</sup> One study suggested that patients with limited exercise capacity could exhibit early resistance to treatment, thus improvements may only occur with longer treatment.<sup>29</sup>

The present trial did not find any significant echocardiographic evidence of improvement or deterioration in hypertrophy, LV geometry, systolic or diastolic function in the intervention or control group. Whilst findings from a few trials agree with our results,<sup>32,33</sup> the VANISH trial showed improvement in myocardial structure and function after treatment with ARBs.<sup>7</sup> Improved diastolic function and reduced LV mass was also reported in pilot studies.<sup>34–36</sup> Evidence from pre-clinical<sup>37,38</sup> and clinical studies<sup>7</sup> suggests that calcium channel blocker or ARB treatment may not only attenuate disease progression but might also promote improvement. However, these benefits were seen prior to or in early phase of LV hypertrophy.

Elevated natriuretic peptide concentrations are associated with adverse outcomes in patients with heart failure irrespective of LVEF.<sup>39</sup> The present study did not show any benefit of sacubitril/valsartan administration on NT-proBNP. We report slightly reduced NT-proBNP levels in the control group, which may not be unusual considering that patients were on optimum dose of recommended medication and were managed according to current guidelines. A previous report based on the PARADIGM-HF trial suggested that treatment with sacubitril/valsartan was almost twice as likely as enalapril to reduce NT-proBNP to values below 1000 pg/ml.<sup>39</sup> Furthermore, the PARAMOUNT heart failure preserved ejection fraction study showed that these effects could be seen as early as 4 weeks with benefits sustained for 36 weeks.<sup>11</sup>

Patient-reported assessments using the MLHFQ showed that sacubitril/valsartan did not impact quality of life unlike reports presented previously.<sup>28,40</sup> Patients in the present study had a median quality-of-life score of 27 with a range between 7 and 41 points out of a possible 105 points, which seems to suggest quality

of life is not severely compromised. This is unsurprising considering over 90% of patients were in NYHA class I and II heart failure categories. However, relatively low functional capacity observed in the present study suggests that quality of life may be independent of peak exercise capacity unlike previously reported in heart failure patients.<sup>41</sup>

## Limitations

In the present study, the following limitations should be considered. Due to the COVID-19 pandemic, recruitment was halted, and number of patients recruited into the study was reduced. Interpretation of the main findings should be considered with caution due to the following limitations. Firstly, the study design was not double-blind, placebo-controlled and thus may lead to bias and limit generalizability of findings. Secondly, our initial power calculation suggested the need of 120 patients to be recruited but the study randomized 115 patients and report missing follow-up data for selected variables in 15–20% of cases. Thirdly, drug titration up to maximum dose was at the discretion of each clinical centre, thus modality of medication administration could have impacted some results, e.g. quality of life. Moreover, prior treatment with ACEi or ARB may have affected our results. Fourthly, echocardiography and CPET analyses was not centralized but performed and interpreted by experts at each centre. Lastly, duration of the intervention of 16 weeks may not be sufficient to allow adaptations that may occur at the cellular and molecular level, further leading to changes in clinical and physiological phenotype.

## Conclusions

This is the first study to evaluate the effect of sacubitril/valsartan on clinical phenotype in non-obstructive HCM in humans. Based on the major findings of the present study, it can be concluded that among patients with non-obstructive HCM, sacubitril/valsartan was well tolerated. It did not however lead to a significant improvement in functional capacity, cardiac function and structure.

## Acknowledgements

We would like to thank all study participants and their family members for taking part in this research. We would also like to thank staff members of participating sites who assisted with patient screening and coordination of research visits.

## Funding

This work received funding from the European Union's Horizon 2020 Research and Innovation Programme under Grant Agreement No. 777204. Nduka C. Okwose, Nenad Filipovic and Djordje G. Jakovljevic are also supported by the European Horizon 2020 Research and Innovation Programme under the grant agreement number 952603. The design of the study, data collection, analyses, interpretation of data, and drafting of the manuscript do not reflect the views and opinions of the funders.

**Conflict of interest:** none declared.

## Appendix A

### SILICOFCM Study Centres and Investigators

- Faculty of Medicine, University of Novi Sad, Novi Sad: Aleksandar Redzek, Marija Bjelobrk, Aleksandra Ilic, Miodrag Golubovic, Tatjana Miljkovic
- Cardiology Department, Faculties of Medicine and Pharmacy, University of Belgrade, Serbia: Ruzica Maksimovic, Stefan Seman, Gorica Ristic, Vidan Markovic, Ratko Lasica.
- Cardiovascular Research, Translational and Clinical Research Institute, Newcastle University, UK: Paul Brennan, Kristian Bailey, Christopher Eggett, Amy Fuller, Alaa Alyahya.
- Department of Internal Medicine II, University Medical Centre Regensburg, Germany: Stefan Wagner, Andrea Baessler, Bernhard Unsoeld.
- Careggi University Hospital, University of Florence, Florence, Italy: Alessia Tomberli.

[Correction added on 11 June 2024, after first online publication: The appendix section containing the name of the investigators has been added in this version]

## References

- Marian AJ, Braunwald E. Hypertrophic cardiomyopathy: Genetics, pathogenesis, clinical manifestations, diagnosis, and therapy. *Circ Res* 2021;121:749–770. <https://doi.org/10.1161/CIRCRESAHA.117.311059>
- Semsarian C, Ingles J, Maron MS, Maron BJ. New perspectives on the prevalence of hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2015;65:1249–1254. <https://doi.org/10.1016/j.jacc.2015.01.019>
- Garfinkel AC, Seidman JG, Seidman CE. Genetic pathogenesis of hypertrophic and dilated cardiomyopathy. *Heart Fail Clin* 2018;14:139–146. <https://doi.org/10.1016/j.hfc.2019.12.004>
- Seferović PM, Polovina M, Bauersachs J, Arad M, Gal TB, Lund LH, et al. Heart failure in cardiomyopathies: A position paper from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2019;21:553–576. <https://doi.org/10.1002/ehf.1461>
- Maron BJ, Rowin EJ, Maron MS. Hypertrophic cardiomyopathy: New concepts and therapies. *Annu Rev Med* 2022;73:363–375. <https://doi.org/10.1146/annurev-med-042220-021539>
- Ommen SR, Mital S, Burke MA, Day SM, Deswal A, Elliott P, et al. 2020 AHA/ACC Guideline for the diagnosis and treatment of patients with hypertrophic cardiomyopathy: A report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2020;142:e558–e631. <https://doi.org/10.1161/CIR.0000000000000937>
- Ho CY, Day SM, Axelsson A, Russell MW, Zahka K, Lever HM, et al. Valsartan in early-stage hypertrophic cardiomyopathy: A randomized phase 2 trial. *Nat Med* 2021;27:1818–1824. <https://doi.org/10.1038/s41591-021-01505-4>
- Zamorano JL, Anastakis A, Borger MA, Borggrefe M, Cecchi F, Charron P, et al. 2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy: The Task Force for the diagnosis and management of hypertrophic cardiomyopathy of the European Society of Cardiology (ESC). *Eur Heart J* 2014;35:2733–2779. <https://doi.org/10.1093/eurheartj/ehu284>
- Ho CY, Meallif ME, Bach RG, Bhattacharya M, Choudhury L, Edelberg JM, et al. Evaluation of mavacamten in symptomatic patients with nonobstructive hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2020;75:2649–2660. <https://doi.org/10.1016/j.jacc.2020.03.064>
- McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, et al.; PARADIGM-HF Investigators and Committees. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014;371:993–1004. <https://doi.org/10.1056/NEJMoa1409077>
- Solomon SD, Zile M, Pieske B, Voors A, Shah A, Kraigher-Krainer E, et al.; Prospective comparison of ARNI with ARB on Management Of heart failure with preserved ejection fraction (PARAMOUNT) Investigators. The angiotensin receptor neprilysin inhibitor LCZ696 in heart failure with preserved ejection fraction: A phase 2 double-blind randomised controlled trial. *Lancet* 2012;380:1387–1395. [https://doi.org/10.1016/S0140-6736\(12\)61227-6](https://doi.org/10.1016/S0140-6736(12)61227-6)
- Armentaro G, D'Arrigo G, Magurno M, Toscani AF, Condoleo V, Miceli S, et al. Impact of sacubitril/valsartan on clinical and echocardiographic parameters in heart failure patients with reduced ejection fraction: Data from a real life 2-year follow-up study. *Front Pharmacol* 2021;12:733475. <https://doi.org/10.3389/fphar.2021.733475>
- Sun Y, Song S, Zhang Y, Mo W, Zhang X, Wang N, et al. Effect of angiotensin receptor neprilysin inhibitors on left atrial remodeling and prognosis in heart failure. *ESC Heart Fail* 2022;9:667–675. <https://doi.org/10.1002/ehf2.13691>
- Landolfo M, Piani F, Esposti DD, Cosentino E, Bacchelli S, Dormi A, et al. Effects of sacubitril valsartan on clinical and echocardiographic parameters of outpatients with heart failure and reduced ejection fraction. *Int J Cardiol Heart Vasc* 2020;31:100656. <https://doi.org/10.1016/j.ijcha.2020.100656>
- Biering-Sorensen T, Shah A, Claggett B, Zile M, Pieske B, Pieske-Kraigher E, et al. The angiotensin receptor neprilysin inhibitor (ARNI), sacubitril/valsartan, improves left ventricular myocardial deformation in heart failure with preserved ejection fraction (PARAMOUNT trial). *J Am Coll Cardiol* 2018;71:A2665 (abstr).
- Beltrán P, Palau P, Domínguez E, Faraudo M, Núñez E, Guri O, et al. Sacubitril/valsartan and short-term changes in the 6-minute walk test: A pilot study. *Int J Cardiol* 2018;252:136–139. <https://doi.org/10.1016/j.ijcard.2019.10.074>
- Tafelmeier M, Baessler A, Wagner S, Unsoeld B, Preveden A, Barlocco F, et al. Design of the SILICOFCM study: Effect of sacubitril/valsartan vs lifestyle intervention on functional capacity in patients with hypertrophic cardiomyopathy. *Clin Cardiol* 2020;43:430–440. <https://doi.org/10.1002/clc.23346>
- Lang RM, Badano LP, Victor MA, Afilalo J, Armstrong A, Ernande L, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: An update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 2015;28:1–39.e14. <https://doi.org/10.1016/j.echo.2014.10.003>
- Guazzi M, Arena R, Halle M, Piepoli MF, Myers J, Lavie CJ. 2016 Focused update: Clinical recommendations for cardiopulmonary exercise testing data assessment in specific patient populations. *Circulation* 2016;133:e694–e711. <https://doi.org/10.1161/CIR.0000000000000406>
- Saberi S, Wheeler M, Bragg-Gresham J, Hornsby W, Agarwal PP, Attali A, et al. Effect of moderate-intensity exercise training on peak oxygen consumption in patients with hypertrophic cardiomyopathy: A randomized clinical trial. *JAMA* 2021;325:1349–1357. <https://doi.org/10.1001/jama.2021.2503>
- Xu Z. Martin Bland (2015): An introduction to medical statistics. *Stat Papers* 2021;58:953–954. <https://doi.org/10.1007/s00362-017-0925-5>
- Coats CJ, Rantell K, Bartnik A, Patel A, Mist B, McKenna WJ, et al. Cardiopulmonary exercise testing and prognosis in hypertrophic cardiomyopathy. *Circ Heart Fail* 2015;8:1022–1031. <https://doi.org/10.1161/CIRCHEARTFAILURE.114.002248>
- Finocchiaro G, Haddad F, Knowles JW, Caleshu C, Pavlovic A, Homburger J, et al. Cardiopulmonary responses and prognosis in hypertrophic cardiomyopathy: A potential role for comprehensive noninvasive hemodynamic assessment. *JACC Heart Fail* 2015;3:408–418. <https://doi.org/10.1016/j.jchf.2014.11.011>
- Desai MY, Bhonsale A, Patel P, Naji P, Smedira NG, Thamilarasan M, et al. Exercise echocardiography in asymptomatic HCM: Exercise capacity, and not LV outflow tract gradient predicts long-term outcomes. *JACC Cardiovasc Imaging* 2014;7:26–36. <https://doi.org/10.1016/j.jcmg.2013.08.010>
- Matsumura Y, Elliott PM, Virdee MS, Sorajja P, Doy Y, McKenna WJ. Left ventricular diastolic function assessed using Doppler tissue imaging in patients with hypertrophic cardiomyopathy: Relation to symptoms and exercise capacity. *Heart* 2002;87:247–251. <https://doi.org/10.1136/heart.87.3.247>
- Ho CY, Sweitzer NK, McDonough B, Maron BJ, Casey SA, Seidman JG, et al. Assessment of diastolic function with Doppler tissue imaging to predict genotype in preclinical hypertrophic cardiomyopathy. *Circulation* 2002;105:2992–2997. <https://doi.org/10.1161/01.cir.0000019070.70491.6d>
- Malfatto G, Ravaro S, Caravita S, Baratto C, Sorropago A, Giglio A, et al. Improvement of functional capacity in sacubitril-valsartan treated patients assessed by cardiopulmonary exercise test. *Acta Cardiol* 2020;75:732–736. <https://doi.org/10.1080/00015385.2019.1669317>
- Pieske B, Wachter R, Shah SJ, Baldrige A, Szczepiety P, Ibram G, et al.; PARALLAX Investigators and Committee members. Effect of sacubitril/valsartan vs standard medical therapies on plasma NT-proBNP concentration and submaximal exercise capacity in patients with heart failure and preserved ejection fraction: The PARALLAX randomized clinical trial. *JAMA* 2021;326:1919–1929. <https://doi.org/10.1001/jama.2021.18463>
- Halle M, Schöbel C, Winzer EB, Bernhardt P, Mueller S, Sieder C, et al. A randomized clinical trial on the short-term effects of 12-week sacubitril/valsartan

- vs. enalapril on peak oxygen consumption in patients with heart failure with reduced ejection fraction: Results from the ACTIVITY-HF study. *Eur J Heart Fail* 2021;**23**:2073–2082. <https://doi.org/10.1002/ehf.2355>
30. Dos Santos MR, de Alves MJ, Jordão CP, CEN P, KTS C, de Souza FR, et al. Sacubitril/valsartan versus enalapril on exercise capacity in patients with heart failure with reduced ejection fraction: A randomized, double-blind, active-controlled study. *Am Heart J* 2021;**239**:1–10. <https://doi.org/10.1016/j.ahj.2021.05.005>
  31. Lau CW, Martens P, Lambeets S, Dupont M, Mullens W. Effects of sacubitril/valsartan on functional status and exercise capacity in real-world patients. *Acta Cardiol* 2019;**74**:405–412. <https://doi.org/10.1080/00015385.2018.1521054>
  32. Axelsson A, Iversen K, Vejlsstrup N, Ho CY, Havndrup O, Kofoed KF, et al. Functional effects of losartan in hypertrophic cardiomyopathy – a randomised clinical trial. *Heart* 2016;**102**:285–291. <https://doi.org/10.1136/heartjnl-2015-308343>
  33. Shimada YJ, Passeri JJ, Baggish AL, O'Callaghan C, Lowry PA, Yannekis G, et al. Effects of losartan on left ventricular hypertrophy and fibrosis in patients with nonobstructive hypertrophic cardiomyopathy. *JACC Heart Fail* 2013;**1**:480–487. <https://doi.org/10.1016/j.jchf.2013.09.001>
  34. Araujo AQ, Arteaga E, Ianni BM, Buck PC, Rabello R, Mady C. Effect of losartan on left ventricular diastolic function in patients with nonobstructive hypertrophic cardiomyopathy. *Am J Cardiol* 2005;**96**:1563–1567. <https://doi.org/10.1016/j.amjcard.2005.07.065>
  35. Yamazaki T, Suzuki JI, Shimamoto R, Tsuji T, Ohmoto-Sekine Y, Ohtomo K, et al. A new therapeutic strategy for hypertrophic nonobstructive cardiomyopathy in humans. A randomized and prospective study with an angiotensin II receptor blocker. *Int Heart J* 2007;**48**:715–724. <https://doi.org/10.1536/ihj.48.715>
  36. Penicka M, Gregor P, Kerekes R, Marek D, Curila K, Krupicka J; Candesaratan use in Hypertrophic And Non-obstructive Cardiomyopathy Estate (CHANCE) Study Investigators. The effects of candesartan on left ventricular hypertrophy and function in nonobstructive hypertrophic cardiomyopathy: A pilot, randomized study. *J Mol Diagn* 2009;**11**:35–41. <https://doi.org/10.2353/jmoldx.2009.080082>
  37. Semsarian C, Ahmad I, Giewat M, Georgakopoulos D, Schmitt JP, McConnell BK, et al. The L-type calcium channel inhibitor diltiazem prevents cardiomyopathy in a mouse model. *J Clin Invest* 2002;**109**:1013–1020. <https://doi.org/10.1172/JCI14677>
  38. Teekakirikul P, Eminaga S, Toka O, Alcalai R, Wang L, Wakimoto H, et al. Cardiac fibrosis in mice with hypertrophic cardiomyopathy is mediated by non-myocyte proliferation and requires Tgf- $\beta$ . *J Clin Invest* 2010;**120**:3520–3529. <https://doi.org/10.1172/JCI42028>
  39. Zile MR, Claggett BL, Prescott MF, McMurray JJV, Packer M, Rouleau JL, et al. Prognostic implications of changes in N-terminal pro-B-type natriuretic peptide in patients with heart failure. *J Am Coll Cardiol* 2016;**68**:2425–2436. <https://doi.org/10.1016/j.jacc.2016.09.931>
  40. Solomon SD, McMurray JJV, Anand IS, Ge J, Lam CSP, Maggioni AP, et al.; PARAGON-HF Investigators and Committees. Angiotensin-neprilysin inhibition in heart failure with preserved ejection fraction. *N Engl J Med* 2019;**381**:1609–1620. <https://doi.org/10.1056/NEJMoa1908655>
  41. Fatrin S, Okwose NC, Bailey K, Velicki L, Popovic D, Ristic A, et al. Haemodynamic determinants of quality of life in chronic heart failure. *BMC Cardiovasc Disord* 2022;**22**:412. <https://doi.org/10.1186/s12872-022-02829-w>